

- (10) 1. Find an equation for the function $f(x)$ if $f''(x) = 6x - 4$ and $f(1) = 1$, $f'(1) = 2$.

A4


- (15) 2. A differentiable function $y = f(x)$ has the following properties:
- (1) The domain is $(-\infty, \infty)$.
 - (2) The only x -intercept is $x = 3$.
 - (3) $\lim_{x \rightarrow \infty} f(x) = -\infty$ and $\lim_{x \rightarrow -\infty} f(x) = 0$.
 - (4) $f(x)$ is negative for $x > 3$ and $f(x)$ is positive for $x < 3$. Moreover, $f(4) = -9$, $f(1) = 2$, and $f(2) = 3$.
 - (5) $f'(x)$ is 0 only at $x = 2$. Moreover, $f'(-1)$ is positive while $f'(7)$ is negative.
 - (6) $f''(x)$ is 0 only at $x = 1$. Moreover, $f''(-3)$ is positive while $f''(7)$ is negative.
- (a) Determine the intervals where the function is increasing and the intervals where it is decreasing.
- (b) Determine the x -values where the function has a local maximum and those where it has a local minimum.
- (c) Determine the intervals where the function is concave up and the intervals where it is concave down.

Continued on next page.

(d) Determine the x -values where the inflection points occur.

(e) Determine all vertical and horizontal asymptotes.

(f) Sketch the graph of a function which has the properties found in parts (a)-(e) above. Show on the graph all intercepts, asymptotes, local maxima and local minima, and points of inflection.



A6

(20) 3. Compute $\frac{dy}{dx}$ in each of the following cases. You don't have to simplify.

(a) $y = \sqrt{6 + \frac{8}{x^5}}$

(b) $y = x^4 \tan^{-1}(9 - 7x)$

(c) $y = \frac{\sin(5 \ln x)}{4 + e^{7x}}$

(d) $y = \int_2^x \frac{t^5}{2 + t^6} dt$

(12) 4. Consider the function $f(x) = e^{-2x}$.

(a) Find an equation of the line tangent to the graph of $y = f(x)$ at the point $P = (1, e^{-2})$.

(b) Use linear approximation to get an estimate for the value of $f(1.2)$.

(c) Use the second derivative to determine whether your estimate for (b) is likely to be high or low.

A8

(12) 5. Consider the equation $x^3 - 7x + 1 = 0$.

(a) Does this equation have a solution in the interval $[-3, -2]$? Justify your answer.

(b) Compute, using Newton's method, the second approximation of a solution by starting with the first approximation $x = -2$. You do not need to simplify your final numerical answer.


(10) 6. Using the limit definition of the derivative, find the derivative of $f(x)$ at $x = 2$ where

$$f(x) = x^2 - x - 1.$$

A10

- (10) 7. Find an equation of the line tangent to the graph of the function given implicitly by the equation $2x^2y - 3xy^2 = -14$ at the point $(2, -1)$.

- (10) 8. An open-top box is to be made by cutting small congruent squares from the corners of a $9\text{ cm} \times 9\text{ cm}$ sheet of metal and bending up the sides. What is largest possible volume of such a box?



A12

(20) 9. Compute the following limits.

(a) $\lim_{x \rightarrow 2^-} \frac{x^2 - 4}{|x - 2|}$

(b) $\lim_{x \rightarrow 2} \frac{x^2 - 4x + 5}{x^2 - 2}$

(c) $\lim_{x \rightarrow \infty} \frac{\sin(x)}{1 + x^2}$

(d) $\lim_{x \rightarrow 0} \frac{e^x + e^{-x} - 2}{e^x - 1 - x}$

- (12) 10. Find the global maximum and minimum values of the function

$$f(x) = x^3 - 12x$$

on the interval $[0, 4]$. You don't have to sketch the graph but you must justify your answer.

Global maximum of f : _____

Global minimum of f : _____

A14

(15) 11. Evaluate the following integrals.

(a) $\int_0^1 x \sin(x^2 + 5) dx$


(b) $\int \left(\sqrt{x} - \frac{2}{\sqrt{x}} \right)^2 dx$

(c) $\int_1^2 \frac{2x^2 - 2}{x} dx$

- (10) 12. A car travels on a restricted-access highway which is 100 miles long. The speed limit on the road is 60 miles per hour. Suppose that $s(t)$, the car's distance from the beginning of the road, is a differentiable function of the time t . If the car enters the road at 9am and leaves the road at 10:15am at the other end of the highway, one and a quarter hours later, explain why the driver should expect to receive a speeding citation (ticket). Your explanation should be in complete English sentences. You should mention a specific result from this course and explain precisely why this result shows that the ticket is valid.

A16

- (12) 13. Let $f(x)$ be a function defined over the interval $[0,4]$, whose graph is shown as below. The graph of f over $[1,3]$ is a semi-circle centered at $(2, 0)$ and of radius 1, the other pieces are straight lines.



- (a) Find $\int_0^4 f(x) dx$ geometrically.

- (b) Compute the Riemann sum for $f(x)$ where $[0, 4]$ is subdivided into 4 subintervals of equal size and the sample points are the right endpoints.

- (c) Let $F(x) = \int_0^x f(t) dt$. Find $F'(2)$ and $F'(3)$.


- (10) 14. Determine the values of the parameters a and b such that the following function $f(x)$ becomes continuous and differentiable at $x = -2$:

$$f(x) = \begin{cases} x^2 + 2x + b & \text{for } x > -2 \\ ax & \text{for } x \leq -2 \end{cases}$$

A18

- (12) 15. Consider the finite region A bounded by the “wedge” $y = |x|$ and the parabola $y = \frac{1}{2}x^2 + \frac{1}{2}$.

(a) Draw a sketch of the region A . [*Hint:* The parabola fits “snuggly” into the wedge.]



(b) Express the area of A in terms of one or more definite integrals.

(c) Compute the area of the region A as a sum of fractions (you do not need to simplify the answer).

- (10) 16. The minute hand of a large tower clock is 2 m long. At each full hour, the tip T of the hand points to the center C of the numeral XII. How fast is the distance between T and C changing when it is 7 : 10 am? [*Hint:* Sketch the triangle formed by C , T , and the center of the clock. Observe that it is isosceles.]

A20

More space for your work.

A2

More space for your work.

- (10) 1. Find an equation for the function $f(x)$ if $f''(x) = 6x + 4$ and $f(2) = 1$, $f'(2) = -1$.

B4


- (15) 2. A differentiable function $y = f(x)$ has the following properties:
- (1) The domain is $(-\infty, \infty)$.
 - (2) The only x -intercept is $x = 0$.
 - (3) $\lim_{x \rightarrow \infty} f(x) = -\infty$ and $\lim_{x \rightarrow -\infty} f(x) = 0$.
 - (4) $f(x)$ is negative for $x > 0$ and $f(x)$ is positive for $x < 0$. Moreover, $f(1) = -9$, $f(-2) = 2$, and $f(-1) = 3$.
 - (5) $f'(x)$ is 0 only at $x = -1$. Moreover, $f'(-4)$ is positive while $f'(4)$ is negative.
 - (6) $f''(x)$ is 0 only at $x = -2$. Moreover, $f''(-6)$ is positive while $f''(4)$ is negative.
- (a) Determine the intervals where the function is increasing and the intervals where it is decreasing.
- (b) Determine the x -values where the function has a local maximum and those where it has a local minimum.
- (c) Determine the intervals where the function is concave up and the intervals where it is concave down.

Continued on next page.

(d) Determine the x -values where the inflection points occur.

(e) Determine all vertical and horizontal asymptotes.

(f) Sketch the graph of a function which has the properties found in parts (a)-(e) above. Show on the graph all intercepts, asymptotes, local maxima and local minima, and points of inflection.



B6

(20) 3. Compute $\frac{dy}{dx}$ in each of the following cases. You don't have to simplify.

(a) $y = \sqrt{3 + \frac{5}{x^9}}$

(b) $y = x^5 \tan^{-1}(8 - 7x)$

(c) $y = \frac{\cos(5 \ln x)}{7 + e^{8x}}$

(d) $y = \int_6^x \frac{t^3}{8 + t^4} dt$

(12) 4. Consider the function $f(x) = e^{3x}$.

(a) Find an equation of the line tangent to the graph of $y = f(x)$ at the point $P = (1, e^3)$.

(b) Use linear approximation to get an estimate for the value of $f(1.1)$.

(c) Use the second derivative to determine whether your estimate for (b) is likely to be high or low.

B8

(12) 5. Consider the equation $x^3 - 7x + 1 = 0$.

(a) Does this equation have a solution in the interval $[-3, -1]$? Justify your answer.

(b) Compute, using Newton's method, the second approximation of a solution by starting with the first approximation $x = -3$. You do not need to simplify your final numerical answer.


(10) 6. Using the limit definition of the derivative, find the derivative of $f(x)$ at $x = 2$ where

$$f(x) = x^2 - 2x + 1.$$

B10

- (10) 7. Find an equation of the line tangent to the graph of the function given implicitly by the equation $2x^2y - 3xy^2 = 14$ at the point $(-2, 1)$.

- (10) 8. An open-top box is to be made by cutting small congruent squares from the corners of a $6\text{ cm} \times 6\text{ cm}$ sheet of metal and bending up the sides. What is largest possible volume of such a box?



B12

(20) 9. Compute the following limits.

(a) $\lim_{x \rightarrow 2^+} \frac{x^2 - 4}{|x - 2|}$

(b) $\lim_{x \rightarrow 1} \frac{x^2 - 4x + 5}{x^2 - 2}$

(c) $\lim_{x \rightarrow \infty} \frac{\cos(x)}{1 + x^2}$

(d) $\lim_{x \rightarrow 0} \frac{\cos(x) - 1}{e^x + e^{-x} - 2}$

- (12) 10. Find the global maximum and minimum values of the function

$$f(x) = x^3 - 12x$$

on the interval $[-1, 3]$. You don't have to sketch the graph but you must justify your answer.

Global maximum of f : _____

Global minimum of f : _____

B14

(15) 11. Evaluate the following integrals.

(a) $\int_0^1 \frac{e^x}{\sqrt{1+e^x}} dx$


(b) $\int \left(\sqrt{x} + \frac{3}{\sqrt{x}} \right)^2 dx$

(c) $\int_1^2 \frac{2x^2 - 1}{x} dx$

- (10) 12. A car travels on a restricted-access highway which is 100 miles long. The speed limit on the road is 60 miles per hour. Suppose that $s(t)$, the car's distance from the beginning of the road, is a differentiable function of the time t . If the car enters the road at 9am and leaves the road at 10:15am at the other end of the highway, one and a quarter hours later, explain why the driver should expect to receive a speeding citation (ticket). Your explanation should be in complete English sentences. You should mention a specific result from this course and explain precisely why this result shows that the ticket is valid.

B16

- (12) 13. Let $f(x)$ be a function defined over the interval $[0,4]$, whose graph is shown as below. The graph of f over $[1,3]$ is a semi-circle centered at $(2, 0)$ and of radius 1, the other pieces are straight lines.



- (a) Find $\int_0^4 f(x) dx$ geometrically.

- (b) Compute the Riemann sum for $f(x)$ where $[0, 4]$ is subdivided into 4 subintervals of equal size and the sample points are the right endpoints.

- (c) Let $F(x) = \int_0^x f(t) dt$. Find $F'(2)$ and $F'(3)$.


- (10) 14. Determine the values of the parameters a and b such that the following function $f(x)$ becomes continuous and differentiable at $x = -1$:

$$f(x) = \begin{cases} x^2 + 3x + b & \text{for } x > -1 \\ ax & \text{for } x \leq -1 \end{cases}$$

B18

- (12) 15. Consider the finite region A bounded by the “wedge” $y = |x|$ and the parabola $y = \frac{1}{4}x^2 + 1$.

(a) Draw a sketch of the region A . [*Hint:* The parabola fits “snuggly” into the wedge.]



(b) Express the area of A in terms of one or more definite integrals.

(c) Compute the area of the region A as a sum of fractions (you do not need to simplify the answer).

- (10) 16. The minute hand of a large tower clock is 2 m long. At each full hour, the tip T of the hand points to the center C of the numeral XII. How fast is the distance between T and C changing when it is 9 : 20 pm? [*Hint*: Sketch the triangle formed by C , T , and the center of the clock. Observe that it is isosceles.]

B20

More space for your work.

B2

More space for your work.

- (10) 1. Find an equation for the function $f(x)$ if $f''(x) = -6x + 4$ and $f(3) = 1$, $f'(3) = -2$.

C4


- (15) 2. A differentiable function $y = f(x)$ has the following properties:
- (1) The domain is $(-\infty, \infty)$.
 - (2) The only x -intercept is $x = 1$.
 - (3) $\lim_{x \rightarrow \infty} f(x) = -\infty$ and $\lim_{x \rightarrow -\infty} f(x) = 0$.
 - (4) $f(x)$ is negative for $x > 1$ and $f(x)$ is positive for $x < 1$. Moreover, $f(2) = -9$, $f(-1) = 2$, and $f(0) = 3$.
 - (5) $f'(x)$ is 0 only at $x = 0$. Moreover, $f'(-3)$ is positive while $f'(5)$ is negative.
 - (6) $f''(x)$ is 0 only at $x = -1$. Moreover, $f''(-5)$ is positive while $f''(5)$ is negative.
- (a) Determine the intervals where the function is increasing and the intervals where it is decreasing.
- (b) Determine the x -values where the function has a local maximum and those where it has a local minimum.
- (c) Determine the intervals where the function is concave up and the intervals where it is concave down.

Continued on next page.

(d) Determine the x -values where the inflection points occur.

(e) Determine all vertical and horizontal asymptotes.

(f) Sketch the graph of a function which has the properties found in parts (a)-(e) above. Show on the graph all intercepts, asymptotes, local maxima and local minima, and points of inflection.



C6

(20) 3. Compute $\frac{dy}{dx}$ in each of the following cases. You don't have to simplify.

(a) $y = \sqrt{2 + \frac{3}{x^7}}$

(b) $y = x^2 \tan^{-1}(2 - 3x)$

(c) $y = \frac{\sin(3 \ln x)}{5 + e^{4x}}$

(d) $y = \int_3^x \frac{t^3}{2 + t^4} dt$

(12) 4. Consider the function $f(x) = e^{-3x}$.

(a) Find an equation of the line tangent to the graph of $y = f(x)$ at the point $P = (1, e^{-3})$.

(b) Use linear approximation to get an estimate for the value of $f(1.2)$.

(c) Use the second derivative to determine whether your estimate for (b) is likely to be high or low.

C8

(12) 5. Consider the equation $x^3 - 7x + 1 = 0$.

(a) Does this equation have a solution in the interval $[1, 3]$? Justify your answer.

(b) Compute, using Newton's method, the second approximation of a solution by starting with the first approximation $x = 3$. You do not need to simplify your final numerical answer.


(10) 6. Using the limit definition of the derivative, find the derivative of $f(x)$ at $x = 2$ where

$$f(x) = x^2 + x - 1.$$

C10

- (10) 7. Find an equation of the line tangent to the graph of the function given implicitly by the equation $2x^2y - 3xy^2 = -16$ at the point $(1, -2)$.

- (10) 8. An open-top box is to be made by cutting small congruent squares from the corners of a $3\text{ cm} \times 3\text{ cm}$ sheet of metal and bending up the sides. What is largest possible volume of such a box?



C12

(20) 9. Compute the following limits.

(a) $\lim_{x \rightarrow -2^+} \frac{x^2 - 4}{|x + 2|}$

(b) $\lim_{x \rightarrow -1} \frac{x^2 - 4x + 5}{x^2 - 2}$

(c) $\lim_{x \rightarrow \infty} \frac{\cos(x)}{e^x}$

(d) $\lim_{x \rightarrow 0} \frac{\cos(x) - 1}{e^x - 1 - x}$

- (12) 10. Find the global maximum and minimum values of the function

$$f(x) = x^3 - 12x$$

on the interval $[0, 3]$. You don't have to sketch the graph but you must justify your answer.

Global maximum of f : _____

Global minimum of f : _____

C14

(15) 11. Evaluate the following integrals.

(a) $\int_1^2 \frac{(\ln(x+1))^2}{x+1} dx$


(b) $\int \left(\sqrt{x} - \frac{1}{\sqrt{x}} \right)^2 dx$

(c) $\int_1^2 \frac{2x^2 - 4}{x} dx$

- (10) 12. A car travels on a restricted-access highway which is 100 miles long. The speed limit on the road is 60 miles per hour. Suppose that $s(t)$, the car's distance from the beginning of the road, is a differentiable function of the time t . If the car enters the road at 9am and leaves the road at 10:15am at the other end of the highway, one and a quarter hours later, explain why the driver should expect to receive a speeding citation (ticket). Your explanation should be in complete English sentences. You should mention a specific result from this course and explain precisely why this result shows that the ticket is valid.

C16

- (12) 13. Let $f(x)$ be a function defined over the interval $[0,4]$, whose graph is shown as below. The graph of f over $[1,3]$ is a semi-circle centered at $(2, 0)$ and of radius 1, the other pieces are straight lines.



- (a) Find $\int_0^4 f(x) dx$ geometrically.

- (b) Compute the Riemann sum for $f(x)$ where $[0, 4]$ is subdivided into 4 subintervals of equal size and the sample points are the right endpoints.

- (c) Let $F(x) = \int_0^x f(t) dt$. Find $F'(2)$ and $F'(3)$.


- (10) 14. Determine the values of the parameters a and b such that the following function $f(x)$ becomes continuous and differentiable at $x = 1$:

$$f(x) = \begin{cases} x^2 - 3x + b & \text{for } x > 1 \\ ax & \text{for } x \leq 1 \end{cases}$$

C18

- (12) 15. Consider the finite region A bounded by the “wedge” $y = |x|$ and the parabola $y = x^2 + \frac{1}{4}$.

(a) Draw a sketch of the region A . [*Hint:* The parabola fits “snuggly” into the wedge.]



(b) Express the area of A in terms of one or more definite integrals.

(c) Compute the area of the region A as a sum of fractions (you do not need to simplify the answer).

- (10) 16. The minute hand of a large tower clock is 1 m long. At each full hour, the tip T of the hand points to the center C of the numeral XII. How fast is the distance between T and C changing when it is 11 : 10 pm? [*Hint*: Sketch the triangle formed by C , T , and the center of the clock. Observe that it is isosceles.]

C20

More space for your work.

C2

More space for your work.

- (10) 1. Find an equation for the function $f(x)$ if $f''(x) = -6x + 2$ and $f(4) = 1$, $f'(4) = 1$.

D4


- (15) 2. A differentiable function $y = f(x)$ has the following properties:
- (1) The domain is $(-\infty, \infty)$.
 - (2) The only x -intercept is $x = 2$.
 - (3) $\lim_{x \rightarrow \infty} f(x) = -\infty$ and $\lim_{x \rightarrow -\infty} f(x) = 0$.
 - (4) $f(x)$ is negative for $x > 2$ and $f(x)$ is positive for $x < 2$. Moreover, $f(3) = -9$, $f(0) = 2$, and $f(1) = 3$.
 - (5) $f'(x)$ is 0 only at $x = 1$. Moreover, $f'(-2)$ is positive while $f'(6)$ is negative.
 - (6) $f''(x)$ is 0 only at $x = 0$. Moreover, $f''(-4)$ is positive while $f''(6)$ is negative.
- (a) Determine the intervals where the function is increasing and the intervals where it is decreasing.
- (b) Determine the x -values where the function has a local maximum and those where it has a local minimum.
- (c) Determine the intervals where the function is concave up and the intervals where it is concave down.

Continued on next page.

(d) Determine the x -values where the inflection points occur.

(e) Determine all vertical and horizontal asymptotes.

(f) Sketch the graph of a function which has the properties found in parts (a)-(e) above. Show on the graph all intercepts, asymptotes, local maxima and local minima, and points of inflection.



D6

(20) 3. Compute $\frac{dy}{dx}$ in each of the following cases. You don't have to simplify.

(a) $y = \sqrt{5 + \frac{2}{x^6}}$

(b) $y = x^3 \tan^{-1}(4 - 6x)$

(c) $y = \frac{\cos(7 \ln x)}{8 + e^{6x}}$

(d) $y = \int_2^x \frac{t^5}{7 + t^8} dt$

(12) 4. Consider the function $f(x) = e^{2x}$.

(a) Find an equation of the line tangent to the graph of $y = f(x)$ at the point $P = (1, e^2)$.

(b) Use linear approximation to get an estimate for the value of $f(1.1)$.

(c) Use the second derivative to determine whether your estimate for (b) is likely to be high or low.

D8

(12) 5. Consider the equation $x^3 - 7x + 1 = 0$.

(a) Does this equation have a solution in the interval $[2, 3]$? Justify your answer.

(b) Compute, using Newton's method, the second approximation of a solution by starting with the first approximation $x = 2$. You do not need to simplify your final numerical answer.


(10) 6. Using the limit definition of the derivative, find the derivative of $f(x)$ at $x = 2$ where

$$f(x) = x^2 + 2x + 1.$$

D10

- (10) 7. Find an equation of the line tangent to the graph of the function given implicitly by the equation $2x^2y - 3xy^2 = 16$ at the point $(-1, 2)$.

- (10) 8. An open-top box is to be made by cutting small congruent squares from the corners of a $12\text{ cm} \times 12\text{ cm}$ sheet of metal and bending up the sides. What is largest possible volume of such a box?



D12

(20) 9. Compute the following limits.

(a) $\lim_{x \rightarrow -2^-} \frac{x^2 - 4}{|x + 2|}$

(b) $\lim_{x \rightarrow -2} \frac{x^2 - 4x + 5}{x^2 - 2}$

(c) $\lim_{x \rightarrow \infty} \frac{\sin(x^2)}{\sqrt{x}}$

(d) $\lim_{x \rightarrow 0} \frac{e^x - 1 - x}{\cos(x) - 1}$

- (12) 10. Find the global maximum and minimum values of the function

$$f(x) = x^3 - 12x$$

on the interval $[1, 3]$. You don't have to sketch the graph but you must justify your answer.

Global maximum of f : _____

Global minimum of f : _____

D14

(15) 11. Evaluate the following integrals.

(a) $\int_0^1 x e^{-x^2+3} dx$


(b) $\int \left(\sqrt{x} + \frac{2}{\sqrt{x}} \right)^2 dx$

(c) $\int_1^2 \frac{2x^2 - 3}{x} dx$

- (10) 12. A car travels on a restricted-access highway which is 100 miles long. The speed limit on the road is 60 miles per hour. Suppose that $s(t)$, the car's distance from the beginning of the road, is a differentiable function of the time t . If the car enters the road at 9am and leaves the road at 10:15am at the other end of the highway, one and a quarter hours later, explain why the driver should expect to receive a speeding citation (ticket). Your explanation should be in complete English sentences. You should mention a specific result from this course and explain precisely why this result shows that the ticket is valid.

D16

- (12) 13. Let $f(x)$ be a function defined over the interval $[0,4]$, whose graph is shown as below. The graph of f over $[1,3]$ is a semi-circle centered at $(2, 0)$ and of radius 1, the other pieces are straight lines.



- (a) Find $\int_0^4 f(x) dx$ geometrically.

- (b) Compute the Riemann sum for $f(x)$ where $[0, 4]$ is subdivided into 4 subintervals of equal size and the sample points are the right endpoints.

- (c) Let $F(x) = \int_0^x f(t) dt$. Find $F'(2)$ and $F'(3)$.


- (10) 14. Determine the values of the parameters a and b such that the following function $f(x)$ becomes continuous and differentiable at $x = 2$:

$$f(x) = \begin{cases} x^2 - 2x + b & \text{for } x > 2 \\ ax & \text{for } x \leq 2 \end{cases}$$

D18

- (12) 15. Consider the finite region A bounded by the “wedge” $y = |x|$ and the parabola $y = \frac{1}{2}(x^2 + 1)$.

(a) Draw a sketch of the region A . [*Hint:* The parabola fits “snuggly” into the wedge.]



(b) Express the area of A in terms of one or more definite integrals.

(c) Compute the area of the region A as a sum of fractions (you do not need to simplify the answer).

- (10) 16. The minute hand of a large tower clock is 1 m long. At each full hour, the tip T of the hand points to the center C of the numeral XII. How fast is the distance between T and C changing when it is 5 : 20 am? [*Hint:* Sketch the triangle formed by C , T , and the center of the clock. Observe that it is isosceles.]

D20

More space for your work.

D2

More space for your work.