


- (8) 1. Find the solution of the differential equation

$$\frac{dy}{dx} = \frac{xy^3}{x^2 + 1}$$

satisfying the initial condition  $y(0) = 3$ . In the answer express  $y$  explicitly as a function of  $x$ .


(12) 2. Below is part of the direction field for the differential equation  $y' = y^2(1 - y)(1 + y)$ .



a) List all numbers  $k$  so that the constant function  $f(x) = k$  is a solution of this differential equation (these are the *equilibrium solutions*).

$k =$  \_\_\_\_\_


b) Sketch a typical solution curve  $y = f(x)$  to this differential equation when  $0 < y(0) < 1$ .



What is  $\lim_{x \rightarrow +\infty} f(x)$ ? \_\_\_\_\_

What is  $\lim_{x \rightarrow -\infty} f(x)$ ? \_\_\_\_\_

- (12) 3. A sequence of cubes has edges made of thin wire. The largest cube has edge length 1 inch, and each successive cube has edge length half the size of the preceding one. The first two cubes are shown here.



- a) What is the total length of wire needed to construct the edges on *all* of the cubes?

- b) What is the total volume enclosed by *all* of the cubes?

- (10) 4. Find the interval of convergence and the radius of convergence of the power series

$$\sum_{n=1}^{\infty} \frac{3^n x^n}{\sqrt{n}}.$$

In addition, determine whether the series is absolutely or conditionally convergent at the boundary points of the interval of convergence.

The radius of convergence is \_\_\_\_\_ .

The interval of convergence is \_\_\_\_\_ .

- (10) 5. The series  $\sum_{n=1}^{\infty} \frac{1}{3n^2 + 5n + 7}$  converges. Find a specific finite sum of rational numbers (quotients of integers) which is within .0001 of the sum of the infinite series. Be sure to explain why your error estimate is correct.

**Hint** Compare the “infinite tail” to something simpler, and analyze that.

(10) 6. a) Suppose the sequence  $\{A_n\}$  is defined by  $A_n = \frac{6 \cdot 4^n + 7n^3}{5 \cdot 4^n + 8n^2}$ . What is  $\lim_{n \rightarrow \infty} A_n$ ?

b) Suppose the sequence  $\{B_n\}$  is defined by  $B_n = \left(1 + \frac{3}{n}\right)^{5n}$ . What is  $\lim_{n \rightarrow \infty} B_n$ ?

**Hint** ln and l'H.

- (10) 7. Find a specific polynomial  $p(x)$  so that  $|p(x) - x^{1/4}| < .001$  for  $16 \leq x \leq 17$ . Be sure to explain why your error estimate is correct.

$n$	$2^n$
1	2
2	4
3	8
4	16
5	32
6	64
7	128
8	256
9	512
10	1,024
11	2,048
12	4,096
13	8,192

(12) 8. a) Use the Taylor series for the exponential function to write  $\int_0^1 e^{-x^2} dx$  as an infinite series. Use summation notation in your answer.

b) Find a specific finite sum of rational numbers (quotients of integers) which is within .00001 of the true value of  $\int_0^1 e^{-x^2} dx$ . Be sure to explain why your error estimate is correct.

$n$	$n!$
1	1
2	2
3	6
4	24
5	120
6	720
7	5,040
8	40,320
9	362,880
10	3,628,800

- (10) 9. Estimate the maximum error committed when  $\cos x$  is replaced by  $1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720}$  for  $x$  in the interval  $[-2, 2]$ . Be sure to explain why your error estimate is correct.

- (6) 10. a) What is the 300<sup>th</sup> Taylor polynomial for  $f(x) = 3 - 11x^5$  centered at  $a = 0$ ? Why?
- b) Explain briefly why  $g(x) = |x|$  has no Taylor series centered at  $a = 0$ .

## Second Exam for Math 152, section 72

November 29, 2001

NAME \_\_\_\_\_

Do all problems, in any order.

Show your work. An answer alone may not receive full credit.

No student notes and no calculators may be used on this exam. A formula sheet will be handed out with the exam.

Problem Number	Possible Points	Points Earned:
1	8	
2	12	
3	12	
4	10	
5	10	
6	10	
7	10	
8	12	
9	10	
10	6	
Total Points Earned:		