(12) 1. Let C be any simple closed curve described in the positive sense in the z plane, and write

$$g(w) = \int_C \frac{z^3 - 5z}{z - w} \, dz \,.$$

a) Show that  $g(w) = 2\pi i(w^3 - 5w)$  if w is inside C.

b) Show that g(w) = 0 when w is outside C.

(12) 2. Compute  $\int_B \frac{z^4}{(z - (1+i))^3} dz$  where *B* is the simple

closed curve shown: the line segment from 0 to 3, followed by the quarter-circular arc centered at 0 from 3 to 3i, followed by the line segment from 3i to 0.

Answer  $-24\pi$ 



$$f(z) = \frac{z+1}{z(z-1)}.$$

Find the Laurent series representing f(z) in the annulus 0 < |z| < 1. Be sure to explain *why* the series you write is valid in that annulus. Find explicit values of the coefficients of  $z^{10}$  and  $z^{-10}$  in the series.

(Partial) **Answer** The coefficients are -2 and 0.

(14) 4. a) Suppose that f is an entire function and there is a positive constant K so that |f(z)| > K for all z. Prove that f must be a constant function.

Hint what can you do with something that is not 0?

b) The exponential function is never 0 and is an entire function. Briefly explain why the exponential function does not contradict the assertion in part a).

(14) 5. If

$$F(z) = \frac{1}{z(z^2 - 1)(z^2 + 6)}$$

compute the integral of F over the circle of radius 2 centered at 0, oriented counterclockwise as usual. Note that  $\sqrt{6} > 2$ .

Answer  $-\frac{\pi i}{21}$ 

## (12) 6. Suppose the following is known about the coefficients of a power series $\sum_{n=0}^{\infty} a_n z^n$ :

All  $a_n$ 's are complex numbers with  $|a_n| \leq 12$ .

Explain why the power series converges for all z with  $|z| \leq \frac{1}{10}$ . Also verify that for those z's the sum of the series is always within a closed disc of radius 17 centered at 0.

Comment 17 is an overestimate!

(12) 7. The function  $g(z) = (1+3z)e^{(z^2)}$  is analytic near 0.

a) Use results about power series of familiar functions to find terms up to and including degree 4 in the Taylor series of g centered at 0.

b) Use your answer to a) to compute  $g^{(4)}(0)$ .

Answer 12

$$\lim_{z \to 0} \frac{(\cos z) - 1 + \frac{z^2}{2!}}{z^4} = L.$$

b) Suppose the function h is defined by

$$h(z) = \begin{cases} \frac{(\cos z) - 1 + \frac{z^2}{2!}}{z^4} & \text{if } z \neq 0\\ L & \text{if } z = 0 \end{cases}$$

where L is the number found in a). Explain carefully why h is entire (analytic in all of the complex numbers).

## Second Exam for Math 403, section 5

April 16, 2001

NAME \_\_\_\_\_

Do all problems, in any order. Show your work. An answer alone <u>will</u> not receive full credit. Answers <u>must</u> be supported by computation and explanation. No notes or calculators may be used on this exam.

Problem Number	Possible Points	Points Earned:
1	12	
2	12	
3	12	
4	14	
5	14	
6	12	
7	12	
8	12	
Total Points Earned:		