

## Math 504: Complex Variables (Spring, 2000)

**J1** Find a distribution solution of  $\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial y^2} = 0$  (the homogeneous wave equation) defined in  $\mathbb{R}^2$  which isn't a continuous function in  $\mathbb{R}^2$ . More precisely, show that there is no function  $w$  continuous in  $\mathbb{R}^2$  with  $w = u$  almost everywhere. Verify your claims, please!

**J2** a) Every function in  $L^1_{\text{loc}}(\mathbb{C})$  is associated to a distribution.

b)  $\log r$  is in  $L^1_{\text{loc}}(\mathbb{C})$ .

c) What is  $\Delta(\log r)$ ?

**Comment** The result of this problem is *very* important. Physicists have believed it for a long time. One proof of the answer to c) uses Green's identities and words like "charge" and "potential". Now mathematicians believe this result too.

**J3** a) Define  $T: \mathcal{O}(\mathbb{C}) \rightarrow \mathbb{C}$  by the formula  $T(f) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!}$ . Prove that  $T(f)$  is well-defined, and that  $T$  is continuous with respect to the u.c.c. topology on  $\mathcal{O}(\mathbb{C})$ .

b)  $T$  is *not* well-defined on  $\mathcal{D}(\mathbb{C})$ .

**Comment** Distributions aren't enough for all aspects of complex analysis!

The next two problems are versions of some in Forster's *Lectures on Riemann Surfaces*.

**J4** Suppose  $\{T_n\}_{n \in \mathbb{N}}$  is a sequence of distributions defined in an open subset  $U$  of  $\mathbb{C}$ . Define " $T_j \rightarrow T$ " (where  $T$  is also a distribution) to mean  $T_j(f) \rightarrow T(f)$  for every  $f \in \mathcal{D}(U)$ . Prove that if  $T_j \rightarrow T$ , then  $\frac{\partial T_j}{\partial x} \rightarrow \frac{\partial T}{\partial x}$ .

**J5** A sequence of continuous functions  $\{f_n\}_{n \in \mathbb{N}}$  defined in an open subset  $U$  of  $\mathbb{C}$  *converges weakly* to a continuous function  $f$  if  $\int_{\mathbb{C}} f_j \phi d\mathcal{L} \rightarrow \int_{\mathbb{C}} f \phi d\mathcal{L}$  for all  $\phi \in \mathcal{D}(U)$ . Here  $\mathcal{L}$  is two-dimensional Lebesgue measure. Prove that if every  $f_j$  is harmonic (resp. holomorphic) then  $f$  must be harmonic (resp. holomorphic).

The next three problems are from J. E. Marsden's text *Elementary Classical Analysis*.  $\delta$  is the Dirac  $\delta$ -function at 0.

**J6** Show that  $\delta''(f) = f''(0)$  for  $f \in \mathcal{D}(\mathbb{R})$ .

**J7** Prove that  $\sqrt{\frac{n}{\pi}} e^{-nx^2} \rightarrow \delta$  as distributions on  $\mathbb{R}$ .

**J8** Find a sequence of continuous functions  $\{g_n\}_{n \in \mathbb{N}}$  on  $\mathbb{R}$  so that  $g_n \rightarrow \delta'$  as distributions.

**J9** Suppose  $f$  is a periodic  $L^1_{\text{loc}}$  function on  $\mathbb{R}$  with period  $2\pi$ . If  $a_n = \frac{1}{2\pi} \int_0^{2\pi} f(t) e^{-int} dt$ , show that  $f = \sum_{-\infty}^{\infty} a_n e^{int}$  as distributions.

**J10** Suppose that  $\{a_n\}_{n \in \mathbb{Z}}$  is a sequence of complex numbers with *polynomial growth*: there are  $K \in \mathbb{N}$  and  $c \in \mathbb{R}$  so that  $|a_n| \leq c|n|^K$  for all  $n \in \mathbb{Z}$ . Then  $\sum_{-\infty}^{\infty} a_n e^{inx}$  converges to a distribution defined on  $\mathbb{R}$ .

**J11** "Let  $\mathbf{L}$  denote the subset of  $\mathbb{R}^2$  defined by  $\{x = 0 \ \& \ y \geq 0\} \cup \{x \geq 0 \ \& \ y = 0\}$  (the union of the non-negative  $x$ - and  $y$ -axes)." \* Let  $S$  be the distribution defined by integrating over  $\mathbf{L}$  \*\*. What is  $\frac{\partial S}{\partial z}$ ? Can you find  $T$  so that  $\frac{\partial S}{\partial z} = T$ ?

\* Of course this is quoted from the beginning of homework A! We end as we began.

\*\* Use  $dx$  and  $dy$  appropriately.