

## Math 504: Complex Variables (Spring, 2000)

In this problem set,  $U$  will be a fixed open subset of  $\mathbb{C}$ . “ $L^2$ ” in the first two problems below is with respect to area measure in the plane.

**C1** a) Suppose that  $f \in L^2(D_\varepsilon(0)) \cap \mathcal{O}(D_\varepsilon(0))$  for some  $\varepsilon > 0$ . Express the  $L^2$  norm of  $f$  in terms of the power series coefficients of  $f$ , and prove that, given  $j \in \mathbb{N} \cup \{0\}$ , there is  $C = C(j, \varepsilon) > 0$  so that  $|f^{(j)}(0)| \leq C \|f\|_{L^2(D_\varepsilon(0))}$ .

b) Suppose  $K$  is compact in  $U$ , and  $\varepsilon > 0$  is such that  $K_\varepsilon \subset U$ . Given  $j \in \mathbb{N} \cup \{0\}$ , prove that there is  $C = C(j, \varepsilon) > 0$  so that  $\|f^{(j)}\|_K \leq C \|f\|_{L^2(K_\varepsilon)}$  for all  $f \in \mathcal{O}(U)$ .

**C2** Please assume the results of the previous problem.

a) Suppose  $\{f_k\}_{k \in \mathbb{N}}$  is a sequence of functions in  $L^2(U) \cap \mathcal{O}(U)$ . Prove that if  $\{f_k\}_{k \in \mathbb{N}}$  is  $L^2$ -Cauchy, then  $\{f_k\}_{k \in \mathbb{N}}$  is u.c.c.-Cauchy.

b) If  $f$  is the u.c.c. limit of the sequence in a), then  $f$  is also the  $L^2$  limit of the sequence.

**C3** Suppose  $\{f_j\}_{j \in \mathbb{N}}$  is a sequence of functions in  $\mathcal{O}(U)$  which converges *pointwise* to  $f$  in  $U$ . Prove that there is a dense open subset  $V$  of  $U$  so that  $f \in \mathcal{O}(V)$  and  $f_j \rightarrow f$  u.c.c. in  $V$ . What happens if the sequence of  $f_j$ 's has pointwise bounded modulus at every  $z \in U$ ?

**C4** There is  $f \in \mathcal{M}(\mathbb{C})$  which has a pole of order  $j$  at each  $j \in \mathbb{N}$  (and those are its only poles). Exhibit such a function precisely, and give  $a, b \in \mathbb{Q}$  so that  $|f(1+i) - (a+bi)| < .001$ .

**C5** There is  $f \in \mathcal{M}(\mathbb{C})$  which has simple poles at  $2^j$  for each  $j \in \mathbb{N}$  (and those are its only poles), and whose principal part at  $2^j$  is  $\frac{3^j}{z - 2^j}$  for each  $j$ . Exhibit such a function precisely, and give  $a, b \in \mathbb{Q}$  so that  $|f(1+i) - (a+bi)| < .001$ .

**C6** (Ridiculous polynomials, **I**) Show that there is a sequence of polynomials  $P_n(z) \in \mathbb{C}[z]$  so that as  $n \rightarrow \infty$ ,  $P_n(0) = 1$  and  $P_n(z) \rightarrow 0$  for  $z \neq 0$ .

**C7** (Ridiculous polynomials, **II**) Show that there is a sequence of polynomials  $P_n(z) \in \mathbb{C}[z]$  so that as  $n \rightarrow \infty$ ,  $P_n(z) \rightarrow 1$  for  $\operatorname{Re} z \geq 0$  and  $P_n(z) \rightarrow 0$  for  $\operatorname{Im} z < 0$ .

**C8** (Ridiculous polynomials, **III**) Show that there is a sequence of polynomials  $P_n(z) \in \mathbb{C}[z]$  so that as  $n \rightarrow \infty$ ,  $P_n(z) \rightarrow 1$  if  $|z| = 1$  and  $P_n(z) \rightarrow 0$  otherwise.

**C9** (Ridiculous polynomials, **IV**) Show that there is a sequence of polynomials  $P_n(z) \in \mathbb{C}[z]$  so that as  $n \rightarrow \infty$ ,  $P_n(z) \rightarrow 0$  for all  $z \in \mathbb{C}$  but  $P_n \not\rightarrow 0$  uniformly on all compact subsets of  $\mathbb{C}$ .

**C10** (Ridiculous polynomials, **V**) Prove that there is  $f \in \mathcal{O}(D_1(0))$  so that for all  $\zeta \in \partial D_1(0)$  and all  $\varepsilon > 0$ , the closure of  $\{f(s\zeta) : 1 - \varepsilon < s < 1\}$  is all of  $\mathbb{C}$ . (For such  $f$ 's, radial limits are wildly non-existent! One way to show that such a function exists is to write it as the u.c.c. limit in  $D_1(0)$  of a sequence of “ridiculous” polynomials.)

**C11** Give some metric  $d$  on  $\mathcal{O}(\mathbb{C})$  where convergence in  $d$  is equivalent to u.c.c. Find positive numbers  $A$  and  $B$  with  $A \leq d(z, z^2) \leq B$ . If the diameter of  $\mathcal{O}(\mathbb{C})$  as measured by  $d$  is finite, please supply a valid  $B$  less than the diameter (include verification of this).