

- (8) 1. Suppose
- $p = (1, 0, 2)$
- ,
- $q = (0, 2, 2)$
- , and
- $r = (1, 1, 1)$
- are points in
- $\mathbb{R}^3$
- .

a) Find a vector orthogonal to the plane through the points  $p$ ,  $q$ , and  $r$ .**Answer** The vector  $\overrightarrow{pq}$  is  $\langle -1, 2, 0 \rangle$  and the vector  $\overrightarrow{pr}$  is  $\langle 0, 1, -1 \rangle$ .The cross product is a vector perpendicular to these vectors:  $\overrightarrow{pq} \times \overrightarrow{pr} = \det \begin{pmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 2 & 0 \\ 0 & 1 & -1 \end{pmatrix} = -2\mathbf{i} - \mathbf{j} - \mathbf{k}$ .b) Find the area of triangle  $pqr$ .**Answer** The area of triangle  $pqr$  is half the magnitude of  $\overrightarrow{pq} \times \overrightarrow{pr}$ , so it is  $\frac{1}{2}\sqrt{(-2)^2 + (-1)^2 + 1^2} = \frac{1}{2}\sqrt{6}$ .

- (12) 2. Suppose
- $\mathbf{V} = \langle 1, 3, -2 \rangle$
- and
- $\mathbf{W} = \langle -1, 2, 1 \rangle$
- . Find vectors
- $\mathbf{V}_1$
- and
- $\mathbf{V}_2$
- so that
- $\mathbf{V} = \mathbf{V}_1 + \mathbf{V}_2$
- ,
- $\mathbf{V}_1$
- is parallel to
- $\mathbf{W}$
- , and
- $\mathbf{V}_2$
- is perpendicular to
- $\mathbf{W}$
- .

**Answer**  $\mathbf{V}_1 = \frac{\mathbf{V} \cdot \mathbf{W}}{|\mathbf{W}|^2} \mathbf{W}$  so compute  $\mathbf{V} \cdot \mathbf{W} = 1 \cdot (-1) + 3 \cdot 2 + (-2) \cdot 1 = -1 + 6 - 2 = 3$  and  $|\mathbf{W}|^2 = (-1)^2 + 2^2 + 1^2 = 6$ . Then  $\mathbf{V}_1 = \frac{3}{6} \langle -1, 2, 1 \rangle = \langle -\frac{1}{2}, 1, \frac{1}{2} \rangle$ . Since  $\mathbf{V} = \mathbf{V}_1 + \mathbf{V}_2$ ,  $\mathbf{V}_2 = \mathbf{V} - \mathbf{V}_1 = \langle 1, 3, -2 \rangle - \langle -\frac{1}{2}, 1, \frac{1}{2} \rangle = \langle \frac{3}{2}, 2, -\frac{5}{2} \rangle$ . If desired, a check that  $\mathbf{W} \perp \mathbf{V}_2$  is:  $\mathbf{W} \cdot \mathbf{V}_2 = (-1) \left(\frac{3}{2}\right) + 2 \cdot 2 + 1 \left(-\frac{5}{2}\right) = 0$ .


- (10) 3. The point
- $(1, 0, 2)$
- is on the plane
- $P$
- whose equation is given by
- $2x - 2y + z = 4$
- .

a) Find parametric equations for the line perpendicular to  $P$  through  $(1, 0, 2)$ .**Answer** The normal vector of  $P$  is  $\langle 2, -2, 1 \rangle$ , the line's direction. Parametric equations are  $\begin{cases} x = 1 + 2t \\ y = 0 - 2t \\ z = 2 + 1t \end{cases}$ .b) Find a point on the line in a) which has distance 5 to  $P$ .**Answer** The distance from  $(1 + 2t, -2t, 2 + t)$  to  $P$  is just the distance to  $(1, 0, 2)$ , and this distance is  $\sqrt{(2t)^2 + (-2t)^2 + t^2} = 3|t|$ . This will be 5 when  $|t| = \frac{5}{3}$  so  $t = \pm \frac{5}{3}$ . The positive root substituted in the parametric equations gives the point  $(1 + \frac{10}{3}, -\frac{10}{3}, 2 + \frac{5}{3}) = (\frac{13}{3}, -\frac{10}{3}, \frac{11}{3})$ .c) Write an equation for a plane parallel to  $P$  whose distance to  $P$  is 5.**Answer** A normal vector is  $\langle 2, -2, 1 \rangle$  and one such plane passes through  $(\frac{13}{3}, -\frac{10}{3}, \frac{11}{3})$ . Therefore an equation for the plane is  $2(x - \frac{13}{3}) - 2(y + \frac{10}{3}) + 1(z - \frac{11}{3}) = 0$ .

- (12) 4. Suppose the position vector of a curve is given by
- $\mathbf{r}(t) = \langle e^t, e^{-t}, \sqrt{2}t \rangle$
- . Find the unit tangent and unit normal vectors when the parameter
- $t = 1$
- . That is, find
- $\mathbf{T}(1)$
- and
- $\mathbf{N}(1)$
- .

**Answer** Since  $\mathbf{r}(t) = \langle e^t, e^{-t}, \sqrt{2}t \rangle$ ,  $\mathbf{r}'(t) = \langle e^t, -e^{-t}, \sqrt{2} \rangle$  and  $|\mathbf{r}'(t)| = \sqrt{(e^t)^2 + (-e^{-t})^2 + (\sqrt{2})^2} = \sqrt{e^{2t} + e^{-2t} + 2} = \sqrt{(e^t + e^{-t})^2} = e^t + e^{-t}$ . So  $\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} = \left\langle \frac{e^t}{e^t + e^{-t}}, \frac{-e^{-t}}{e^t + e^{-t}}, \frac{\sqrt{2}}{e^t + e^{-t}} \right\rangle$ , and  $\frac{d}{dt} \mathbf{T}(t) = \left\langle \frac{e^t(e^t + e^{-t}) - (e^t - e^{-t})e^t}{(e^t + e^{-t})^2}, \frac{(-1)^2 e^{-t}(e^t + e^{-t}) - (e^t - e^{-t})(-e^{-t})}{(e^t + e^{-t})^2}, \frac{-(e^t - e^{-t})\sqrt{2}}{(e^t + e^{-t})^2} \right\rangle$ .  $\mathbf{T}(1) = \left\langle \frac{e}{e+e^{-1}}, \frac{-e^{-1}}{e+e^{-1}}, \frac{\sqrt{2}}{e+e^{-1}} \right\rangle$ .  $\mathbf{T}'(1) = \left\langle \frac{e(e+e^{-1}) - (e-e^{-1})e}{(e+e^{-1})^2}, \frac{e^{-1}(e+e^{-1}) + e^{-1}(e-e^{-1})}{(e+e^{-1})^2}, \frac{-(e-e^{-1})\sqrt{2}}{(e+e^{-1})^2} \right\rangle = \left\langle \frac{2}{(e+e^{-1})^2}, \frac{2}{(e+e^{-1})^2}, \frac{-(e-e^{-1})\sqrt{2}}{(e+e^{-1})^2} \right\rangle$ .  $\mathbf{N}(1)$  is a unit vector in the direction of  $\mathbf{T}'(1)$ , so consider  $\langle 2e, 2e, -(e^2 - 1)\sqrt{2} \rangle$ , a positive multiple of  $\mathbf{T}'(1)$  whose length is  $\sqrt{8e^2 + 2(e^4 - 2e^2 + 1)} = \sqrt{2(e^4 + 2e^2 + 1)} = \sqrt{2}(e^2 + 1)$ . "Thus"  $\mathbf{N}(1) = \left\langle \frac{2e}{\sqrt{2}(e^2 + 1)}, \frac{2e}{\sqrt{2}(e^2 + 1)}, \frac{-(e^2 - 1)\sqrt{2}}{\sqrt{2}(e^2 + 1)} \right\rangle$ .


- (12) 5. A point is moving along the curve below in the direction indicated. Its motion is parameterized by arc length,
- $s$
- , so that it is moving at unit speed. Arc length is measured from the point
- $P$
- (both backward and forward). The curve is intended to continue indefinitely both forward and backward in
- $s$
- , with its forward motion curling more and more tightly around the indicated circle,
- $B$
- , and, backward, curling more and more tightly around the other circle,
- $A$
- . Near
- $P$
- the curve is parallel to the indicated line segment.

Sketch a graph of the curvature,  $\kappa$ , as a function of the arc length,  $s$ . What are  $\lim_{s \rightarrow +\infty} \kappa(s)$  and  $\lim_{s \rightarrow -\infty} \kappa(s)$ ?

Use complete English sentences to briefly explain the numbers you give.

\* I cheated, sort of. I computed this by hand, and got a horrible mess. I had Maple simplify the answer which I "reverse engineered" to get the displayed computation. Any correct mess submitted is certainly acceptable!

**Answer** As  $s \rightarrow -\infty$ ,  $\kappa \rightarrow 1$ : the curve is getting closer to a circle of radius 1 with curvature = 1. There should be an interval of length  $\approx 4$  centered at  $s = 0$  where the curve is “flat”,  $\kappa = 0$ . As  $s \rightarrow \infty$ ,  $\kappa \rightarrow \frac{1}{2}$ : the curve is getting closer to a circle of radius 2 with curvature =  $\frac{1}{2}$ . The graph should decrease and then increase. Also,  $\lim_{s \rightarrow -\infty} \kappa(s) = 1$  and  $\lim_{s \rightarrow +\infty} \kappa(s) = \frac{1}{2}$ .



- (10) 6. Find the linear approximation of the function  $f(x, y) = \sqrt{20 - x^2 - 7y^2}$  at  $(2, 1)$  and use it to approximate  $f(1.95, 1.08)$ .

**Answer**  $f(2, 1) = \sqrt{20 - 2^2 - 7 \cdot 1^2} = \sqrt{9} = 3$ ;  $f_x = \frac{1}{2}(20 - x^2 - 7y^2)^{-\frac{1}{2}}(-2x)$  so  $f_x(2, 1) = \frac{1}{2} \cdot \frac{1}{3}(-2 \cdot 2) = -\frac{2}{3}$ ;  $f_y = \frac{1}{2}(20 - x^2 - 7y^2)^{-\frac{1}{2}}(-14y)$  so  $f_y(2, 1) = \frac{1}{2} \cdot \frac{1}{3}(-14 \cdot 1) = -\frac{7}{3}$ . The linear approximation of  $f(x, y)$  at  $(2, 1)$  is  $f(2 + \Delta x, 1 + \Delta y) \approx 3 - \frac{2}{3}\Delta x - \frac{7}{3}\Delta y$ . To approximate  $f(1.95, 1.08)$  take  $\Delta x = -.05$  and  $\Delta y = .08$ . The result is  $3 + \frac{2}{3}(.05) - \frac{7}{3}(.08)^{**}$ .

- (12) 7. a) If  $z = f(x^2 - 3y)$ , show that  $3\frac{\partial z}{\partial x} + 2x\frac{\partial z}{\partial y} = 0$ .

**Answer**  $z_x = f'(x^2 - 3y)(2x)$ ;  $z_y = f'(x^2 - 3y)(-3)$ . Therefore  $3\frac{\partial z}{\partial x} + 2x\frac{\partial z}{\partial y} = 3f'(x^2 - 3y)(2x) + (2x)f'(x^2 - 3y)(-3) = (6x - 6x)f'(x^2 - 3y) = 0$ .

b) If  $z = f(x^2 - 3y)$ , then  $\frac{\partial^2 z}{\partial x^2} = A(x)f''(x^2 - 3y) + B(x)f'(x^2 - 3y)$  where  $A(x)$  and  $B(x)$  are simple functions of  $x$  alone or constants. What are  $A(x)$  and  $B(x)$ ?

**Answer** Let's  $\frac{\partial}{\partial x}$  the equation  $z_x = f'(x^2 - 3y)(2x)$ . We need to apply the product rule and the chain rule. The result is  $f''(x^2 - 3y)(2x)^2 + f'(x^2 - 3y)2$  so that  $A(x) = (2x)^2$  or  $4x^2$  and  $B(x) = 2$ .

- (12) 8. Suppose that  $F(x, y, z) = x^2 + 3yz$  and  $p = (-3, 2, -1)$ .

a) Find the maximum directional derivative of  $F$  at  $p$  and write a unit vector pointing in the direction this maximum value occurs.

**Answer**  $\nabla F(x, y, z) = \langle 2x, 3z, 3y \rangle$  so that  $\nabla F(-3, 2, -1) = \langle 2(-3), 3(-1), 3 \cdot (2) \rangle = \langle -6, -3, 6 \rangle$ . The maximum directional derivative is  $|\nabla F(-3, 2, -1)| = \sqrt{(-6)^2 + (-3)^2 + 6^2} = \sqrt{81} = 9$ . A unit vector in the desired direction is  $\frac{1}{|\nabla F(-3, 2, -1)|} \nabla F(-3, 2, -1)$  which is  $\langle -\frac{6}{9}, -\frac{3}{9}, \frac{6}{9} \rangle$ .

b) Suppose  $C = F(-3, 2, -1)$ . Compute  $C$  and write an equation for the plane tangent to the surface  $F(x, y, z) = C$  at the point  $p$ .

**Answer**  $F(-3, 2, -1) = (-3)^2 - 3 \cdot 2(-1) = 3$ , so  $C = 3$ .  $\nabla F(-3, 2, -1)$  gives a normal vector, so an equation for a plane tangent to  $F(x, y, z) = 3$  at  $p$  is  $-6(x - 3) - 3(y - 2) + 6(z + 1) = 0$ .

- (12) 9. a) If  $f(x, y, z) = x^2 + y^2$ , compute  $\nabla f(x, y, z)$ . What are  $f(2, 1, 2)$  and  $\nabla f(2, 1, 2)$ ?


**Answer**  $\nabla f(x, y, z) = \langle 2x, 2y, 0 \rangle$ .  $f(2, 1, 2) = 2^2 + 1^2 = 5$  and  $\nabla f(2, 1, 2) = \langle 4, 2, 0 \rangle$ .

b) If  $g(x, y, z) = x^2 + y^2 + z^2 - xy - yz$ , compute  $\nabla g(x, y, z)$ . What are  $g(2, 1, 2)$  and  $\nabla g(2, 1, 2)$ ?

**Answer**  $\nabla g(x, y, z) = \langle 2x - y, 2y - x - z, 2z - y \rangle$ .  $g(2, 1, 2) = 2^2 + 1^2 + 2^2 - 2 - 2 = 5$  and  $\nabla g(2, 1, 2) = \langle 3, -2, 3 \rangle$ .

c) The point  $(2, 1, 2)$  is on both the surface  $x^2 + y^2 = 5$ , a circular cylinder whose axis of symmetry is the  $z$ -axis, and the surface  $x^2 + y^2 + z^2 - xy - yz = 5$ , an ellipsoid tilted with respect to the coordinate axes. The surfaces intersect in a curve. The surfaces and the curve are shown in the picture to the right. Find a vector tangent to that curve at  $(2, 1, 2)$ . Your answers to a) and b) can be used here.

**Answer** A vector tangent to the curve is perpendicular to both surface normals.  $\nabla f(2, 1, 2) \times \nabla g(2, 1, 2)$  is such a vector, so  $\det \begin{pmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 4 & 2 & 0 \\ 3 & -2 & 3 \end{pmatrix} = 6\mathbf{i} - 12\mathbf{j} - 14\mathbf{k}$ , one valid answer.



\*\* Maple reports the “true value” is about 2.834 and the approximation is about 2.847, if this matters.