

Your goal here is to compute $\int_{-\infty}^{\infty} \frac{1}{x^2 + x + 1} dx$ using the Residue Theorem.

1 We may write $\int_{-\infty}^{\infty} \frac{1}{x^2 + x + 1} dx = \lim_{R \rightarrow \infty} \int_{I_R} f(z) dz$ where I_R is the interval from $-R$ to R on the real line. f is an analytic function with two isolated singularities. It is given by a formula:

$f(z) =$ _____.

2 The isolated singularities of f are at the complex numbers $A =$ _____ in the upper half plane and $B =$ _____ in the lower half plane.

The type of the isolated singularity at A is (circle one)

A POLE A REMOVABLE SINGULARITY AN ESSENTIAL SINGULARITY

3 We can factor the denominator of f and rewrite its formula as follows:


$f(z) =$ _____.

4 The function f has an isolated singularity at A in the upper half plane. Since f has a _____ (insert here a useful and precise descriptive phrase about f 's singularity at A), the residue of f at A is easy to compute.

That residue is _____.

5 Suppose S_R is the counterclockwise oriented semicircle of radius R in the upper half plane centered at 0. We will write $I_R + S_R$ to mean the simple closed curve obtained by following I_R by S_R . If R is large enough (as shown) so A is inside $I_R + S_R$, the Residue Theorem tells us that

the value of $\int_{I_R + S_R} f(z) dz$ is _____.



6 If $|z| = R$ where R is a large positive number, then the “reverse triangle inequality” applied to the original formula for f allows us to underestimate the denominator of $|f(z)|$ in terms of R (note that there will be several expressions that are subtracted):

So the denominator of $|f(z)| \geq$ _____.

7 The length of S_R is _____. The ML inequality allows us to overestimate the modulus of $\int_{S_R} f(z) dz$ in terms of R .

$$\left| \int_{S_R} f(z) dz \right| \leq \underline{\hspace{2cm}}.$$

8 We therefore conclude that $\lim_{R \rightarrow \infty} \int_{S_R} f(z) dz =$ _____.

9 Combine the results of **8** and **5** to compute the value of $\lim_{R \rightarrow \infty} \int_{I_R} f(z) dz$.

The value of this limit is _____.

10 Putting it all together, we finally can write

$$\int_{-\infty}^{\infty} \frac{1}{x^2 + x + 1} dx = \underline{\hspace{2cm}}.$$

Maple reports that this quantity is approximately 3.62759 87284 68435 7012.