

Formula sheet for the first exam in Math 135:F2, summer 2006

Function	Derivative
k (const.)	0
x^n	nx^{n-1}
e^x	e^x
$\sin x$	$\cos x$
$\cos x$	$-\sin x$
$\tan x$	$(\sec x)^2$
$Cf(x)$	$Cf'(x)$ (C const.)
$f(x) + g(x)$	$f'(x) + g'(x)$
$f(x) \cdot g(x)$	$f'(x) \cdot g(x) + f(x) \cdot g'(x)$
$\frac{f(x)}{g(x)}$	$\frac{f'(x)g(x) - g'(x)f(x)}{g(x)^2}$
$f(g(x))$	$f'(g(x)) \cdot g'(x)$

Exponential properties	
$a^{b+c} = a^b \cdot a^c$	$a^{-b} = 1/a^b$
$(a^b)^c = a^{bc}$	$e \approx 2.718$
$a^0 = 1$	$e^{\ln a} = a$ if $a > 0$

Logarithm properties		
$\ln(a \cdot b) = \ln a + \ln b$	$\ln(a^b) = b \ln(a)$	
$\ln(a/b) = \ln(a) - \ln(b)$	$\ln(\frac{1}{b}) = -\ln(b)$	
$\ln(e^a) = a$	$\ln(1) = 0$	$\ln(e) = 1$

Miscellaneous formulas

If $a \neq 0$, the **roots** of $ax^2 + bx + c = 0$ are $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Distance between (a, b) and (c, d) : $\sqrt{(a - c)^2 + (b - d)^2}$.

Circle center (h, k) and radius r : $(x - h)^2 + (y - k)^2 = r^2$.

f is **continuous** at w if $\lim_{x \rightarrow w} f(x)$ exists and equals $f(w)$.

Intermediate Value Theorem

If $f(x)$ is continuous in $a \leq x \leq b$, then $f(x)$'s values include all numbers between $f(a)$ and $f(b)$.

$f'(x)$, the **derivative of f at x** , is defined to be $\lim_{h \rightarrow 0} \frac{f(x+h)-f(x)}{h}$.

Area and Volume Formulas

Rectangle $A = \text{LENGTH} \cdot \text{WIDTH}$

Circle $A = \pi \text{ RADIUS}^2$

Triangle $A = \frac{1}{2} \text{ BASE} \cdot \text{HEIGHT}$

Box $V = \text{LENGTH} \cdot \text{WIDTH} \cdot \text{HEIGHT}$

Cylinder $V = \pi \text{ RADIUS}^2 \cdot \text{HEIGHT}$

Sphere $V = \frac{4}{3}\pi \text{ RADIUS}^3$

Special values of trig functions

θ	$\sin \theta$	$\cos \theta$	$\tan \theta$
0	0	1	0
$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
$\frac{\pi}{4}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	1
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
$\frac{\pi}{2}$	1	0	Undefined