

POSITIVE OPERATORS AND HAUSDORFF DIMENSION OF INVARIANT SETS

ROGER D. NUSSBAUM, AMIT PRIYADARSHI, AND SJOERD VERDUYN LUNEL

ABSTRACT. In this paper we obtain theorems which give the Hausdorff dimension of the invariant set for a finite family of contraction mappings which are “infinitesimal similitudes” on a complete, perfect metric space. Our work generalizes the graph-directed construction of Mauldin and Williams [13] and is related in its general setting to results of Schief [22], but differs crucially in that the mappings need not be similitudes. We use the theory of positive linear operators and generalizations of the Krein-Rutman theorem to characterize the Hausdorff dimension as the unique value of $\sigma > 0$ for which $r(L_\sigma) = 1$, where L_σ , $\sigma > 0$, is a naturally associated family of positive linear operators and $r(L_\sigma)$ denotes the spectral radius of L_σ . We also indicate how these results can be generalized to countable families of infinitesimal similitudes. The intent here is foundational: to derive a basic formula in its proper generality and to emphasize the utility of the theory of positive linear operators in this setting. Later work will explore the usefulness of the basic theorem and its functional analytic setting in studying questions about Hausdorff dimension.

1. INTRODUCTION

Given N contraction mappings θ_i , $1 \leq i \leq N$, on a complete metric space (X, d) , there exists a unique, nonempty compact set C such that $C = \bigcup_{i=1}^N \theta_i(C)$. C is called an invariant set or an attractor for the family $\{\theta_i\}_{i=1}^N$. A general problem is to obtain theorems which allow the accurate estimation of the Hausdorff dimension of C . A well studied case is when the maps θ_i are “similitudes”, i.e., when for $1 \leq i \leq N$, there exists r_i , $0 < r_i < 1$, with $d(\theta_i(x), \theta_i(y)) = r_i d(x, y)$ for all $x, y \in X$. If, in addition, X is a normed linear space with metric d derived from the norm on X and if the similitudes θ_i are onto maps (which is necessarily true if X is finite dimensional), then a theorem of Mazur and Ulam (see [14] or [23]) implies that each θ_i is an affine linear map. Moran [15] and Hutchinson [7] have studied the case that each θ_i , $1 \leq i \leq N$, is an affine linear similitude on a finite dimensional normed linear space X . Provided the “pieces” $\theta_i(C)$ do not overlap too much, they have proved that the Hausdorff dimension of C is the same as the “similarity dimension” α and is determined by the equation $\sum_{i=1}^N r_i^\alpha = 1$. More precisely, one needs the “open set condition”, i.e., the assumption that there exists a nonempty, open set U such that the sets $\theta_i(U)$ are contained in U and are pairwise disjoint. Mauldin and Williams [13] have generalized the idea of “self-similarity” to

2000 *Mathematics Subject Classification.* Primary 37F35, 28A80; Secondary 37C30, 47B65.

Key words and phrases. Hausdorff dimension, iterated function systems, positive operators, spectral radius.

The first author was supported in part by NSF Grant DMS-0701171.

“graph self-similarity” which allows a larger class of sets like C , but still the maps in question are affine linear similitudes. Schief [22] works in a setting similar to ours and considers similitudes on general complete metric spaces and obtains analogous results. Interestingly, he has shown that in this generality, the open set condition is no longer sufficient and must be strengthened to the “strong open set condition”.

It is of considerable interest (see [2, 3]) to allow maps θ_i which may not be affine linear. For example, in studying subsets of \mathbb{R} defined by properties of their continued fraction expansions, one is led to maps $\theta_i : [0, 1] \rightarrow [0, 1]$ defined by $\theta_i(x) = (x + m_i)^{-1}$, m_i a positive integer.

In this paper we shall consider a complete, perfect metric space (X, d) and maps $\theta_i : X \rightarrow X$, $1 \leq i \leq N$, where θ_i is a contraction mapping and an “infinitesimal similitude” (instead of a similitude) for $1 \leq i \leq N$. If G is a bounded open subset of \mathbb{C} , $\theta_j : G \rightarrow G$ is analytic and $\overline{\theta_j(G)} \subset G$ for $1 \leq j \leq N$ and $X = \bigcup_{j=1}^N \overline{\theta_j(G)}$ with appropriate metric d , we obtain an important example for which the mappings $\theta_j : X \rightarrow X$ are contractions and infinitesimal similitudes. For the general definition of infinitesimal similitudes and their properties, see the beginning of Section 3. If C denotes the nonempty, compact invariant set for $\{\theta_i\}_{i=1}^N$ and if $\theta_i(C)$ and $\theta_j(C)$ are disjoint for $1 \leq i < j \leq N$, we shall obtain below a formula for the Hausdorff dimension of C . In fact, we shall obtain such a formula in a setting similar to that of Mauldin and Williams, but using contractions and infinitesimal similitudes, rather than affine linear contractions which are similitudes.

The classical Krein-Rutman theorem (see [8]) considers a positive (in the sense of mapping a suitable cone to itself), compact, linear map $T : X \rightarrow X$ which has positive spectral radius r and asserts the existence of a positive eigenvector v with $T(v) = rv$. Generalizations, particularly allowing noncompact T , can be found in [1, 9, 10, 18, 19, 21]. Our approach in this paper will be to use generalizations of the Krein-Rutman theorem. To each nonnegative real σ we shall associate a positive linear operator L_σ on a Banach space Y of continuous functions. We shall prove that L_σ has a positive eigenvector with eigenvalue $r(L_\sigma)$, the spectral radius of L_σ . We shall prove that σ_0 , the desired Hausdorff dimension, is the unique value of $\sigma \geq 0$ for which $r(L_\sigma) = 1$. We shall not use the thermodynamic formalism. Curiously, we have found no references to the Krein-Rutman theorem in the Hausdorff dimension literature, despite its relevance. Analogues of the operator L_σ we consider are sometimes called “Perron-Frobenius operators” or “Frobenius-Ruelle operators”, although the theory originally developed by Perron and Frobenius is restricted to matrices with nonnegative entries, and generalizations to infinite dimensions pose substantial difficulties.

For the convenience of the reader we shall now state our main theorem in the simpler setting of iterated function systems on a compact, perfect metric space. For the more general case see section 3.

Let (S, d) be a compact, perfect metric space. If $\theta : S \rightarrow S$, we shall say that θ is an infinitesimal similitude at $t \in S$ if for any sequences $(s_k)_k$ and $(t_k)_k$ with $s_k \neq t_k$ for $k \geq 1$ and $s_k \rightarrow t$, $t_k \rightarrow t$, the limit

$$(1.1) \quad \lim_{k \rightarrow \infty} \frac{d(\theta(s_k), \theta(t_k))}{d(s_k, t_k)} =: (D\theta)(t)$$

exists and is independent of the particular sequences $(s_k)_k$ and $(t_k)_k$. We shall say that θ is an infinitesimal similitude on S if θ is an infinitesimal similitude at t for all $t \in S$.

Assume that for $1 \leq i \leq N$, $\theta_i : S \rightarrow S$ is a Lipschitz map with Lipschitz constant $c_i \leq c < 1$. Then we know that there exists a unique nonempty, compact set $C \subset S$ with

$$C = \bigcup_{i=1}^N \theta_i(C).$$

Assume the map $\theta_i : S \rightarrow S$ is an infinitesimal similitude on S and the map $t \rightarrow (D\theta_i)(t)$ is a strictly positive Hölder continuous function on S for $1 \leq i \leq N$. For $\sigma \geq 0$, define $L_\sigma : C(S) \rightarrow C(S)$ by

$$(1.2) \quad (L_\sigma f)(t) = \sum_{i=1}^N ((D\theta_i)(t))^\sigma f(\theta_i(t)).$$

It follows (see Theorem 5.4 in [19]) that L_σ has a strictly positive eigenvector u_σ with eigenvalue equal to the spectral radius $r(L_\sigma)$ of L_σ . We also have the following lemma.

Lemma 1.1. *The map $\sigma \mapsto r(L_\sigma)$ is continuous and strictly decreasing. Furthermore, there is a unique $\sigma_0 \geq 0$ such that $r(L_{\sigma_0}) = 1$.*

We are now ready to state the theorem about the Hausdorff dimension of the invariant set C .

Theorem 1.2. *Let $\theta_i : S \rightarrow S$, $1 \leq i \leq N$, be infinitesimal similitudes and assume that the map $t \mapsto (D\theta_i)(t)$ is a strictly positive Hölder continuous function on S . Assume that $\theta_i : S \rightarrow S$ is a Lipschitz map with Lipschitz constant $c_i \leq c < 1$ and let C denote the unique, nonempty, compact set such that*

$$C = \bigcup_{i=1}^N \theta_i(C).$$

Further, assume that θ_i , $1 \leq i \leq N$, satisfy

$$\theta_i(C) \cap \theta_j(C) = \emptyset \text{ for } 1 \leq i, j \leq N, i \neq j$$

and are one-to-one on C . Then the Hausdorff dimension, $\dim(C)$ of C is given by the unique σ_0 such that $r(L_{\sigma_0}) = 1$.

To see that this is a special case of our general theory of later sections, let $V = \{1\}$, $\mathcal{E} = \{1, 2, \dots, N\}$ and $\Gamma = V \times \mathcal{E}$ in the terminology below.

We should remark that our proofs require that the pieces $\theta_i(C)$, $1 \leq i \leq N$, be pairwise disjoint. It would be very interesting to find variant arguments which allowed some overlap. For instance, we do not know if the strong open set condition is sufficient to get the results in this generality.

The paper is organised as follows. In §2 we introduce the basic set-up and prove the existence of the invariant set. In §3 we discuss the Perron-Frobenius operators. In §4 we give the main theorems about the Hausdorff dimension of the invariant set. In §5 the results are extended to infinite iterated function systems, and we show that a class of complex continued fraction previously studied by Mauldin and Urbanski [12] is also covered by our theorems. In §6 we show that choosing an appropriate metric gives a large class of examples of iterated function systems to which our theory can be applied. For instance, we discuss the Carathéodory-Reiffen-Finsler (CRF) metric on bounded open subsets of \mathbb{C} .

2. INVARIANT SETS

Let V and \mathcal{E} be finite sets and for each $v \in V$, let (S_v, d_v) be a complete metric space. Let Γ be a subset of $V \times \mathcal{E}$, and $\alpha : \Gamma \rightarrow V$. For each $(v, e) \in \Gamma$, let $\theta_{(v,e)} : S_v \rightarrow S_{\alpha(v,e)}$ be a Lipschitz map with $\text{Lip}(\theta_{(v,e)}) \leq c < 1$. Recall that a map $\psi : (S_1, d_1) \rightarrow (S_2, d_2)$ is said to be Lipschitz if there is a constant c such that $d_2(\psi(s), \psi(t)) \leq c d_1(s, t) \forall s, t \in S_1$, and

$$\text{Lip}(\psi) := \sup \left\{ \frac{d_2(\psi(s), \psi(t))}{d_1(s, t)} : s, t \in S_1, s \neq t \right\}.$$

Notation and assumptions will be as in the preceding paragraph for the remainder of the paper. We shall keep in mind two important particular cases.

Example 2.1. (*The Mauldin-Williams graph*) Let V be the set of vertices and \mathcal{E} be the set of edges of a directed multigraph. Let $i(e)$ and $t(e)$ denote the initial and terminal vertices of edge $e \in \mathcal{E}$. The set Γ is defined by $(v, e) \in \Gamma$ if and only if $v = t(e)$. The map α in this case is $\alpha(v, e) = i(e)$. See chapter 4.3 in [4] for a discussion of the Mauldin-Williams graph.

Example 2.2. Let (T, d) be a bounded complete metric space. Assume that $T = \bigcup_{k=1}^p T_k$, where each T_k is a closed subset of T and $T_k \cap T_l = \emptyset$ for $k \neq l$. For $1 \leq i \leq m$, let $\theta_i : T \rightarrow T$ be a continuous map such that $\theta_i(T_k) \subset T_{\nu(i,k)}$, $1 \leq k \leq p$, where $\nu(i, k) \in \{1, 2, \dots, p\}$, and $\text{Lip}(\theta_i|_{T_k}) \leq c < 1$. In this case, we take $V = \{k : 1 \leq k \leq p\}$, $S_k = T_k$ for $1 \leq k \leq p$, $\mathcal{E} = \{i : 1 \leq i \leq m\}$, $\Gamma = V \times \mathcal{E}$. The map $\alpha(k, i) = \nu(i, k)$, $1 \leq k \leq p$, $1 \leq i \leq m$, and $\theta_{(k,i)} = \theta_i|_{T_k}$.

For $u \in V$, define

$$\Gamma_u = \{(v, e) \in \Gamma : \alpha(v, e) = u\}$$

and

$$E_u = \{e \in \mathcal{E} : (u, e) \in \Gamma\}.$$

For $n \geq 1$, define

$$\Gamma^{(n)} = \{[(v_1, e_1), \dots, (v_n, e_n)] : (v_i, e_i) \in \Gamma, \alpha(v_{i+1}, e_{i+1}) = v_i, 1 \leq i \leq n-1\}.$$

For $u \in V, n \geq 1$, define

$$\Gamma_u^{(n)} = \{[(v_1, e_1), \dots, (v_n, e_n)] \in \Gamma^{(n)} : \alpha(v_1, e_1) = u\}.$$

Define $V_\infty = \{u \in V : \Gamma_u^{(n)} \neq \emptyset \quad \forall n \geq 1\}$.

Before we prove the next theorem we need to recall the definition of the *Hausdorff metric*. Let (S, d) be a complete metric space. If $A \subset S$, we define the diameter of A by

$$\text{diam}(A) = \sup\{d(s, t) : s, t \in A\}.$$

We shall say that A is *bounded* if $\text{diam}(A) < \infty$. For $A \subset S$ and $s \in S$ we define

$$d(s, A) = \inf\{d(s, a) : a \in A\}.$$

If $A \subset S$ and $\delta > 0$, we define $N_\delta(A)$ by

$$N_\delta(A) = \{s \in S : d(s, A) < \delta\}.$$

If A and B are nonempty, closed, bounded subsets of S , we define

$$D(A, B) = \inf\{\delta > 0 : A \subset N_\delta(B) \text{ and } B \subset N_\delta(A)\}.$$

If $\mathcal{B}(S)$ denotes the collection of nonempty, closed, bounded subsets of S , then it follows that $(\mathcal{B}(S), D)$ is a metric space. The metric D is called the Hausdorff

metric. It is known (see [16], Exercise 7, pages 280-281) that if (S, d) is a complete metric space, then $(\mathcal{B}(S), D)$ is also a complete metric space.

Theorem 2.3. *Let $V, \mathcal{E}, \Gamma, \alpha$ be given as before, and $\theta_{(v,e)} : S_v \rightarrow S_{\alpha(v,e)}$ be a Lipschitz map with $\text{Lip}(\theta_{(v,e)}) \leq c < 1$ for all $(v, e) \in \Gamma$. Assume $\Gamma_u \neq \emptyset$ for all $u \in V$. Then there exists a unique list $(C_v)_{v \in V}$ of nonempty closed bounded sets $C_v \subset S_v$ such that*

$$(2.1) \quad C_u = \overline{\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(C_v)}$$

for all $u \in V$. Furthermore, C_v is indeed compact for all $v \in V$, so we may remove the closure in the above equation.

Proof. Let $\mathcal{B}(S_v)$ denote the collection of closed, bounded nonempty subsets of S_v with the Hausdorff metric D_v . Then since S_v is a complete metric space, we know that $\mathcal{B}(S_v)$ is a complete metric space. So the finite cartesian product $\prod_{v \in V} \mathcal{B}(S_v)$ with the sup metric is also a complete metric space. Define the map $\Theta : \prod_{v \in V} \mathcal{B}(S_v) \rightarrow \prod_{v \in V} \mathcal{B}(S_v)$ by

$$\Theta((A_v)_{v \in V}) = \left(\overline{\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(A_v)} \right)_{u \in V}$$

Note that for any $u \in V$, $\overline{\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(A_v)}$ is nonempty because Γ_u is nonempty by assumption, and A_v is nonempty for each $v \in V$. Also it is bounded because $\theta_{(v,e)}(A_v)$, being the image of a bounded set A_v under a Lipschitz map $\theta_{(v,e)}$, is bounded, and a finite union of bounded sets is bounded. Thus the map Θ is well defined.

We claim that Θ is a contraction map. Let $A = (A_v)_{v \in V}$ and $B = (B_v)_{v \in V}$ be in $\prod_{v \in V} \mathcal{B}(S_v)$. Then $D(A, B) = \max_{v \in V} D_v(A_v, B_v)$ and

$$D(\Theta(A), \Theta(B)) = \max_{u \in V} D_u \left(\overline{\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(A_v)}, \overline{\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(B_v)} \right).$$

Let $\delta > D(A, B)$, and take any $(v, e) \in \Gamma_u$, $a_v \in A_v$. Since $D_v(A_v, B_v) \leq D(A, B) < \delta$, there exists $b_v \in B_v$ such that $d_v(a_v, b_v) < \delta$. Then

$$d_u(\theta_{(v,e)}(a_v), \theta_{(v,e)}(b_v)) \leq c d_v(a_v, b_v) < c\delta.$$

This shows that $\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(A_v)$ is contained in a $c\delta$ -neighborhood of

$$\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(B_v).$$

Interchanging the roles of A_v and B_v yields a corresponding inclusion, so

$$D_u \left(\overline{\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(A_v)}, \overline{\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(B_v)} \right) \leq c\delta$$

for all $u \in V$. Hence, $D(\Theta(A), \Theta(B)) \leq c\delta$. Since $\delta > D(A, B)$ was arbitrary, we have proved that $D(\Theta(A), \Theta(B)) \leq cD(A, B)$.

Therefore we have a contraction map Θ on a complete metric space. By the contraction mapping theorem, Θ has a unique fixed point, say $(C_v)_{v \in V}$. Thus we have $C_u = \overline{\bigcup_{(v,e) \in \Gamma_u} \theta_{(v,e)}(C_v)}$.

To see that C_v is compact for all $v \in V$ let us restrict the map Θ to $\prod_{v \in V} \mathcal{K}(S_v)$, where $\mathcal{K}(S_v)$ denotes the collection of nonempty compact subsets of S_v with the Hausdorff metric D_v . It is a straightforward exercise to prove that $\prod_{v \in V} \mathcal{K}(S_v)$ is a closed subset of the complete metric space $\prod_{v \in V} \mathcal{B}(S_v)$, so $\prod_{v \in V} \mathcal{K}(S_v)$ is itself a complete metric space. Then Θ maps the complete metric space $\prod_{v \in V} \mathcal{K}(S_v)$ into itself, and is a contraction map as seen above. Thus Θ has a unique fixed point in $\prod_{v \in V} \mathcal{K}(S_v)$. The fixed point must be the same as $(C_v)_{v \in V}$, since otherwise the original map Θ would have two fixed points. Therefore C_v is compact for all $v \in V$, and the theorem is proved. \square

Remark 2.4. The assumption $\Gamma_u \neq \emptyset$ for all $u \in V$ in the above theorem may be too strong for some examples. A weaker assumption under which we can prove the existence of an invariant list is $V_\infty \neq \emptyset$. Note that $\Gamma_u \neq \emptyset$ for all $u \in V$ implies $V_\infty = V$, so $V_\infty \neq \emptyset$.

First we claim that $u \in V_\infty$ implies that there exists $v \in V_\infty$ with $(v, e) \in \Gamma_u$ for some $e \in \mathcal{E}$. Suppose not. Then for all $(v, e) \in \Gamma_u$, $v \notin V_\infty$. This implies, since V is a finite set, that there exists $n \geq 1$ such that $\Gamma_v^{(n)} = \emptyset$ for all $(v, e) \in \Gamma_u$. But since $u \in V_\infty$, there exists $[(v_1, e_1), (v_2, e_2), \dots, (v_{n+1}, e_{n+1})] \in \Gamma_u^{(n+1)}$, which implies $[(v_2, e_2), \dots, (v_{n+1}, e_{n+1})] \in \Gamma_{v_1}^{(n)}$ and since $\alpha(v_1, e_1) = u$, $(v_1, e_1) \in \Gamma_u$. This contradicts $\Gamma_v^{(n)} = \emptyset$ for all $(v, e) \in \Gamma_u$. Hence the claim.

Now consider the map $\Theta : \prod_{v \in V_\infty} \mathcal{B}(S_v) \rightarrow \prod_{v \in V_\infty} \mathcal{B}(S_v)$ by

$$\Theta((A_v)_{v \in V_\infty}) = \left(\overline{\bigcup_{\substack{(v,e) \in \Gamma_u \\ v \in V_\infty}} \theta_{(v,e)}(A_v)} \right)_{u \in V_\infty}$$

Note that Θ is well defined because of the above claim. Again by the contraction mapping theorem, we have $(C_v)_{v \in V_\infty}$, $C_v \subset S_v$ is compact such that

$$C_u = \bigcup_{\substack{(v,e) \in \Gamma_u \\ v \in V_\infty}} \theta_{(v,e)}(C_v)$$

Equivalently, under the assumption that $V_\infty \neq \emptyset$, all we have done is replaced V by $\hat{V} := V_\infty$, Γ by $\hat{\Gamma} := \{(v, e) \in \Gamma \mid v \in \hat{V}\}$ and α by $\hat{\alpha} := \alpha|_{\hat{\Gamma}}$, and applied Theorem 2.3.

Remark 2.5. Applying the previous result to Example 2.2, we get nonempty compact sets $C_k \subset T_k$ for $k \in V_\infty$ such that

$$C_k = \bigcup_{\substack{\nu(i,l)=k \\ l \in V_\infty}} \theta_i(C_l) \text{ for } k \in V_\infty.$$

If we let $C = \bigcup_{i \in V_\infty} C_i$, then C is a nonempty compact set and it satisfies

$$C = \bigcup_{i=1}^m \theta_i(C).$$

Thus we have a compact invariant set for the family of maps θ_i , $1 \leq i \leq m$.

Remark 2.6. We can relax the condition $\text{Lip}(\theta_{(v,e)}) \leq c < 1$ for all $(v,e) \in \Gamma$ in the previous theorem to the following weaker condition. Suppose for some fixed $n \geq 1$ the composition of any n of the maps $\theta_{(v,e)}$, whenever the composition is defined, is Lipschitz with Lipschitz constant $\leq c < 1$. Then it is easy to see that Θ^n is a contraction map on a complete metric space, where Θ is the map defined in the proof of the theorem. It is then well known that the map Θ has a unique fixed point. Thus the conclusion of the previous theorem holds under this weaker assumption.

3. PERRON-FROBENIUS OPERATORS

From now on, let $V = \{1, 2, \dots, p\}$ with S_1, S_2, \dots, S_p the corresponding complete metric spaces. We do not necessarily assume that S_j , $1 \leq j \leq p$, is compact. Let

$$X_i = C_b(S_i) = \{f : S_i \rightarrow \mathbb{R} : f \text{ is continuous and bounded}\}$$

for $1 \leq i \leq p$ with $\|f\| = \sup_{s \in S_i} |f(s)|$.

Define a linear map $A : X_1 \times X_2 \times \dots \times X_p \rightarrow X_1 \times X_2 \times \dots \times X_p$ by

$$(3.1) \quad (Af)_j(s) = \sum_{e \in E_j} b_{(j,e)}(s) f_{\alpha(j,e)}(\theta_{(j,e)}(s)) \quad \text{for } s \in S_j$$

where $f = (f_1, f_2, \dots, f_p)$ and the functions $b_{(j,e)} \in X_j$ are given. We assume throughout this section that $E_j = \{e \in \mathcal{E} : (j,e) \in \Gamma\}$ is nonempty for all $j \in V$.

Define for $M > 0, \lambda \geq 0, 1 \leq j \leq p$

$$(3.2) \quad K_j(M, \lambda) = \{f \in X_j : 0 \leq f(s) \leq f(t) \exp(M(d_j(s,t))^\lambda) \text{ for all } s, t \in S_j\}.$$

Remark 3.1. From the definition, it follows that if $f \in K_j(M, \lambda)$ and $f(t) = 0$ for some $t \in S_j$, then $f(s) = 0$ for all $s \in S_j$. Thus $f \in K_j(M, \lambda)$ implies that either f is identically zero on S_j or f is strictly positive on S_j .

If Y is a real Banach space, a closed set $K \subset Y$ is called a closed *cone* if $\lambda K + \mu K \subset K$ for all $\lambda \geq 0, \mu \geq 0$ and $K \cap (-K) = \{0\}$.

The following lemma follows by the same argument used in Lemma 5.4, p.89, in [19]. We give the proof for the reader's convenience.

Lemma 3.2. *Let $K_j := K_j(M, \lambda)$ be as defined by eq. (3.2) with $\lambda > 0$. Then K_j is a closed cone in $(X_j, \|\cdot\|)$, and $\{f \in K_j : \|f\| \leq 1\}$ is equicontinuous.*

Proof. It is easy to verify that K_j is a closed cone, and the proof is left to the reader. To prove the equicontinuity of $\{f \in K_j : \|f\| \leq 1\}$ let $f \in K_j$ with $\|f\| \leq 1$. We claim that for any $s, t \in S_j$ we have

$$|f(s) - f(t)| \leq M(d_j(s,t))^\lambda.$$

According to the previous remark, either f is identically zero on S_j or f is strictly positive on S_j . The inequality is obvious in the first case. In the later case, we may assume that $0 < f(s) \leq f(t) \leq 1$. The definition of K_j implies that

$$|\ln(f(s)) - \ln(f(t))| \leq M(d_j(s,t))^\lambda.$$

The mean value theorem implies that for some ξ with $\ln(f(s)) \leq \xi \leq \ln(f(t)) \leq 0$ we have

$$\begin{aligned} |f(s) - f(t)| &= \exp(\ln(f(t))) - \exp(\ln(f(s))) \\ &= \exp(\xi) |\ln(f(t)) - \ln(f(s))| \leq M(d_j(s,t))^\lambda. \end{aligned}$$

Since this is true for any $f \in K_j$ with $\|f\| \leq 1$, equicontinuity follows. \square

Lemma 3.3. *Assume for some $M_0 > 0$ and $\lambda > 0$, $b_{(j,e)} \in K_j(M_0, \lambda)$ for all $(j, e) \in \Gamma$. Then there exists $M > 0$ so that the map A defined above maps $\prod_{i=1}^p K_i(M, \lambda)$ into itself.*

Proof. Let $f_i \in K_i(M, \lambda)$ for $1 \leq i \leq p$ and $s, t \in S_j$. Then

$$(Af)_j(s) = \sum_{e \in E_j} b_{(j,e)}(s) f_{\alpha(j,e)}(\theta_{(j,e)}(s))$$

Since $b_{(j,e)} \in K_j(M_0, \lambda)$, $b_{(j,e)}(s) \leq b_{(j,e)}(t) \exp((M_0(d_j(s, t)))^\lambda)$. Also

$$\begin{aligned} f_{\alpha(j,e)}(\theta_{(j,e)}(s)) &\leq f_{\alpha(j,e)}(\theta_{(j,e)}(t)) \exp(M(d_{\alpha(j,e)}(\theta_{(j,e)}(s), \theta_{(j,e)}(t)))^\lambda) \\ &\leq f_{\alpha(j,e)}(\theta_{(j,e)}(t)) \exp(Mc^\lambda(d_j(s, t))^\lambda). \end{aligned}$$

Thus

$$\begin{aligned} (Af)_j(s) &= \sum_{e \in E_j} b_{(j,e)}(s) f_{\alpha(j,e)}(\theta_{(j,e)}(s)) \\ &\leq \sum_{e \in E_j} b_{(j,e)}(t) f_{\alpha(j,e)}(\theta_{(j,e)}(t)) \exp((M_0 + Mc^\lambda)(d_j(s, t))^\lambda) \\ &= (Af)_j(t) \exp((M_0 + Mc^\lambda)(d_j(s, t))^\lambda). \end{aligned}$$

So, if we choose M such that $M_0 + Mc^\lambda \leq M$, which can be done because $c < 1$ and $\lambda > 0$, then

$$(Af)_j(s) \leq (Af)_j(t) \exp(M(d_j(s, t))^\lambda),$$

so $(Af)_j \in K_j(M, \lambda)$ for $1 \leq j \leq p$. \square

We should note that observations similar to Lemma 3.3 have been made earlier by other authors. See the proof of Theorem 5.4 in [19] and [2], for example.

We shall use the following notations.

$$\bar{\Gamma}^{(n)} := \{[(j_1, e_1), \dots, (j_n, e_n)] : (j_i, e_i) \in \Gamma, 1 \leq i \leq n, \alpha(j_i, e_i) = j_{i+1}, 1 \leq i < n\}.$$

$$\bar{\Gamma}_j^{(n)} := \{[(j_1, e_1), \dots, (j_n, e_n)] \in \bar{\Gamma}^{(n)} : j_1 = j\}.$$

We shall also use (J, E) where $J = (j_1, \dots, j_n)$, $E = (e_1, \dots, e_n)$ as a shorthand notation for $[(j_1, e_1), \dots, (j_n, e_n)] \in \bar{\Gamma}^{(n)}$.

For $(J, E) = [(j_1, e_1), \dots, (j_n, e_n)] \in \bar{\Gamma}^{(n)}$, define for $s \in S_{j_1}$

$$b_{(J,E)}(s) := b_{(j_1, e_1)}(s) b_{(j_2, e_2)}(\theta_{(j_1, e_1)}(s)) \cdots b_{(j_n, e_n)}(\theta_{(j_{n-1}, e_{n-1})} \circ \cdots \circ \theta_{(j_1, e_1)}(s))$$

and

$$\theta_{(J,E)}(s) := \theta_{(j_n, e_n)} \circ \cdots \circ \theta_{(j_1, e_1)}(s).$$

Let us compute A^2 .

$$\begin{aligned} (A^2 f)_{j_1}(s) &= (A(Af))_{j_1}(s) \\ &= \sum_{e_1: (j_1, e_1) \in \Gamma} b_{(j_1, e_1)}(s) (Af)_{\alpha(j_1, e_1)}(\theta_{(j_1, e_1)}(s)) \end{aligned}$$

Using

$$(Af)_{\alpha(j_1, e_1)}(\theta_{(j_1, e_1)}(s)) = \sum_{\substack{e_2: (j_2, e_2) \in \Gamma \\ j_2 = \alpha(j_1, e_1)}} b_{(j_2, e_2)}(\theta_{(j_1, e_1)}(s)) f_{\alpha(j_2, e_2)}(\theta_{(j_2, e_2)}(\theta_{(j_1, e_1)}(s))),$$

we get

$$(A^2 f)_{j_1}(s) = \sum_{(J,E) \in \bar{\Gamma}_{j_1}^{(2)}} b_{(J,E)}(s) f_{\alpha(j_2, e_2)}(\theta_{(J,E)}(s)).$$

This computation suggests the formula for A^n given in the following lemma.

Lemma 3.4. *Let $n \geq 1$. Then for $f = (f_1, f_2, \dots, f_p) \in \prod_{i=1}^p X_i$ and $1 \leq j_1 \leq p$*

$$(3.3) \quad (A^n f)_{j_1}(s) = \sum_{(J,E) \in \bar{\Gamma}_{j_1}^{(n)}} b_{(J,E)}(s) f_{\alpha(j_n, e_n)}(\theta_{(J,E)}(s)), \quad s \in S_{j_1}$$

Also the operator norm of A^n is given by

$$(3.4) \quad \|A^n\| = \max_{1 \leq j \leq p} \sup_{s \in S_j} \sum_{(J,E) \in \bar{\Gamma}_j^{(n)}} b_{(J,E)}(s).$$

Proof. The equation for A^n follows by a simple induction on n . If $f \in \prod_{i=1}^p X_i$ with $\|f\| \leq 1$, i.e., $|f_j(s)| \leq 1 \forall s \in S_j, 1 \leq j \leq p$ then the equation for $(A^n f)_j(s)$ gives

$$|(A^n f)_j(s)| \leq \sum_{(J,E) \in \bar{\Gamma}_j^{(n)}} b_{(J,E)}(s)$$

Taking supremum over $s \in S_j$ and then maximum over $1 \leq j \leq p$ gives

$$\|A^n\| \leq \max_{1 \leq j \leq p} \sup_{s \in S_j} \sum_{(J,E) \in \bar{\Gamma}_j^{(n)}} b_{(J,E)}(s).$$

If we take $f = (f_1, f_2, \dots, f_p)$ where f_j is identically equal to one on S_j then

$$(A^n f)_j(s) = \sum_{(J,E) \in \bar{\Gamma}_j^{(n)}} b_{(J,E)}(s)$$

Therefore we get the equation for $\|A^n\|$. □

Lemma 3.5. *Let $(S_j)_{j=1}^p$ be bounded complete metric spaces, assume that $\Gamma_j := \{(k, e) \in \Gamma : \alpha(k, e) = j\}$ is nonempty for $1 \leq j \leq p$, and let $(C_j)_{j=1}^p$ be the unique invariant list of compact sets given by Theorem 2.3. Let $(J, E) = [(j_1, e_1), \dots, (j_n, e_n)] \in \bar{\Gamma}^{(n)}$ and $\theta_{(J,E)}(s) = \theta_{(j_n, e_n)} \circ \dots \circ \theta_{(j_1, e_1)}(s)$, $s \in S_{j_1}$. Then there exists $M_1 > 0$ such that for all $n \geq 1$*

$$d_{\alpha(j_n, e_n)}(\theta_{(J,E)}(s), C_{\alpha(j_n, e_n)}) \leq M_1 c^n \quad \forall s \in S_{j_1}$$

where $c < 1$ is the constant such that $\text{Lip}(\theta_{(j,e)}) \leq c$ for all $(j, e) \in \Gamma$.

Proof. Since the metric spaces (S_j) , $1 \leq j \leq p$ are bounded we can find M_1 so that $d_j(s, C_j) \leq M_1 \forall s \in S_j, 1 \leq j \leq p$. Let $(j, e) \in \Gamma$ and $s \in S_j$. Then we can find $t \in C_j$ such that $d_j(s, t) \leq M_1$. Since $\theta_{(j,e)}(t) \in C_{\alpha(j,e)}$,

$$d_{\alpha(j,e)}(\theta_{(j,e)}(s), C_{\alpha(j,e)}) \leq d_{\alpha(j,e)}(\theta_{(j,e)}(s), \theta_{(j,e)}(t)) \leq c d_j(s, t) \leq c M_1.$$

The result now follows easily by an induction on n . □

Let us recall the definition of Kuratowski's *measure of noncompactness* β . If (S, d) is a metric space and $B \subset S$ is a bounded set, then $\beta(B)$ is defined by

$$\beta(B) = \inf\{\delta > 0 : B = \cup_{j=1}^k B_j, k < \infty \text{ and } \text{diam}(B_j) \leq \delta \text{ for } 1 \leq j \leq k\}.$$

Suppose that K is a closed cone in a Banach space Y and $L : Y \rightarrow Y$ is a bounded linear map with $L(K) \subset K$. Define

$$\|L\|_K = \sup\{\|L(y)\| : y \in K, \|y\| \leq 1\}.$$

Define $r_K(L)$, the *cone spectral radius* of L , and $\sigma_K(L)$, by

$$r_K(L) := \lim_{n \rightarrow \infty} \|L^n\|_K^{1/n} \text{ and}$$

$$\sigma_K(L) := \limsup_{n \rightarrow \infty} (\beta(L^n(U)))^{1/n}$$

where $U = \{y \in K : \|y\| \leq 1\}$.

It is a special case of Theorem 4.10 in [10] that if $\sigma_K(L) < r_K(L)$, then there exists $y \in K \setminus \{0\}$ with $L(y) = ry$, $r = r_K(L)$. (Note that the definition in [10] of $\rho_K(L)$, the cone essential spectral radius of L , satisfies $\rho_K(L) \leq \sigma_K(L)$. The definition of $\rho_K(L)$ in [10] differs from that in [9] and [18]. It is shown in [10] that the earlier definition has some serious deficiencies.) We shall use this result to prove the existence of a non-zero eigenvector for the map A given by (3.1). Alternately, if all the sets S_j were compact, an argument similar to the one used to prove Theorem 5.4 in [19] would give the following theorem.

Theorem 3.6. *Consider the map A defined on $\prod_{j=1}^p C_b(S_j)$ by*

$$(Af)_j(s) = \sum_{e \in E_j} b_{(j,e)}(s) f_{\alpha(j,e)}(\theta_{(j,e)}(s)) \quad \text{for } s \in S_j$$

where $f = (f_1, f_2, \dots, f_p)$. Assume that S_j is a bounded, complete metric space for $1 \leq j \leq p$. Assume also that $\Gamma_j \neq \emptyset$ and $E_j \neq \emptyset$ for $1 \leq j \leq p$ and that for some $M_0 > 0$ and $\lambda > 0$, $b_{(j,e)} \in K_j(M_0, \lambda)$ for all $(j, e) \in \Gamma$. Let K be the cone $\prod_{j=1}^p K_j(M, \lambda)$, where $M_0 + c^\lambda M \leq M$. Then $\|A^n\|_K = \|A^n\|$ for all $n \geq 1$, where $\|A^n\|$ is given by equation (3.4), and $r_K(A) = \lim_{n \rightarrow \infty} \|A^n\|^{1/n}$. If $r_K(A) > 0$, there exists $u = (u_1, u_2, \dots, u_p) \in K \setminus \{0\}$ with $Au = ru$, where $r = r_K(A)$. If $b_{(j,e)}(s) > 0$ for all $(j, e) \in \Gamma$ and all $s \in S_j$, then $r_K(A) > 0$.

Proof. It is enough to show that $\sigma_K(A) < r_K(A)$. Let $(C_j)_{j=1}^p$ be the unique invariant list of nonempty compact sets given by Theorem 2.3 and let $C = \prod_{j=1}^p C_j$. Let $\mathcal{U} = \{f \in K : \|f\| \leq 1\}$. From Lemma 3.2, we know that \mathcal{U} is equicontinuous. Let us write $\mathcal{U}|_C = \{f|_C : f \in \mathcal{U}\}$. Then $\mathcal{U}|_C$ is a bounded equicontinuous family of functions from the compact set C into \mathbb{R}^p . So, by Ascoli's theorem, it is totally bounded. Therefore, given $\epsilon > 0$, we can write $\mathcal{U} = \bigcup_{l=1}^q \mathcal{U}_l$, $q < \infty$, such that $\|f|_C - g|_C\| \leq \epsilon$ provided f and g are in the same \mathcal{U}_l . Let $f, g \in \mathcal{U}$ and $1 \leq j \leq p$. Then using Lemma 3.4, we have for $s \in S_j$

$$|(A^n f)_j(s) - (A^n g)_j(s)| \leq \sum_{(J,E) \in \bar{\Gamma}_j^{(n)}} b_{(J,E)}(s) |f_{\alpha(j_n, e_n)}(\theta_{(J,E)}(s)) - g_{\alpha(j_n, e_n)}(\theta_{(J,E)}(s))|.$$

Using Lemma 3.5, there exists $\tau \in C_{\alpha(j_n, e_n)}$ with $d(\theta_{(J,E)}(s), \tau) \leq M_1 c^n$.

Since $f_{\alpha(j_n, e_n)} \in K_{\alpha(j_n, e_n)}(M, \lambda)$ and $\|f\| \leq 1$, we have as in the proof of Lemma 3.2

$$|f_{\alpha(j_n, e_n)}(\theta_{(J,E)}(s)) - f_{\alpha(j_n, e_n)}(\tau)| \leq M d(\theta_{(J,E)}(s), \tau)^\lambda \leq M(M_1 c^n)^\lambda.$$

The same is true for g . Also, if we assume that $f, g \in \mathcal{U}_l$ for some l , $1 \leq l \leq q$ then $|f_{\alpha(j_n, e_n)}(\tau) - g_{\alpha(j_n, e_n)}(\tau)| \leq \epsilon$. Therefore, by triangle inequality

$$|f_{\alpha(j_n, e_n)}(\theta_{(J,E)}(s)) - g_{\alpha(j_n, e_n)}(\theta_{(J,E)}(s))| \leq \epsilon + 2M(M_1 c^n)^\lambda \text{ if } f, g \in \mathcal{U}_l.$$

So, if $f, g \in \mathcal{U}_l$ and $1 \leq j \leq p$,

$$|(A^n f)_j(s) - (A^n g)_j(s)| \leq (\epsilon + 2M(M_1 c^n)^\lambda) \sum_{(J,E) \in \bar{\Gamma}_j^{(n)}} b_{(J,E)}(s).$$

Taking supremum over $s \in S_j$ and max over $1 \leq j \leq p$ and using equation (3.4), we get

$$\|A^n f - A^n g\| \leq (\epsilon + 2M(M_1 c^n)^\lambda) \|A^n\|$$

for $f, g \in \mathcal{U}_l$, $1 \leq l \leq q$.

Thus $A^n(\mathcal{U}) = \bigcup_{l=1}^q A^n(\mathcal{U}_l)$ with

$$\text{diam}(A^n(\mathcal{U}_l)) \leq (\epsilon + 2M(M_1 c^n)^\lambda) \|A^n\|$$

So, $\beta(A^n(\mathcal{U})) \leq (\epsilon + 2M(M_1 c^n)^\lambda) \|A^n\|$.

Since $\epsilon > 0$ was arbitrary, $\beta(A^n(\mathcal{U})) \leq 2M(M_1 c^n)^\lambda \|A^n\|$ which implies

$$(\beta(A^n(\mathcal{U})))^{\frac{1}{n}} \leq (2MM_1^\lambda)^{\frac{1}{n}} c^\lambda \|A^n\|^{\frac{1}{n}}.$$

In general, it is obviously true that $\|A^n\|_K \leq \|A^n\|$. On the other hand, if $f = (f_1, f_2, \dots, f_p)$ and $f_j(s) = 1$ for all $s \in S_j$, then $f \in K$; and we have seen in the proof of Lemma 3.4 that $\|A^n(f)\| = \|A^n\|$. It follows that $\|A^n\|_K = \|A^n\|$ for all $n \geq 1$ and that $r_K(A) = \lim_{n \rightarrow \infty} \|A^n\|^{\frac{1}{n}} = r(A)$, where $\|A^n\|$ is given by equation (3.4) and $r(A)$ denotes the spectral radius of A . Taking the limit in our estimate for $(\beta(A^n(\mathcal{U})))^{\frac{1}{n}}$ gives

$$\sigma_K(A) = \lim_{n \rightarrow \infty} (\beta(A^n(\mathcal{U})))^{\frac{1}{n}} \leq (c^\lambda) r_K(A).$$

If $r_K(A) > 0$, it follows (because $0 < c < 1$) that $\sigma_K(A) < r_K(A)$, and we are done. If we assume that $b_{(j,e)}(s) > 0$ for all $(j,e) \in \Gamma$ and $s \in S_j$, then because $b_{(j,e)} \in K_j(M_0, \lambda)$, there exists $\delta > 0$ such that $b_{(j,e)}(s) \geq \delta$ for all $(j,e) \in \Gamma$ and $s \in S_j$, and it follows easily that $r_K(A) \geq \delta > 0$. \square

Remark 3.7. Suppose $u = (u_1, u_2, \dots, u_p)$ is a nonzero eigenvector of the linear map A with eigenvalue $r(A)$ given by Theorem 3.6. Then for any $1 \leq j \leq p$, either u_j is identically zero or u_j is strictly positive on S_j . To see this note that $u_j \in K_j(M, \lambda)$, so

$$0 \leq u_j(s) \leq u_j(t) \exp(M(d_j(s,t))^\lambda) \text{ for all } s, t \in S_j.$$

Thus $u_j(t) = 0$ for some $t \in S_j$ will imply that $u_j(s) = 0$ for all $s \in S_j$. Also, since u is nonzero, at least one of the coordinate functions u_j is strictly positive.

Remark 3.8. In general (in the context of the Krein-Rutman Theorem), if $r_K(L) = 0$, it need not be true that there exist $v \in K \setminus \{0\}$ with $L(v) = 0$. Suppose, however, that K and A are as in Theorem 3.6, that $b_{(j,e)} \in K_j(M_0, \lambda)$ for all $(j,e) \in \Gamma$ and that $r_K(A) = 0$. We claim that there exists an integer N such that $A^N = 0$ and that there exists $v \in K \setminus \{0\}$ with $A(v) = 0$. Because S_j is bounded and $b_{(j,e)} \in K_j(M_0, \lambda)$, we have already seen that either $b_{(j,e)}(s) = 0$ for all $s \in S_j$ or there exists $\delta_{(j,e)} > 0$ with $b_{(j,e)}(s) \geq \delta_{(j,e)}$ for all $s \in S_j$. Let $\mathcal{P} = \{(j,e) \in \Gamma : b_{(j,e)}(s) > 0 \text{ for all } s \in S_j\}$. Because \mathcal{P} is a finite set (since Γ is finite), there exists $\delta > 0$ with $b_{(j,e)}(s) \geq \delta$ for all $s \in S_j$ and for all $(j,e) \in \mathcal{P}$. For $n \geq 1$, define $\bar{\mathcal{P}}^{(n)} \subset \bar{\Gamma}^{(n)}$ by

$$\bar{\mathcal{P}}^{(n)} = \{(J, E) \in \bar{\Gamma}^{(n)} : (j_k, e_k) \in \mathcal{P} \text{ for } 1 \leq k \leq n\}.$$

If $\bar{\mathcal{P}}^{(n)}$ is nonempty, it easily follows from equation (3.4) that $\|A^n\| \geq \delta^n$, so if $\bar{\mathcal{P}}^{(n)}$ is nonempty for all $n \geq 1$, $r_K(A) \geq \delta$, contrary to our assumption. Thus there must exist an integer N such that $\bar{\mathcal{P}}^{(n)}$ is empty for all $n \geq N$. However, if $(J, E) \in \bar{\Gamma}^{(n)}$ and $(J, E) \notin \bar{\mathcal{P}}^{(n)}$, $b_{(J,E)}(s) = 0$ for all $s \in S_j$, so we find that $A^n = 0$ for all $n \geq N$. If $w \in K \setminus \{0\}$, let $p \leq N$ be the least positive integer such that $A^p(w) = 0$. If we define $v = A^{p-1}(w) \in K \setminus \{0\}$, $A(v) = 0$.

Remark 3.9. If β denotes the Kuratowski measure of noncompactness on a Banach space X , K denotes a closed cone in X and $\Lambda : X \rightarrow X$ is a bounded linear map such that $\Lambda(K) \subset K$, define $\beta_K(\Lambda)$ by

$$\beta_K(\Lambda) := \inf\{\lambda > 0 : \beta(\Lambda(T)) \leq \lambda\beta(T) \text{ for all bounded sets } T \subset K\}.$$

If $L : X \rightarrow X$ is a bounded linear map such that $L(K) \subset K$, one can prove that $\lim_{n \rightarrow \infty} (\beta_K(L^n))^{\frac{1}{n}}$ exists. If $\lim_{n \rightarrow \infty} (\beta_K(L^n))^{\frac{1}{n}} < r_K(L)$, Theorem 2.2 in [18] implies that there exists $u \in K \setminus \{0\}$ with $L(u) = ru$ and $r := r_K(L)$.

One might hope that Theorem 2.2 in [18] could be used to prove Theorem 3.6. However, if the metric spaces S_j , $1 \leq j \leq p$, in Theorem 3.6 are not compact, Theorem 2.2 in [18] is, in general, not applicable, even in very simple special cases. To illustrate this we work in the Hilbert space $l^2(\mathbb{N})$, and we let $\{e_j | j \geq 1\}$ denote the usual orthonormal basis of $l^2(\mathbb{N})$. In the notation of Theorem 3.6, we take $S_1 = \{x \in l^2(\mathbb{N}) : \|x\| \leq 1\}$, $V = \{1\}$, \mathcal{E} to be a set with exactly one point. Let $X_1 = C_b(S_1)$ and, in the notation of Lemma 3.2, let $K = K(M, 1)$, where M is a fixed constant with $M \geq 1$. Define a bounded linear operator $A : X_1 \rightarrow X_1$ by

$$(Af)(x) = f\left(\frac{1}{2}x\right)$$

for $x \in S_1$. Obviously, A is a trivial example of the operators considered in Theorem 3.6, and $A(K(M, 1)) \subset K(M, 1)$ and $Au = u$, where $u \in K(M, 1)$ denotes the function identically equal to 1. However, we claim that for each integer $n \geq 1$, there is a bounded set $T_n \subset K(M, 1)$ such that $\beta(A^n(T_n)) = \beta(T_n) = (\frac{1}{2})^n$; and this implies that $\lim_{n \rightarrow \infty} (\beta_K(A^n))^{\frac{1}{n}} = 1 = r_K(A)$, so Theorem 2.2 of [18] cannot be used (at least with the cones $K(M, 1)$) even in this trivial situation.

To construct the sets T_n , for each integer $n \geq 1$ define $\phi_n : [0, 1] \rightarrow [1, 1 + 2^{-n}]$ by $\phi_n(s) = 1 + 2^{-n}$ for $2^{-n} \leq s \leq 1$ and $\phi_n(s) = 1 + s$ for $0 \leq s \leq 2^{-n}$, so ϕ_n is Lipschitz with Lipschitz constant 1. Let $\langle \cdot, \cdot \rangle$ denote the inner product on $l^2(\mathbb{N})$, and define $\psi_{j,n}(x)$ and T_n by

$$\psi_{j,n}(x) = \phi_n(|\langle x, e_j \rangle|) \text{ and } T_n = \{\psi_{j,n} : j \geq 1\}.$$

The reader can verify that $T_n \subset K(M, 1)$ for all $n \geq 1$ and, that, for a fixed $n \geq 1$, and for $1 \leq j < k < \infty$,

$$\|\psi_{j,n} - \psi_{k,n}\| = \sup\{|\psi_{j,n}(x) - \psi_{k,n}(x)| : x \in S_1\} = |\psi_{j,n}(e_j) - \psi_{k,n}(e_j)| = 2^{-n}.$$

Since $(A^n \psi_{j,n})(x) = \phi_n(2^{-n}|\langle x, e_j \rangle|)$, the same argument shows that for a fixed $n \geq 1$ and for $1 \leq j < k < \infty$,

$$\|A^n(\psi_{j,n}) - A^n(\psi_{k,n})\| = |\psi_{j,n}(2^{-n}e_j) - \psi_{k,n}(2^{-n}e_j)| = 2^{-n}.$$

Using these equations, it is easy to verify that $\beta(T_n) = \beta(A^n(T_n)) = 2^{-n}$.

4. HAUSDORFF DIMENSION

Recall that a metric space (S, d) is called *perfect* if every point of S is a limit point of S , i.e., for each $s \in S$, there exists a sequence $(s_k)_k$ in S such that $s_k \neq s$ for all k and $s_k \rightarrow s$ as $k \rightarrow \infty$.

Let (S_1, d_1) be a perfect metric space and (S_2, d_2) a metric space. A map $\theta : S_1 \rightarrow S_2$ is said to be an *infinitesimal similitude* at $s \in S_1$ if for any sequences $(s_k)_k$ and $(t_k)_k$ in S_1 with $s_k \neq t_k$ for $k \geq 1$ and $s_k \rightarrow s, t_k \rightarrow s$, the limit

$$(4.1) \quad \lim_{k \rightarrow \infty} \frac{d_2(\theta(s_k), \theta(t_k))}{d_1(s_k, t_k)} =: (D\theta)(s)$$

exists and is independent of the particular sequences $(s_k)_k$ and $(t_k)_k$. We shall say that θ is an infinitesimal similitude on S_1 if θ is an infinitesimal similitude at s for all $s \in S_1$. Notice that the assumption that S_1 is perfect implies that for every $s \in S_1$, there exist sequences $(s_k)_k$ and $(t_k)_k$ as above.

We list some basic properties of infinitesimal similitudes that we shall need.

Lemma 4.1. *Let (S_1, d_1) be a perfect metric space and (S_2, d_2) a metric space. If $\theta : S_1 \rightarrow S_2$ is an infinitesimal similitude, then $s \mapsto (D\theta)(s)$ is continuous.*

Proof. We argue by contradiction and assume that $s \mapsto (D\theta)(s)$ is not continuous. Then there exist $\epsilon > 0$ and $s \in S_1$ and a sequence $(s_k)_k$ in S_1 with $d_1(s_k, s) \rightarrow 0$ as $k \rightarrow \infty$ such that

$$|(D\theta)(s_k) - (D\theta)(s)| > \epsilon > 0.$$

Since S_1 is perfect and θ is an infinitesimal similitude, for each $k \geq 1$, there exist t_k and w_k in S_1 with $t_k \neq w_k, 0 < d_1(s_k, t_k) < \frac{1}{k}, 0 < d_1(s_k, w_k) < \frac{1}{k}$ and

$$\left| \frac{d_2(\theta(t_k), \theta(w_k))}{d_1(t_k, w_k)} - (D\theta)(s_k) \right| < \frac{\epsilon}{4}.$$

Since $s_k \rightarrow s$ as $k \rightarrow \infty$, it follows that $t_k \rightarrow s$ and $w_k \rightarrow s$ as $k \rightarrow \infty$ as well. So, by definition,

$$\frac{d_2(\theta(t_k), \theta(w_k))}{d_1(t_k, w_k)} \rightarrow (D\theta)(s) \text{ as } k \rightarrow \infty.$$

So, for k large enough, $|(D\theta)(s_k) - (D\theta)(s)| < \frac{\epsilon}{2}$, a contradiction. Hence, $s \mapsto (D\theta)(s)$ is continuous. \square

The following lemma states an analogue of the chain rule for infinitesimal similitudes.

Lemma 4.2. *Let $(S_j, d_j), j = 1, 2$, be perfect metric spaces and let (S_3, d_3) be a metric space. Let $\theta : S_1 \rightarrow S_2$ and $\psi : S_2 \rightarrow S_3$ be given. If θ is an infinitesimal similitude at $s \in S_1$ and ψ is an infinitesimal similitude at $\theta(s) \in S_2$, then $\psi \circ \theta$ is an infinitesimal similitude at s and*

$$(4.2) \quad (D(\psi \circ \theta))(s) = (D\psi)(\theta(s))(D\theta)(s).$$

Proof. Let $(s_k)_k$ and $(t_k)_k$ be sequences in S_1 with $s_k \neq t_k, k \geq 1, s_k \rightarrow s, t_k \rightarrow s$ as $k \rightarrow \infty$. Then $\theta(s_k) \rightarrow \theta(s), \theta(t_k) \rightarrow \theta(s)$ as $k \rightarrow \infty$. We consider two cases.

Case I. Assume that $(D\theta)(s) \neq 0$. We claim that there exists a positive integer k_0 with $\theta(s_k) \neq \theta(t_k)$ for all $k \geq k_0$. If not, there exists a subsequence $k_i \rightarrow \infty$ such that $\theta(s_{k_i}) = \theta(t_{k_i})$ for $i \geq 1$. Writing $\sigma_i = s_{k_i}$ and $\tau_i = t_{k_i}$, we have that

$\sigma_i \rightarrow s$, $\tau_i \rightarrow s$, $\sigma_i \neq \tau_i$ and $(D\theta)(s) = \lim_{i \rightarrow \infty} \frac{d_2(\theta(\sigma_i), \theta(\tau_i))}{d_1(\sigma_i, \tau_i)} = 0$, which contradicts our assumption. It follows that, for $k \geq k_0$, we can write

$$\frac{d_3(\psi(\theta(s_k)), \psi(\theta(t_k)))}{d_1(s_k, t_k)} = \frac{d_3(\psi(\theta(s_k)), \psi(\theta(t_k)))}{d_2(\theta(s_k), \theta(t_k))} \frac{d_2(\theta(s_k), \theta(t_k))}{d_1(s_k, t_k)}.$$

As $k \rightarrow \infty$, the limit of the right hand side exists and equals $(D\psi)(\theta(s))(D\theta)(s)$, so the limit of the left hand side exists and equation (4.2) is satisfied.

Notice that if s_k and t_k are sequences with $s_k \rightarrow s$, $t_k \rightarrow s$, $s_k \neq t_k$ for all k and $\theta(s_k) \neq \theta(t_k)$ for all $k \geq k_0$, then the argument above proves that

$$\lim_{k \rightarrow \infty} \frac{d_3(\psi(\theta(s_k)), \psi(\theta(t_k)))}{d_1(s_k, t_k)} = (D\psi)(\theta(s))(D\theta)(s),$$

even if $(D\theta)(s) = 0$.

Case II. Assume that $(D\theta)(s) = 0$. Let s_k and t_k be sequences in S_1 with $s_k \rightarrow s$, $t_k \rightarrow s$ and $s_k \neq t_k$ for all $k \geq 1$. If there exists $k_0 \geq 1$ such that $\theta(s_k) \neq \theta(t_k)$ for all $k \geq k_0$, the argument above shows that

$$\lim_{k \rightarrow \infty} \frac{d_3(\psi(\theta(s_k)), \psi(\theta(t_k)))}{d_1(s_k, t_k)} = (D\psi)(\theta(s))(D\theta)(s) = 0.$$

If there exists k_1 such that $\theta(s_k) = \theta(t_k)$ for all $k \geq k_1$, we certainly have that

$$\lim_{k \rightarrow \infty} \frac{d_3(\psi(\theta(s_k)), \psi(\theta(t_k)))}{d_1(s_k, t_k)} = 0.$$

Thus we can assume that $K_1 := \{k \geq 1 : \theta(s_k) \neq \theta(t_k)\}$ and $K_2 := \{k \geq 1 : \theta(s_k) = \theta(t_k)\}$ are infinite sets. However, our previous argument (Case I) shows that

$$\lim_{k \rightarrow \infty, k \in K_1} \frac{d_3(\psi(\theta(s_k)), \psi(\theta(t_k)))}{d_1(s_k, t_k)} = (D\psi)(\theta(s))(D\theta)(s) = 0,$$

and it is clear that

$$\lim_{k \rightarrow \infty, k \in K_2} \frac{d_3(\psi(\theta(s_k)), \psi(\theta(t_k)))}{d_1(s_k, t_k)} = 0,$$

so we conclude that

$$\lim_{k \rightarrow \infty} \frac{d_3(\psi(\theta(s_k)), \psi(\theta(t_k)))}{d_1(s_k, t_k)} = 0.$$

□

The following lemma gives a “mean value theorem” and will be crucial in the proof of the main theorem.

Lemma 4.3. *Suppose that (S_1, d_1) and (S_2, d_2) are bounded, complete metric spaces, that (S_1, d_1) is perfect and that $\theta : S_1 \rightarrow S_2$ is an infinitesimal similitude. Also, assume that θ is Lipschitz and that $(D\theta)(s) > 0$ for all $s \in S_1$. Let $K \subset S_1$ be a compact, nonempty set. Then for each $\mu > 1$, there exists an open neighborhood U_μ of K and a positive number $\epsilon = \epsilon(\mu)$ such that for every $s, t \in U_\mu$ with $0 < d_1(s, t) < \epsilon(\mu)$,*

$$\mu^{-1}(D\theta)(s) \leq \frac{d_2(\theta(s), \theta(t))}{d_1(s, t)} \leq \mu(D\theta)(s)$$

Proof. For $(s, t) \in S_1 \times S_1$ with $s \neq t$, define $F(s, t) = \frac{d_2(\theta(s), \theta(t))}{d_1(s, t)}$. If $(s, s) \in S_1 \times S_1$, define $F(s, s) = (D\theta)(s)$. Because we assume that θ is Lipschitz on S_1 , there is a constant M_1 with $F(s, t) \leq M_1$ for all $(s, t) \in S_1 \times S_1$.

We claim that F is continuous on $S_1 \times S_1$. It suffices to prove that if $(s_k, t_k) \rightarrow (s, s)$, then $F(s_k, t_k) \rightarrow F(s, s) = (D\theta)(s)$. If $s_k \neq t_k$ for all $k \geq k_0$, we know that $F(s_k, t_k) \rightarrow (D\theta)(s)$ by the definition of $(D\theta)(s)$. If $s_k = t_k$ for all $k \geq k_1$, $F(s_k, t_k) = (D\theta)(s_k)$ for $k \geq k_1$, and Lemma 4.1 implies that $(D\theta)(s_k) \rightarrow (D\theta)(s)$. Thus we can assume that $J_1 := \{k : s_k \neq t_k\}$ and $J_2 := \{k : s_k = t_k\}$ are infinite sets. But in this case, the same reasoning implies that $\lim_{k \rightarrow \infty, k \in J_1} F(s_k, t_k) = (D\theta)(s)$ and $\lim_{k \rightarrow \infty, k \in J_2} F(s_k, t_k) = (D\theta)(s)$, so $\lim_{k \rightarrow \infty} F(s_k, t_k) = (D\theta)(s)$.

Lemma 4.1 implies that $s \mapsto (D\theta)(s)$ is continuous on S_1 , and $(s, t) \mapsto F(s, t)$ is continuous on $S_1 \times S_1$. Thus, since $(D\theta)(s) > 0$, if we define $G(s, t)$ by $G(s, t) = \frac{F(s, t)}{(D\theta)(s)}$, $(s, t) \mapsto G(s, t)$ is continuous on $S_1 \times S_1$ and $G(s, s) = 1$. Since $K \times K$ is compact, $G|_{K \times K}$ is uniformly continuous, so given $\mu > 1$, there exists $\epsilon(\mu) > 0$ with

$$\mu^{-1} < G(s, t) < \mu$$

for all $(s, t) \in K \times K$ with $d_1(s, t) \leq \epsilon(\mu)$. We claim that there exists an open neighborhood U_μ of K such that for all $s, t \in U_\mu$ with $d_1(s, t) \leq \epsilon(\mu)$, $\mu^{-1} < G(s, t) < \mu$. We argue by contradiction and suppose not. For m a positive integer, let $V_m = \{s \in S_1 : d_1(s, K) < \frac{1}{m}\}$. By assumption, there exist $s_m, t_m \in V_m$ with $d_1(s_m, t_m) \leq \epsilon(\mu)$ and $G(s_m, t_m) < \mu^{-1}$ or $G(s_m, t_m) > \mu$. Because $d_1(s_m, K) \rightarrow 0$ and $d_1(t_m, K) \rightarrow 0$, we can, by taking a subsequence, assume that $s_m \rightarrow s \in K$ and $t_m \rightarrow t \in K$ and $d_1(s, t) \leq \epsilon(\mu)$. By continuity of G , we either have $G(s, t) \leq \mu^{-1}$ or $G(s, t) \geq \mu$. However, because $s, t \in K$ and $d_1(s, t) \leq \epsilon(\mu)$, $\mu^{-1} < G(s, t) < \mu$, a contradiction. Thus an open set U_μ exists and, in fact, we can take $U_\mu = V_m$ for some $m \geq 1$. \square

Remark 4.4. For each $\epsilon > 0$, define $\mu(\epsilon) \geq 1$ to be the infimum of numbers $\mu > 1$ such that $\mu^{-1}(D\theta)(s) \leq \frac{d_2(\theta(s), \theta(t))}{d_1(s, t)} \leq \mu(D\theta)(s)$ for $s, t \in K$ with $0 < d_1(s, t) \leq \epsilon$. Lemma 4.3 implies that $\lim_{\epsilon \rightarrow 0+} \mu(\epsilon) = 1$, and clearly $\mu(\epsilon)$ is an increasing function of ϵ for $\epsilon > 0$.

Throughout the remainder of this section we shall make the following assumption.

H4.1 Let $V = \{1, 2, \dots, p\}$ and S_1, S_2, \dots, S_p be bounded, complete, perfect metric spaces. Let \mathcal{E} be a finite set, $\Gamma \subset V \times \mathcal{E}$ and $\alpha : \Gamma \rightarrow V$. For each $(j, e) \in \Gamma$, $\theta_{(j, e)} : S_j \rightarrow S_{\alpha(j, e)}$ is a Lipschitz map with $\text{Lip}(\theta_{(j, e)}) \leq c < 1$. Also, $\Gamma_i = \{(j, e) \in \Gamma : \alpha(j, e) = i\} \neq \emptyset$ for $1 \leq i \leq p$ and $E_j = \{e \in \mathcal{E} : (j, e) \in \Gamma\} \neq \emptyset$ for $1 \leq j \leq p$.

If H4.1 is satisfied then Theorem 2.3 implies that there exists a unique list of nonempty compact sets $C_j \subset S_j$, $1 \leq j \leq p$ with

$$(4.3) \quad C_i = \bigcup_{(j, e) \in \Gamma_i} \theta_{(j, e)}(C_j) \quad \text{for } 1 \leq i \leq p.$$

We shall further assume the following.

H4.2 For each $(j, e) \in \Gamma$, the map $\theta_{(j, e)} : S_j \rightarrow S_{\alpha(j, e)}$, given in H4.1, is an infinitesimal similitude and $(D\theta_{(j, e)})(s) > m > 0$ for all $s \in S_j$.

Notice that since $\theta_{(j, e)}$ is Lipschitz with $\text{Lip}(\theta_{(j, e)}) \leq c$, if $\theta_{(j, e)}$ is an infinitesimal similitude, we have $(D\theta_{(j, e)})(s) \leq c$ for all $s \in S_j$.

Assume that H4.1 and H4.2 are satisfied. For $\sigma \geq 0$, define

$$L_\sigma : \prod_{j=1}^p C_b(S_j) \rightarrow \prod_{j=1}^p C_b(S_j)$$

by

$$(4.4) \quad (L_\sigma f)_j(s) = \sum_{e \in E_j} ((D\theta_{(j,e)})(s))^\sigma f_{\alpha(j,e)}(\theta_{(j,e)}(s)) \text{ for } s \in S_j, 1 \leq j \leq p.$$

Recall that a map $f : (S, d) \rightarrow \mathbb{R}$ is said to be *Hölder continuous with Hölder exponent* $\lambda > 0$ if there exists a constant $C \in (0, \infty)$ such that

$$|f(s) - f(t)| \leq C(d(s, t))^\lambda \text{ for all } s, t \in S.$$

Let us assume the following.

H4.3 For each $(j, e) \in \Gamma$, the map $s \mapsto (D\theta_{(j,e)})(s)$ is Hölder continuous with Hölder exponent $\lambda > 0$, where λ is independent of $(j, e) \in \Gamma$.

Lemma 4.5. *If H4.1, H4.2 and H4.3 hold, and $\sigma \geq 0$, then the map*

$$s \mapsto ((D\theta_{(j,e)})(s))^\sigma$$

is in the cone $K_j(M_0, \lambda)$ defined by eq. (3.2) for some $M_0 > 0$ (depending on σ).

Proof. Fix $(j, e) \in \Gamma$ and $\sigma \geq 0$. Let $f(s) = (D\theta_{(j,e)})(s)$. The hypotheses H4.1 and H4.2 imply that $0 < m < f(s) \leq c$. By H4.3

$$|f(s) - f(t)| \leq C(d_j(s, t))^\lambda \text{ for all } s, t \in S_j.$$

Let $s, t \in S_j$. By the mean value theorem, there exists ξ between $f(s)$ and $f(t)$ such that

$$|\ln(f(s)) - \ln(f(t))| = \frac{1}{\xi} |f(s) - f(t)| \leq \frac{1}{\xi} C(d_j(s, t))^\lambda \leq \frac{1}{m} C(d_j(s, t))^\lambda$$

So, $f(s) \leq f(t) \exp(\frac{1}{m} C(d_j(s, t))^\lambda)$ which implies

$$f(s)^\sigma \leq f(t)^\sigma \exp(M_0(d_j(s, t))^\lambda) \text{ where } M_0 = \frac{\sigma C}{m}.$$

This completes the proof. \square

Now applying Theorem 3.6 to the linear map L_σ defined in (4.4), we get an eigenvector $u_\sigma \in K \setminus \{0\}$ with $L_\sigma u_\sigma = r(L_\sigma) u_\sigma$ and $r(L_\sigma) > 0$.

Lemma 4.6. *The map $\sigma \mapsto r(L_\sigma)$ is continuous and strictly decreasing. Furthermore, there is a unique $\sigma_0 \geq 0$ such that $r(L_{\sigma_0}) = 1$.*

Proof. Let u_σ be the positive eigenvector of L_σ with eigenvalue $r(L_\sigma)$. Let us write $b_{(j,e)}(t) = (D\theta_{(j,e)})(t)$ for $(j, e) \in \Gamma$. We know that $0 < m \leq b_{(j,e)}(t) \leq c < 1$ for all $t \in S_j$. Let $0 \leq \sigma < \sigma'$. Then

$$(b_{(j,e)}(t))^{\sigma'} = (b_{(j,e)}(t))^{\sigma' - \sigma} (b_{(j,e)}(t))^\sigma \leq c^{\sigma' - \sigma} (b_{(j,e)}(t))^\sigma$$

Therefore, $(b_{(j,e)}(t))^\sigma \geq \mu(b_{(j,e)}(t))^{\sigma'}$, where $\mu = (\frac{1}{c})^{\sigma'-\sigma} > 1$. It follows that for all $t \in S_j$

$$\begin{aligned} (L_\sigma u_{\sigma'})_j(t) &= \sum_{e \in E_j} (b_{(j,e)}(t))^\sigma (u_{\sigma'})_{\alpha(j,e)}(\theta_{(j,e)}(t)) \\ &\geq \mu \sum_{e \in E_j} (b_{(j,e)}(t))^{\sigma'} (u_{\sigma'})_{\alpha(j,e)}(\theta_{(j,e)}(t)) \\ &= \mu (L'_\sigma u_{\sigma'})_j(t) = \mu r(L_{\sigma'}) (u_{\sigma'})_j(t), \end{aligned}$$

so $L_\sigma u_{\sigma'} \geq \mu r(L_{\sigma'}) u_{\sigma'}$, where the inequality has the natural coordinate-wise interpretation. Iterating this inequality k times, we obtain

$$L_\sigma^k u_{\sigma'} \geq (\mu r(L_{\sigma'}))^k u_{\sigma'}$$

If e denotes the function identically equal to one in each component, we have $u_{\sigma'} \leq \|u_{\sigma'}\| e$. Thus

$$L_\sigma^k u_{\sigma'} \leq L_\sigma^k (\|u_{\sigma'}\| e) \leq \|u_{\sigma'}\| L_\sigma^k(e).$$

Taking norms, we get

$$\|u_{\sigma'}\| \|L_\sigma^k(e)\| \geq \|L_\sigma^k u_{\sigma'}\| \geq (\mu r(L_{\sigma'}))^k \|u_{\sigma'}\|.$$

So, $\|L_\sigma^k\| \geq \|L_\sigma^k(e)\| \geq (\mu r(L_{\sigma'}))^k$ from which it follows that

$$r(L_\sigma) = \lim_{k \rightarrow \infty} \|L_\sigma^k\|^{1/k} \geq \mu r(L_{\sigma'}).$$

Since $\mu > 1$, we have proved that $r(L_\sigma) > r(L_{\sigma'})$.

Next we prove the continuity of $\sigma \mapsto r(L_\sigma)$. Let $\sigma > 0$ be fixed. Given $\nu < 1$, select $\delta > 0$ such that

$$\nu (b_{(j,e)}(t))^\sigma \leq (b_{(j,e)}(t))^{\sigma'} \leq \nu^{-1} (b_{(j,e)}(t))^\sigma \text{ for } t \in S_j, |\sigma - \sigma'| \leq \delta.$$

Then, using the argument as above, we have $\nu r(L_\sigma) \leq r(L_{\sigma'}) \leq \nu^{-1} r(L_\sigma)$ whenever $|\sigma - \sigma'| \leq \delta$. Since $\nu < 1$ was arbitrary, this proves that $\sigma \mapsto r(L_\sigma)$ is continuous.

Since $\|L_0^k\| \geq \|L_0^k e\| \geq 1$, we see that $r(L_0) \geq 1$. Also if $|\mathcal{E}|$ denotes the cardinality of \mathcal{E} , then from the definition of L_σ and using the fact $D\theta_{(j,e)}(t) \leq c$ for all $t \in S_j$ and $(j,e) \in \Gamma$, it is clear that $\|L_\sigma\| \leq |\mathcal{E}| c^\sigma \rightarrow 0$ as $\sigma \rightarrow \infty$, so $r(L_\sigma) \leq \|L_\sigma\| \rightarrow 0$ as $\sigma \rightarrow \infty$. It follows by the continuity and strict monotonicity of $\sigma \rightarrow r(L_\sigma)$ that there exists a unique $\sigma_0 \geq 0$ such that $r(L_{\sigma_0}) = 1$. \square

Definition 4.7. We define *strong connectedness* to be the property that for each pair j and k in V there exists for some $n \geq 1$, $(J, E) = [(j_1, e_1), \dots, (j_n, e_n)]$ such that $(j_i, e_i) \in \Gamma$, for $1 \leq i \leq n$, $j_1 = j$, $\alpha(j_i, e_i) = j_{i+1}$, $1 \leq i < n$ and $\alpha(j_n, e_n) = k$. Note that in this case we have a map $\theta_{(J,E)} = \theta_{(j_n, e_n)} \circ \dots \circ \theta_{(j_1, e_1)}$ which maps S_j into S_k . Note also (compare H4.1) that strong connectedness implies that $\Gamma_i \neq \emptyset$ for $1 \leq i \leq p$ and $E_j \neq \emptyset$ for $1 \leq j \leq p$.

From now on we shall always assume strong connectedness.

H4.4 The property of strong connectedness is satisfied.

Lemma 4.8. *Assume that the hypotheses H4.1, H4.2, H4.3 and H4.4 are satisfied and let $u_\sigma \in K \setminus \{0\}$ be an eigenvector of L_σ with eigenvalue $r(L_\sigma)$. Then each component $(u_\sigma)_j$ is a strictly positive function on S_j for $1 \leq j \leq p$. Furthermore, there are constants l and L with $0 < l \leq L < \infty$ such that for every j , $1 \leq j \leq p$*

$$(4.5) \quad l \leq (u_\sigma)_j(t) \leq L \quad \text{for all } t \in S_j$$

Proof. Suppose for some j , $1 \leq j \leq p$, $(u_\sigma)_j$ equals zero at some point in S_j . Then, since $(u_\sigma)_j \in K_j(M, \lambda)$, it follows, as shown in Remark 3.1, that $(u_\sigma)_j$ is identically equal to zero on S_j . Fix a k , $1 \leq k \leq p$. By strong connectedness, there exist $n \geq 1$ and $(J', E') = [(j_1, e_1), \dots, (j_n, e_n)] \in \bar{\Gamma}_j^{(n)}$ with $j_1 = j$, and $\alpha(j_n, e_n) = k$. Since $L_\sigma u_\sigma = r(L_\sigma)u_\sigma$, it follows that $L_\sigma^n u_\sigma = (r(L_\sigma))^n u_\sigma$. So, using the formula for L_σ^n given by lemma 3.4 with A replaced by L_σ , we get

$$(r(L_\sigma))^n (u_\sigma)_j(s) = \sum_{(J,E) \in \bar{\Gamma}_j^{(n)}} b_{(J,E)}(s) (u_\sigma)_{\alpha(j_n, e_n)}(\theta_{(J,E)}(s)),$$

where $b_{(j,e)}(s) = ((D\theta_{(j,e)})(s))^\sigma$. The left hand side in the above equation is zero because $(u_\sigma)_j(s) = 0$. Thus, since each term in the sum in the right hand is nonnegative, it follows that each term equals zero. In particular, $(u_\sigma)_k(\theta_{(J,E)}(s)) = 0$ since $b_{(J,E)}(s)$ is strictly positive by H4.2. This implies $(u_\sigma)_k$ is identically equal to zero on S_k . Since this is true for any k , $1 \leq k \leq p$, we arrive at a contradiction that u_σ is identically zero. Thus $(u_\sigma)_j$ is a strictly positive function on S_j for $1 \leq j \leq p$.

Since each S_j is bounded, there is a $D < \infty$ such that $\text{diam}(S_j) \leq D$ for $1 \leq j \leq p$. Then, since $(u_\sigma)_j \in K_j(M, \lambda)$, it follows that

$$0 < (u_\sigma)_j(s) \leq (u_\sigma)_j(t) \exp(MD^\lambda) \text{ for all } s, t \in S_j.$$

From this it is easy to see that there are constants $0 < l \leq L < \infty$ such that

$$l \leq (u_\sigma)_j(t) \leq L \text{ for all } t \in S_j, 1 \leq j \leq p.$$

□

Let $C_j \subset S_j$, $1 \leq j \leq p$ be the invariant list of nonempty compact sets such that

$$C_i = \bigcup_{(j,e) \in \Gamma_i} \theta_{(j,e)}(C_j) \text{ for } 1 \leq i \leq p.$$

Our goal is to determine the Hausdorff dimension of the sets C_i .

Let us recall the definition of Hausdorff measure and Hausdorff dimension. Suppose (X, d) is a metric space and $A \subset X$. We define, for $\epsilon > 0$ and $\sigma \geq 0$,

$$\mathcal{H}_\epsilon^\sigma(A) = \inf \left\{ \sum_{k=1}^{\infty} (\text{diam}(A_k))^\sigma : A \subset \bigcup_{k=1}^{\infty} A_k, \text{diam}(A_k) < \epsilon \right\}.$$

It follows that $\mathcal{H}_\epsilon^\sigma$ is an outer measure. For a given $\sigma \geq 0$, the function $\epsilon \mapsto \mathcal{H}_\epsilon^\sigma(A)$ is decreasing and we define

$$\mathcal{H}^\sigma(A) = \lim_{\epsilon \rightarrow 0^+} \mathcal{H}_\epsilon^\sigma(A) = \sup_{\epsilon > 0} \mathcal{H}_\epsilon^\sigma(A).$$

It follows that \mathcal{H}^σ is a Borel measure and is called *Hausdorff σ -dimensional measure*. It is not hard to prove that there is a unique number $\sigma_0 \geq 0$ such that $\mathcal{H}^\sigma(A) = \infty$ for $0 \leq \sigma < \sigma_0$ and $\mathcal{H}^\sigma(A) = 0$ for $\sigma > \sigma_0$. The number σ_0 is called the *Hausdorff dimension* of A . We refer the reader to [5] and [11] for the basic properties of Hausdorff measure.

First we shall prove that Hausdorff dimension of C_i is independent of i , $1 \leq i \leq p$, under the assumption of strong connectedness.

Lemma 4.9. *Assume that the hypotheses H4.1, H4.2, H4.3 and H4.4 are satisfied and let $C_j \subset S_j$, $1 \leq j \leq p$ be the unique invariant list of compact, nonempty sets such that*

$$C_i = \bigcup_{(j,e) \in \Gamma_i} \theta_{(j,e)}(C_j) \text{ for } 1 \leq i \leq p.$$

Then $\dim(C_j)$, the Hausdorff dimension of C_j , is independent of j for $1 \leq j \leq p$.

Proof. First we claim that $\dim(\theta_{(j,e)}(C_j)) = \dim(C_j)$ for any $(j,e) \in \Gamma$. Since $\theta_{(j,e)}$ is a Lipschitz map with Lipschitz constant c , $\mathcal{H}^\sigma(\theta_{(j,e)}(C_j)) \leq c^\sigma \mathcal{H}^\sigma(C_j)$ for any $\sigma \geq 0$. This implies $\dim(\theta_{(j,e)}(C_j)) \leq \dim(C_j)$. To prove the other inequality, we first claim that there exist $m_0 > 0$ and $\delta > 0$ such that $d(\theta_{(j,e)}(s), \theta_{(j,e)}(t)) \geq m_0 d(s,t)$ for all $s, t \in C_j$ with $d(s,t) \leq \delta$. We abuse notation here by letting d denote d_j and $d_{\alpha(j,e)}$. We argue by contradiction. If the claim is false, then for each positive integer k there exist $s_k, t_k \in C_j$ with $0 < d(s_k, t_k) \leq k^{-1}$ and $d(\theta_{(j,e)}(s_k), \theta_{(j,e)}(t_k)) < k^{-1} d(s_k, t_k)$. Since C_j is compact, by taking a subsequence we can assume that $s_k \rightarrow s$ and $t_k \rightarrow s$. But this implies that $D\theta_{(j,e)}(s) = 0$, which contradicts H4.2. Thus $m_0 > 0$ and $\delta > 0$ exist, and since C_j is compact, we can write $C_j = \bigcup_{l=1}^p C_{j,l}$, where $\text{diam}(C_{j,l}) \leq \delta$ and $p < \infty$. It follows easily from the definition of Hausdorff dimension that there exists l such that the Hausdorff dimension of $C_{j,l}$ equals the Hausdorff dimension of C_j . Also, by our construction, $\theta_{(j,e)}|_{C_{j,l}}$ is one-to-one and $(\theta_{(j,e)}|_{C_{j,l}})^{-1}$ is Lipschitz. This implies that

$$\dim(C_j) = \dim(C_{j,l}) \leq \dim(\theta_{(j,e)}(C_{j,l})) \leq \dim(\theta_{(j,e)}(C_j)),$$

and we have shown that $\dim(C_j) = \dim(\theta_{(j,e)}(C_j))$.

Now, since $\theta_{(j,e)}(C_j) \subset C_{\alpha(j,e)}$, $\dim(C_{\alpha(j,e)}) \geq \dim(\theta_{(j,e)}(C_j)) = \dim(C_j)$ for all $(j,e) \in \Gamma$. Let $1 \leq j \leq p$ and $1 \leq k \leq p$. By strong connectedness, there exists $[(j_1, e_1), \dots, (j_n, e_n)]$ such that $j_1 = j$, $\alpha(j_i, e_i) = j_{i+1}$, $1 \leq i < n$ and $\alpha(j_n, e_n) = k$. So,

$$\dim(C_k) = \dim(C_{\alpha(j_n, e_n)}) \geq \dim(C_{j_n}) \geq \dim(C_{j_{n-1}}) \geq \dots \geq \dim(C_{j_1}) = \dim(C_j).$$

Since j and k were arbitrary, it follows that $\dim(C_j) = \dim(C_k)$ for all $1 \leq j, k \leq p$. \square

We introduce a ‘weighted’ Hausdorff measure using the strictly positive eigenvector u_σ of L_σ with eigenvalue $r(L_\sigma)$. Let $1 \leq j \leq p$. Define for $A_j \subset S_j$ and $\epsilon > 0$,

$$(4.6) \quad \tilde{\mathcal{H}}_\epsilon^\sigma(A_j) = \inf \left\{ \sum_{k=1}^{\infty} (u_\sigma)_j(\xi_{jk}) (\text{diam}(A_{jk}))^\sigma : A_j \subset \bigcup_{k=1}^{\infty} A_{jk}, \right. \\ \left. \xi_{jk} \in A_{jk}, \text{diam}(A_{jk}) < \epsilon \right\}.$$

From Lemma 4.8, we know that there exist constants $0 < l \leq L < \infty$ such that for $1 \leq j \leq p$, $l \leq (u_\sigma)_j(t) \leq L$ for all $t \in S_j$. This implies that, for $A_j \subset S_j$, $\tilde{\mathcal{H}}_\epsilon^\sigma(A_j)$ and $\mathcal{H}_\epsilon^\sigma(A_j)$ are equivalent

$$(4.7) \quad l\mathcal{H}_\epsilon^\sigma(A_j) \leq \tilde{\mathcal{H}}_\epsilon^\sigma(A_j) \leq L\mathcal{H}_\epsilon^\sigma(A_j)$$

Theorem 4.10. *Assume that the hypotheses H4.1, H4.2, H4.3 and H4.4 are satisfied and let $C_j \subset S_j$, $1 \leq j \leq p$ be the unique invariant list of compact, nonempty sets such that*

$$C_i = \bigcup_{(j,e) \in \Gamma_i} \theta_{(j,e)}(C_j) \text{ for } 1 \leq i \leq p.$$

If $\dim(C_i)$ denotes the Hausdorff dimension of (C_i) and σ_0 denotes the unique nonnegative real number such that $r(L_{\sigma_0}) = 1$, then $\dim(C_i) \leq \sigma_0$ for $1 \leq i \leq p$.

Proof. Fix $\epsilon > 0$. Take $\delta > 0$ and $\sigma > 0$. We can choose a covering $\{A_{jk}\}_{k=1}^{\infty}$ of C_j and points $\xi_{jk} \in A_{jk}$ such that $\text{diam}(A_{jk}) < \epsilon$ and

$$(4.8) \quad \sum_{k=1}^{\infty} (u_{\sigma})_j(\xi_{jk}) (\text{diam}(A_{jk}))^{\sigma} \leq \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(C_j) + \delta.$$

Since $C_i = \bigcup_{(j,e) \in \Gamma_i} \theta_{(j,e)}(C_j)$, we have that $\{\theta_{(j,e)}(A_{jk}) : 1 \leq k < \infty, (j,e) \in \Gamma_i\}$ is a covering of C_i with

$$\text{diam}(\theta_{(j,e)}(A_{jk})) \leq c \text{diam}(A_{jk}) < c\epsilon.$$

Furthermore, using Lemma 4.3 and Remark 4.4, it is easy to see that there exists $\mu_{(j,e)}(\epsilon) > 1$ with $\mu_{(j,e)}(\epsilon) \rightarrow 1$ as $\epsilon \rightarrow 0+$ such that

$$(4.9) \quad \text{diam}(\theta_{(j,e)}(A_{jk})) \leq \mu_{(j,e)}(\epsilon) (D\theta_{(j,e)})(\xi_{jk}) \text{diam}(A_{jk}).$$

Let $\mu(\epsilon) = \max_{(j,e) \in \Gamma} \mu_{(j,e)}(\epsilon)$.

$$\begin{aligned} \tilde{\mathcal{H}}_{c\epsilon}^{\sigma}(C_i) &\leq \sum_{k=1}^{\infty} \sum_{(j,e) \in \Gamma_i} (u_{\sigma})_i(\theta_{(j,e)}(\xi_{jk})) (\text{diam}(\theta_{(j,e)}(A_{jk})))^{\sigma} \\ &\leq (\mu(\epsilon))^{\sigma} \sum_{k=1}^{\infty} \sum_{(j,e) \in \Gamma_i} (u_{\sigma})_i(\theta_{(j,e)}(\xi_{jk})) ((D\theta_{(j,e)})(\xi_{jk}))^{\sigma} (\text{diam}(A_{jk}))^{\sigma}. \end{aligned}$$

Summing over i , $1 \leq i \leq p$, we have

$$\sum_{i=1}^p \tilde{\mathcal{H}}_{c\epsilon}^{\sigma}(C_i) \leq (\mu(\epsilon))^{\sigma} \sum_{i=1}^p \sum_{k=1}^{\infty} \sum_{(j,e) \in \Gamma_i} (u_{\sigma})_i(\theta_{(j,e)}(\xi_{jk})) ((D\theta_{(j,e)})(\xi_{jk}))^{\sigma} (\text{diam}(A_{jk}))^{\sigma}.$$

Rearranging the sum, we get

$$\begin{aligned} \sum_{i=1}^p \tilde{\mathcal{H}}_{c\epsilon}^{\sigma}(C_i) &\leq (\mu(\epsilon))^{\sigma} \sum_{j=1}^p \sum_{k=1}^{\infty} (\text{diam}(A_{jk}))^{\sigma} \sum_{i:(j,e) \in \Gamma_i} (u_{\sigma})_i(\theta_{(j,e)}(\xi_{jk})) ((D\theta_{(j,e)})(\xi_{jk}))^{\sigma} \\ &= (\mu(\epsilon))^{\sigma} \sum_{j=1}^p \sum_{k=1}^{\infty} (L_{\sigma} u_{\sigma})_j(\xi_{jk}) (\text{diam}(A_{jk}))^{\sigma} \\ &= (\mu(\epsilon))^{\sigma} r(L_{\sigma}) \sum_{j=1}^p \sum_{k=1}^{\infty} (u_{\sigma})_j(\xi_{jk}) (\text{diam}(A_{jk}))^{\sigma}. \end{aligned}$$

Thus, using (4.8), we get

$$\sum_{i=1}^p \tilde{\mathcal{H}}_{c\epsilon}^{\sigma}(C_i) \leq (\mu(\epsilon))^{\sigma} r(L_{\sigma}) \sum_{j=1}^p \left(\tilde{\mathcal{H}}_{\epsilon}^{\sigma}(C_j) + \delta \right).$$

Since $c < 1$, $\tilde{\mathcal{H}}_\epsilon^\sigma(C_i) \leq \tilde{\mathcal{H}}_{c\epsilon}^\sigma(C_i)$. Also $\delta > 0$ was arbitrary. Therefore,

$$(4.10) \quad \sum_{i=1}^p \tilde{\mathcal{H}}_\epsilon^\sigma(C_i) \leq (\mu(\epsilon))^\sigma r(L_\sigma) \sum_{i=1}^p \tilde{\mathcal{H}}_\epsilon^\sigma(C_i).$$

Using Lemma 4.6, $r(L_\sigma) < 1$ for all $\sigma > \sigma_0$. Since $\mu(\epsilon) \rightarrow 1$ as $\epsilon \rightarrow 0$, given $\sigma > \sigma_0$, we can choose $\epsilon > 0$ small so that $(\mu(\epsilon))^\sigma r(L_\sigma) < 1$. By the definition, $\tilde{\mathcal{H}}_\epsilon^\sigma(C_i) < \infty$ because we can take a finite ϵ -cover of the compact set C_i . Thus, if $\sigma > \sigma_0$, (4.10) can hold only if

$$\lim_{\epsilon \rightarrow 0^+} \sum_{i=1}^p \tilde{\mathcal{H}}_\epsilon^\sigma(C_i) = 0.$$

This implies for each i , $1 \leq i \leq p$ and $\sigma > \sigma_0$, $\lim_{\epsilon \rightarrow 0^+} \tilde{\mathcal{H}}_\epsilon^\sigma(C_i) = 0$, and hence using (4.7), $\lim_{\epsilon \rightarrow 0^+} \mathcal{H}_\epsilon^\sigma(C_i) = 0$, i.e., $\mathcal{H}^\sigma(C_i) = 0$ for all $\sigma > \sigma_0$. Thus, by the definition of Hausdorff dimension, $\dim(C_i) \leq \sigma_0$. \square

We define for $0 < \eta < \epsilon$, and $A_j \subset S_j$

$$(4.11) \quad \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j) := \inf \left\{ \sum_{k=1}^{\infty} (u_\sigma)_j(\xi_{jk}) (\text{diam}(A_{jk}))^\sigma : A_j \subset \bigcup_{k=1}^{\infty} A_{jk}, \right. \\ \left. \xi_{jk} \in A_{jk}, \eta < \text{diam}(A_{jk}) < \epsilon \right\}.$$

The quantity $\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j)$ will be technically useful later, primarily because it is strictly positive whenever it is defined. However, caution is necessary in using this tool. It is easy to see that $\tilde{\mathcal{H}}^\sigma(A_j)$ and $\tilde{\mathcal{H}}_\epsilon^\sigma(A_j)$ depend only on the metric space (A_j, d_j) . In contrast, $\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j)$ depends also on S_j . If $A_j \subset T_j \subset S_j$ one could give an analogous definition in which one only allows sets $A_{jk} \subset T_j$:

$$(4.12) \quad \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j; T_j) := \inf \left\{ \sum_{k=1}^{\infty} (u_\sigma)_j(\xi_{jk}) (\text{diam}(A_{jk}))^\sigma : A_j \subset \bigcup_{k=1}^{\infty} A_{jk} \subset T_j, \right. \\ \left. \xi_{jk} \in A_{jk}, \eta < \text{diam}(A_{jk}) < \epsilon \right\}.$$

In this cumbersome notation, $\tilde{\mathcal{H}}_\epsilon^\sigma(A_j) = \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j; S_j)$ for $A_j \subset S_j$. If A_j is compact and $A_j \subset T_j \subset S_j$, it may happen that $\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j; T_j)$ and $\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j; S_j)$ are both defined but are unequal. For the unique list of nonempty compact sets $C_j \subset S_j$, $1 \leq j \leq p$, ensured by Theorem 2.3 and for $A_j \subset C_j$, we shall, in our later work, sometimes consider $\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j; C_j)$ rather than $\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j; S_j)$.

In general, if $A_j \subset T_j \subset S_j$ and A_j is compact and $0 < \eta < \epsilon$, in order that $\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j; T_j)$ be defined and finite, it is necessary and sufficient that there exist sets $A_{jk} \subset T_j$, $1 \leq k \leq m < \infty$, with $\eta < \text{diam}(A_{jk}) < \epsilon$ for $1 \leq k \leq m$ and $A_j \subset \bigcup_{k=1}^m A_{jk}$. The existence or non-existence of such sets may be a delicate question. If A_j contains an isolated point of T_j , such sets A_{jk} do not exist for all small ϵ and all η with $0 < \eta < \epsilon$. If T_j is a complete perfect metric space, $A_j \subset T_j$ is compact and $\epsilon > 0$, $\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(A_j; T_j)$ will be defined for all sufficiently small $\eta > 0$. To see this, use compactness of A_j to find finitely many open balls $B_k \subset T_j$, $1 \leq k \leq n$, with radius $r < \epsilon/2$ and centers in A_j , such that $A_j \subset \bigcup_{k=1}^n B_k$. Since

T_j is perfect, each B_k contains an accumulation point, so $\text{diam}(B_k) > \eta_k > 0$ and $\text{diam}(B_k) > \eta := \min\{\eta_k : 1 \leq k \leq n\}$ and $\text{diam}(B_k) \leq 2r < \epsilon$.

Lemma 4.11. *Assume that H4.1, H4.2, H4.3 and H4.4 are satisfied. Let $1 \leq j \leq p$ and let A_j be a compact subset of S_j . If $\sigma \geq 0$ and $\epsilon > 0$, then*

$$\lim_{\eta \rightarrow 0^+} \tilde{\mathcal{H}}_{\epsilon, \eta}^{\sigma}(A_j) = \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(A_j).$$

Proof. For $0 < \eta < \epsilon$, we have $\tilde{\mathcal{H}}_{\epsilon, \eta}^{\sigma}(A_j) \geq \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(A_j)$ because the infimum is taken over a smaller set. So,

$$\lim_{\eta \rightarrow 0^+} \tilde{\mathcal{H}}_{\epsilon, \eta}^{\sigma}(A_j) \geq \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(A_j).$$

To prove the reverse inequality, take $\delta > 0$ and choose a covering $\{A_{jk} : 1 \leq k < \infty\}$ of A_j by sets A_{jk} with $\text{diam}(A_{jk}) < \epsilon$, $1 \leq k < \infty$ such that

$$\inf \left\{ \sum_{k=1}^{\infty} (u_{\sigma})_j(\xi_{jk})(\text{diam}(A_{jk}))^{\sigma} : \xi_{jk} \in A_{jk} \right\} \leq \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(A_j) + \delta.$$

Without loss of generality, we can assume that the sets $A_{jk}, k \geq 1$ are open. Since A_j is compact, there exists a finite open subcover of A_j , so there exists an integer $l < \infty$ such that

$$A_j \subset \bigcup_{k=1}^l A_{jk}.$$

Let $0 < \eta_0 < \epsilon$ be such that $\eta_0 < \min_{1 \leq k \leq l} \text{diam}(A_{jk})$. Then, for $0 < \eta < \eta_0$, we have

$$\tilde{\mathcal{H}}_{\epsilon, \eta}^{\sigma}(A_j) \leq \inf \left\{ \sum_{k=1}^l (u_{\sigma})_j(\xi_{jk})(\text{diam}(A_{jk}))^{\sigma} : \xi_{jk} \in A_{jk} \right\}.$$

So, for every $\delta > 0$, there exists $\eta_0, 0 < \eta_0 < \epsilon$, such that

$$\tilde{\mathcal{H}}_{\epsilon, \eta}^{\sigma}(A_j) \leq \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(A_j) + \delta \quad \text{for } 0 < \eta < \eta_0.$$

This shows

$$\lim_{\eta \rightarrow 0^+} \tilde{\mathcal{H}}_{\epsilon, \eta}^{\sigma}(A_j) \leq \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(A_j)$$

and completes the proof of the lemma. \square

Lemma 4.12. *Assume that H4.1, H4.2, H4.3 and H4.4 are satisfied. Let $1 \leq j \leq p$ and let A_j be a compact subset of S_j . Let $\sigma > 0$ be such that $\mathcal{H}^{\sigma}(A_j) = 0$. Then for every ϵ_1 and ϵ_2 with $0 < \epsilon_1 < \epsilon_2$, there exists an $\eta_0 > 0$ such that for any $B_j \subset A_j$*

$$\tilde{\mathcal{H}}_{\epsilon_1, \eta}^{\sigma}(B_j) = \tilde{\mathcal{H}}_{\epsilon_2, \eta}^{\sigma}(B_j) \quad \text{for } 0 < \eta < \eta_0.$$

Proof. Since $\mathcal{H}^{\sigma}(A_j) = 0$, it follows that $\mathcal{H}_{\epsilon}^{\sigma}(A_j) = 0$ for every $\epsilon > 0$. By using (4.7), it also follows that $\tilde{\mathcal{H}}_{\epsilon}^{\sigma}(A_j) = 0$ for every $\epsilon > 0$. So, by Lemma 4.11,

$$\lim_{\eta \rightarrow 0^+} \tilde{\mathcal{H}}_{\epsilon_2, \eta}^{\sigma}(A_j) = 0.$$

This implies that there exists $\eta_0 > 0$ such that for $0 < \eta < \eta_0$

$$\tilde{\mathcal{H}}_{\epsilon_2, \eta}^{\sigma}(A_j) < l\epsilon_1^{\sigma},$$

where, as before, $l > 0$ is such that $(u_{\sigma})_j(t) \geq l$ for all $t \in S_j$.

If $B_j \subset A_j$, then $\tilde{\mathcal{H}}_{\epsilon_2, \eta}^\sigma(B_j) \leq \tilde{\mathcal{H}}_{\epsilon_2, \eta}^\sigma(A_j) < l\epsilon_1^\sigma$. Therefore, given $\delta > 0$, there exists a covering $\{B_{j_k} : k \geq 1\}$ of B_j such that $\eta < \text{diam}(B_{j_k}) < \epsilon_2$ for $k \geq 1$ and

$$\inf \left\{ \sum_{k=1}^{\infty} (u_\sigma)_j(\xi_{j_k}) (\text{diam}(B_{j_k}))^\sigma : \xi_{j_k} \in B_{j_k} \right\} \leq \tilde{\mathcal{H}}_{\epsilon_2, \eta}^\sigma(B_j) + \delta < l\epsilon_1^\sigma.$$

Next we claim that actually $\text{diam}(B_{j_k}) < \epsilon_1$ for all $k \geq 1$. Suppose not, then there exists an index k_1 such that $\text{diam}(B_{j_{k_1}}) \geq \epsilon_1$. By considering the term corresponding to index k_1 in the sum and using $(u_\sigma)_j(\xi_{j_k}) \geq l$, we get

$$\inf \left\{ \sum_{k=1}^{\infty} (u_\sigma)_j(\xi_{j_k}) (\text{diam}(B_{j_k}))^\sigma : \xi_{j_k} \in B_{j_k} \right\} \geq l\epsilon_1^\sigma$$

which gives a contradiction. Thus $\text{diam}(B_{j_k}) < \epsilon_1$ for all $k \geq 1$ and we conclude that

$$\tilde{\mathcal{H}}_{\epsilon_1, \eta}^\sigma(B_j) \leq \tilde{\mathcal{H}}_{\epsilon_2, \eta}^\sigma(B_j) + \delta \quad \text{for } 0 < \eta < \eta_0.$$

Since $\delta > 0$ was arbitrary, we conclude

$$\tilde{\mathcal{H}}_{\epsilon_1, \eta}^\sigma(B_j) \leq \tilde{\mathcal{H}}_{\epsilon_2, \eta}^\sigma(B_j) \quad \text{for } 0 < \eta < \eta_0.$$

Since $\tilde{\mathcal{H}}_{\epsilon, \eta}^\sigma(B_j)$ is a decreasing function of ϵ , the reverse inequality is obvious. Thus, we obtain

$$\tilde{\mathcal{H}}_{\epsilon_1, \eta}^\sigma(B_j) = \tilde{\mathcal{H}}_{\epsilon_2, \eta}^\sigma(B_j) \quad \text{for } 0 < \eta < \eta_0. \quad \square$$

If $C_j \subset S_j$, $1 \leq j \leq p$, is the unique invariant list guaranteed by Theorem 2.3, it is convenient in the arguments below to work in the compact sets C_j rather than S_j . For this to be permissible, we must first show that each set C_j is a perfect metric space. Our first lemma in this direction has essentially been established in the proof of Lemma 4.9 and is left to the reader.

Lemma 4.13. *Assume hypotheses H4.1, H4.2, H4.3 and H4.4 and let $C_j \subset S_j$, $1 \leq j \leq p$, be the invariant sets guaranteed by Theorem 2.3. Then there exist $m_0 > 0$ and $\delta_0 > 0$ such that for all $(j, e) \in \Gamma$ and for all $s, t \in S_j$ with $s \in C_j$ and $d_j(s, t) < \delta_0$,*

$$d_{\alpha(j, e)}(\theta_{(j, e)}(s), \theta_{(j, e)}(t)) \geq m_0 d_j(s, t).$$

Lemma 4.14. *Assume hypotheses H4.1, H4.2, H4.3 and H4.4 and let $C_j \subset S_j$, $1 \leq j \leq p$, be the invariant sets guaranteed by Theorem 2.3. Then either (a) each C_j , $1 \leq j \leq p$, is a perfect metric space or (b) each C_j , $1 \leq j \leq p$, is a finite set. If $\theta_{(j, e)}|_{C_j}$ is one-to-one for all $(j, e) \in \Gamma$ and each C_j is a finite set, then each C_j contains exactly one element for $1 \leq j \leq p$.*

Proof. Assume that C_i is not a finite set for some i , $1 \leq i \leq p$. It follows that there exists $\tau_i \in C_i$ such that τ_i is an accumulation point of C_i . If $1 \leq k \leq p$, H4.4 implies that there exist $(j_l, e_l) \in \Gamma$, $1 \leq l \leq n$, with $\alpha(j_l, e_l) = j_{l+1}$ for $1 \leq l < n$, $j_1 = i$ and $\alpha(j_n, e_n) = k$. Writing $(J, E) = [(j_1, e_1), (j_2, e_2), \dots, (j_n, e_n)]$, $\theta_{(J, E)}(\tau_i) \in C_k$. Also, because each $\theta_{(j, e)}$ is a contraction, Lemma 4.13 implies that $\theta_{(J, E)}$ is one-to-one on $B_{r_0}(\tau_i) := \{t \in C_i : d_i(t, \tau_i) < r_0\}$, where $r_0 = \delta_0/2$ and δ_0 is as in Lemma 4.13. Using this fact, we see that $\theta_{(J, E)}(\tau_i) := \tau_k$ is an accumulation point of C_k .

If $V = \{1, 2, \dots, p\}$, Θ is as in the proof of Theorem 2.3 and A_j is any closed, bounded, nonempty subset of S_j for $1 \leq j \leq p$, then, because Θ is a contraction map

in the Hausdorff metric, $\Theta^n((A_1, A_2, \dots, A_p))$ converges in the Hausdorff metric to (C_1, C_2, \dots, C_p) as $n \rightarrow \infty$. We apply this result to $A_j = \{\tau_j\}$, where τ_j is an accumulation point of C_j for $1 \leq j \leq p$. If k , $1 \leq k \leq p$, and $\sigma \in C_k$ are fixed, it follows that for each $n \geq 1$, there exists $(J^n, E^n) = [(j_1^n, e_1^n), (j_2^n, e_2^n) \dots, (j_n^n, e_n^n)]$ with $(j_i^n, e_i^n) \in \Gamma$ for $1 \leq i \leq n$, $\alpha(j_i^n, e_i^n) = (j_{i+1}^n, e_{i+1}^n)$ for $1 \leq i < n$, $\alpha(j_n^n, e_n^n) = k$ and $\theta_{(J^n, E^n)}(\tau_{j_1^n}) \rightarrow \sigma$ as $n \rightarrow \infty$. For notational convenience, write $i_n := j_1^n$. If r_0 and δ_0 are as above and $V_n = \{t \in C_{i_n} : d(t, \tau_{i_n}) < r_0\}$, $\theta_{(J^n, E^n)}|_{V_n}$ is one-to-one and $\text{diam}(\theta_{(J^n, E^n)}(V_n)) \leq \delta_0 c^n$, where $\text{Lip}(\theta_{(j,e)}) \leq c < 1$ for all $(j, e) \in \Gamma$. Because we also know that τ_{i_n} is an accumulation point of C_{i_n} , it follows that there exist points $s_n, t_n \in \theta_{(J^n, E^n)}(V_n)$, $s_n \neq t_n$ and necessarily $\lim_{n \rightarrow \infty} s_n = \lim_{n \rightarrow \infty} t_n = \sigma$. This, in turn, implies that σ is an accumulation point of C_k and that C_k is a perfect metric space for $1 \leq k \leq p$.

If $\theta_{(j,e)}|_{C_j}$ is one-to-one for each $(j, e) \in \Gamma$ and if each C_j is a finite set, we claim that each C_j is a one point set. If not, there exists j , $1 \leq j \leq p$, such that C_j contains at least two points. Using H4.4 and the fact that each $\theta_{(i,e)}|_{C_i}$ is one-to-one, it follows that C_k contains at least two points for $1 \leq k \leq p$. Define $\rho = \inf\{d_j(s, t) : s, t \in C_j, s \neq t, 1 \leq j \leq p\}$, so $\rho > 0$, and select j and $s, t \in C_j$ with $d_j(s, t) = \rho$. Because $\theta_{(j,e)}$ is a contraction and is one-to-one on C_j , $\theta_{(j,e)}(s) \neq \theta_{(j,e)}(t)$ and $d_{\alpha(j,e)}(\theta_{(j,e)}(s), \theta_{(j,e)}(t)) < \rho$, a contradiction. \square

Assume H4.1, H4.2, H4.3 and H4.4. If $f = (f_1, f_2, \dots, f_p) \in \prod_{j=1}^p C(C_j)$, where (C_1, C_2, \dots, C_p) is as in Theorem 2.3, we can define $\Lambda_\sigma : \prod_{j=1}^p C(C_j) \rightarrow \prod_{j=1}^p C(C_j)$ by eq. (4.4). Furthermore, if each C_j is a perfect metric space (compare Lemma 4.14) and $\psi_{(j,e)} : C_j \rightarrow C_{\alpha(j,e)}$ is defined by $\psi_{(j,e)}(s) = \theta_{(j,e)}(s)$ for $s \in C_j$, then $D\psi_{(j,e)}(s)$ is defined for all $s \in C_j$ and $D\psi_{(j,e)}(s) = D\theta_{(j,e)}(s)$ for $s \in C_j$. In any event, by using Lemma 4.8, it follows easily that the spectral radius of L_σ equals the spectral radius of Λ_σ and is the eigenvalue of the strictly positive eigenvector u_σ of Lemma 4.8. The corresponding positive eigenvector v_σ for Λ_σ is given by $(v_\sigma)_j(s) = (u_\sigma)_j(s)$ for $s \in C_j$. If each C_j is a finite set, $\prod_{j=1}^p C(C_j)$ is finite dimensional and determining $r(\Lambda_\sigma) = r(L_\sigma)$ is a finite dimensional problem.

Lemma 4.15. *Assume that hypotheses H 5.1, H 5.2, H 5.3 and H 5.4 are satisfied and let $C_j \subset S_j$, $1 \leq j \leq p$ be the unique invariant list such that*

$$C_i = \bigcup_{(j,e) \in \Gamma_i} \theta_{(j,e)}(C_j) \text{ for } 1 \leq i \leq p.$$

Also assume that the map $\theta_{(j,e)}|_{C_j}$ is one-to-one for all $(j, e) \in \Gamma$ and that, for some j , C_j has more than one element. Suppose that for all $(j, e) \in \Gamma$ and $(j', e') \in \Gamma$ such that $(j, e) \neq (j', e')$ and $\alpha(j, e) = \alpha(j', e')$ we have that $\theta_{(j,e)}(C_j) \cap \theta_{(j',e')}(C_{j'}) = \emptyset$. Finally, suppose, for $\nu > 0$, that $\{\Lambda_\beta^i : \beta \in \mathcal{A}_i(\nu)\}$ is a partition of C_i consisting of compact subsets Λ_β^i of C_i with $\text{diam}(\Lambda_\beta^i) < \nu$. Then there exist $\epsilon_0 > 0$ and $\nu_0 > 0$ such that for $0 < \epsilon < \epsilon_0$, $0 < \nu < \nu_0$, $0 < \eta < m\epsilon$ and $1 \leq i \leq p$,

$$(4.13) \quad \mu(\nu)r(L_\sigma)\tilde{\mathcal{H}}_{\mu_1(\nu)\epsilon, \eta}^\sigma(\Lambda_\beta^i; C_i) \leq \sum_{e \in E_i} \tilde{\mathcal{H}}_{m\epsilon, \eta}^\sigma(\theta_{(i,e)}(\Lambda_\beta^i); C_{\alpha(i,e)}),$$

where $\mu(\nu) \rightarrow 1$ and $\mu_1(\nu) \rightarrow 1$ as $\nu \rightarrow 0+$ and $(D\theta_{(i,e)})(t) > m$ for all $t \in C_i$ and all $(i, e) \in \Gamma$.

Remark. For compact sets $B \subset C_k$, we shall always use $\tilde{\mathcal{H}}_{\epsilon_1, \eta_1}^\sigma(B; C_k)$ in the following proof, so for notational simplicity we shall write $\tilde{\mathcal{H}}_{\epsilon_1, \eta_1}^\sigma(B)$ instead of $\tilde{\mathcal{H}}_{\epsilon_1, \eta_1}^\sigma(B; C_k)$.

We interpret eq. (4.13) as meaning that η with $0 < \eta < m\epsilon$ is such that $\tilde{\mathcal{H}}_{m\epsilon, \eta}^\sigma(\theta_{(i,e)}(\Lambda_\beta^i))$ is defined and finite for all $e \in E_i$ and that this implies that for $\epsilon^* := \mu_1(\nu)\epsilon$, $\tilde{\mathcal{H}}_{\epsilon^*, \eta}^\sigma(\Lambda_\beta^i)$ is defined and finite and eq. (4.13) is satisfied.

Proof. Lemma 4.14 implies that each C_j , $1 \leq j \leq p$, is a compact, perfect metric space. Since we assume that $\theta_{(j,e)}(C_j) \cap \theta_{(j',e')}(C_{j'}) = \emptyset$ whenever $(j,e) \neq (j',e')$ and $\alpha(j,e) = \alpha(j',e')$, we can select ϵ_0 so that $d_k(s,t) > \epsilon_0$ whenever $(j,e) \neq (j',e') \in \Gamma$, $\alpha(j,e) = \alpha(j',e') = k$, $s \in \theta_{(j,e)}(C_j)$ and $t \in \theta_{(j',e')}(C_{j'})$. Since, for $1 \leq k \leq p$, $C_k = \bigcup_{(j,e) \in \Gamma_k} \theta_{(j,e)}(C_j)$, it follows that, as a subset of the metric space (C_k, d_k) , $\theta_{(j,e)}(C_j)$ is both compact and open for all $(j,e) \in \Gamma$ with $\alpha(j,e) = k$. Furthermore, denoting by $N_\delta(B)$ the closed δ neighborhood in C_k of a compact set $B \subset C_k$, $N_\delta(\theta_{(j,e)}(C_j)) = \theta_{(j,e)}(C_j)$ for all $(j,e) \in \Gamma$ with $\alpha(j,e) = k$ and all δ with $0 < \delta \leq \epsilon_0$.

Fix $\nu > 0$, $1 \leq i \leq p$, Λ_β^i with $\beta \in \mathcal{A}_i(\nu)$. Suppose that ϵ_0 is as above. By decreasing ϵ_0 further we can also assume that $\epsilon_0 \leq (1-c)\nu/2m$. Suppose that $0 < \epsilon \leq \epsilon_0$, $0 < \eta < m\epsilon$ and that $\tilde{\mathcal{H}}_{m\epsilon, \eta}^\sigma(\theta_{(i,e)}(\Lambda_\beta^i))$ is defined and finite for all $e \in E_i$. For any $\delta > 0$, there exists an open (in the relative topology of $C_{\alpha(i,e)}$), necessarily finite covering of $\theta_{(i,e)}(\Lambda_\beta^i)$, $\{A_j : 1 \leq j \leq k\}$, in $C_{\alpha(i,e)}$, such that $\eta < \text{diam}(A_j) < m\epsilon$ for $1 \leq j \leq k$ and

$$(4.14) \quad \inf \left\{ \sum_{j=1}^k (u_\sigma)_{\alpha(i,e)}(\xi_j) (\text{diam}(A_j))^\sigma : \xi_j \in A_j \right\} \leq \tilde{\mathcal{H}}_{m\epsilon, \eta}^\sigma(\theta_{(i,e)}(\Lambda_\beta^i)) + \delta.$$

We denote by $N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i))$ the closed $m\epsilon$ neighborhood of $\theta_{(i,e)}(\Lambda_\beta^i)$ in $C_{\alpha(i,e)}$; and because $m\epsilon \leq \epsilon_0$, we observe that $N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i)) \subset \theta_{(i,e)}(C_i)$. We have that $\text{diam}(A_j) < m\epsilon$, and we can assume that A_j intersects $\theta_{(i,e)}(\Lambda_\beta^i)$ for $1 \leq j \leq k$, so

$$(4.15) \quad \theta_{(i,e)}(\Lambda_\beta^i) \subset \bigcup_{j=1}^k A_j \subset N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i)) \subset \theta_{(i,e)}(C_i).$$

Since $\text{diam}(\theta_{(i,e)}(\Lambda_\beta^i)) \leq c \text{diam}(\Lambda_\beta^i) < c\nu$,

$$(4.16) \quad \text{diam}(N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i))) \leq c\nu + 2m\epsilon < \nu$$

since we assume that $\epsilon < \epsilon_0 \leq (1-c)\nu/2m$. Since $\theta_{(i,e)}|_{C_i}$ is one-to-one, we derive from eq. (4.15) that we have $\theta_{(i,e)}^{-1} : \theta_{(i,e)}(C_i) \rightarrow C_i$ and

$$(4.17) \quad \Lambda_\beta^i \subset \bigcup_{j=1}^k \theta_{(i,e)}^{-1}(A_j) \subset C_i.$$

Using Lemma 4.3 and Remark 4.4, there exists $\mu_1(\nu)$ with $\mu_1(\nu) \rightarrow 1+$ as $\nu \rightarrow 0+$ and

$$(4.18) \quad \mu_1(\nu)^{-1} (D\theta_{(i,e)}^{-1})(y) \leq \frac{d(\theta_{(i,e)}^{-1}(x), \theta_{(i,e)}^{-1}(y))}{d(x,y)} \leq \mu_1(\nu) (D\theta_{(i,e)}^{-1})(y)$$

for all $x, y \in \theta_{(i,e)}(C_i)$ with $0 < d(x, y) < \nu$ and, in particular, for all $x, y \in N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i))$. Note that $\mu_1(\nu)$ can be chosen independent of $(i, e) \in \Gamma$. We write d instead of d_i or $d_{\alpha(i,e)}$ here and below. In particular, for any $x, y \in A_j$, $1 \leq j \leq k$,

$$d(\theta_{(i,e)}^{-1}(x), \theta_{(i,e)}^{-1}(y)) \leq \mu_1(\nu)(D\theta_{(i,e)}^{-1})(y)d(x, y) \leq \mu_1(\nu)\left(\frac{1}{(D\theta_{(i,e)})(\theta_{(i,e)}^{-1}(y))}\right)\text{diam}(A_j).$$

Using the compactness of $\theta_{(i,e)}^{-1}(N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i)))$, we see that there exists $\tau_1 \in \theta_{(i,e)}^{-1}(N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i)))$ with

$$(4.19) \quad \text{diam}(\theta_{(i,e)}^{-1}(A_j)) \leq \mu_1(\nu)\frac{1}{(D\theta_{(i,e)})(\tau_1)}\text{diam}(A_j).$$

Since $\text{diam}(A_j) < m\epsilon$, eq. (4.19) implies

$$\text{diam}(\theta_{(i,e)}^{-1}(A_j)) < \mu_1(\nu)\left(\frac{1}{m}\right)m\epsilon = \mu_1(\nu)\epsilon.$$

By choosing $x_0, y_0 \in N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i))$ with $d(x_0, y_0) = \text{diam}(A_j)$, we also obtain

$$\text{diam}(\theta_{(i,e)}^{-1}(A_j)) \geq d(\theta_{(i,e)}^{-1}(x_0), \theta_{(i,e)}^{-1}(y_0)) \geq \mu_1(\nu)^{-1}\left(\frac{1}{(D\theta_{(i,e)})(\theta_{(i,e)}^{-1}(y_0))}\right)\text{diam}(A_j)$$

Since $\text{diam}(A_j) > \eta$ and $(D\theta_{(i,e)})(\theta_{(i,e)}^{-1}(y_0)) \leq c < 1$, we find that $\text{diam}(\theta_{(i,e)}^{-1}(A_j)) \geq \mu_1(\nu)^{-1}c^{-1}\eta$. It follows that, assuming we originally chose ν_0 such that $\mu_1(\nu)c < 1$ for $0 < \nu < \nu_0$, we have $\text{diam}(\theta_{(i,e)}^{-1}(A_j)) > \eta$. Since $\Lambda_\beta^i \subset \bigcup_{j=1}^k \theta_{(i,e)}^{-1}(A_j)$, we conclude that

$$\tilde{\mathcal{H}}_{\mu_1(\nu)\epsilon, \eta}^\sigma(\Lambda_\beta^i) \leq \inf \left\{ \sum_{j=1}^k (u_\sigma)_i(\zeta_j)(\text{diam}(\theta_{(i,e)}^{-1}(A_j)))^\sigma : \zeta_j \in \theta_{(i,e)}^{-1}(A_j) \right\}.$$

Using eq. (4.19) and writing $\epsilon^* = \mu_1(\nu)\epsilon$,

$$\tilde{\mathcal{H}}_{\epsilon^*, \eta}^\sigma(\Lambda_\beta^i) \leq \frac{\mu_1(\nu)^\sigma}{((D\theta_{(i,e)})(\tau_1))^\sigma} \inf \left\{ \sum_{j=1}^k \frac{(u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\zeta_j))(\text{diam}(A_j))^\sigma (u_\sigma)_i(\zeta_j)}{(u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\zeta_j))} : \zeta_j \in \theta_{(i,e)}^{-1}(A_j) \right\}.$$

Choose $\tau_2 \in \theta_{(i,e)}^{-1}(N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i)))$ such that

$$\frac{(u_\sigma)_i(\tau_2)}{(u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\tau_2))} \geq \frac{(u_\sigma)_i(\zeta_j)}{(u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\zeta_j))}$$

for all $\zeta_j \in \theta_{(i,e)}^{-1}(A_j)$, $1 \leq j \leq k$. Using this together with eq. (4.14), we get

$$\tilde{\mathcal{H}}_{\epsilon^*, \eta}^\sigma(\Lambda_\beta^i) \leq \frac{\mu_1(\nu)^\sigma}{((D\theta_{(i,e)})(\tau_1))^\sigma} \frac{(u_\sigma)_i(\tau_2)}{(u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\tau_2))} (\tilde{\mathcal{H}}_{m\epsilon, \eta}^\sigma(\theta_{(i,e)}(\Lambda_\beta^i)) + \delta).$$

Since $\delta > 0$ was arbitrary,

$$(4.20) \quad \tilde{\mathcal{H}}_{\epsilon^*, \eta}^\sigma(\Lambda_\beta^i) \leq \frac{\mu_1(\nu)^\sigma}{((D\theta_{(i,e)})(\tau_1))^\sigma} \frac{(u_\sigma)_i(\tau_2)}{(u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\tau_2))} \tilde{\mathcal{H}}_{m\epsilon, \eta}^\sigma(\theta_{(i,e)}(\Lambda_\beta^i)).$$

The final step consists of replacing τ_1, τ_2 by a $\xi \in \Lambda_\beta^i$. From eq. (4.18), it follows that

$$\text{diam}(\theta_{(i,e)}^{-1}(N_{m\epsilon}(\theta_{(i,e)}(\Lambda_\beta^i)))) \leq \mu_1(\nu) \frac{1}{m} \nu \leq \kappa \nu,$$

where κ is independent of ν . In particular, we have that $d(\tau_1, \tau_2) \leq \kappa \nu$, so by continuity, there exists a function $\mu_2(\nu)$ such that $\mu_2(\nu) \rightarrow 1$ as $\nu \rightarrow 0$ and

$$\mu_2(\nu) ((D\theta_{(i,e)})(\xi))^\sigma \frac{(u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\xi))}{(u_\sigma)_i(\xi)} \leq ((D\theta_{(i,e)})(\tau_1))^\sigma \frac{(u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\tau_2))}{(u_\sigma)_i(\tau_2)},$$

$\xi \in \Lambda_\alpha^i$. Using this, eq. (4.20) implies

$$\tilde{\mathcal{H}}_{m\epsilon, \eta}^\sigma(\theta_{(i,e)}(\Lambda_\beta^i)) \geq \mu(\nu) \frac{(u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\xi))}{(u_\sigma)_i(\xi)} ((D\theta_{(i,e)})(\xi))^\sigma \tilde{\mathcal{H}}_{\epsilon^\sigma, \eta}^\sigma(\Lambda_\beta^i),$$

where $\mu(\nu) = \mu_2(\nu)(\mu_1(\nu))^{-\sigma}$. Now, we sum over $e \in E_i$, and use the fact that

$$\sum_{e \in E_i} ((D\theta_{(i,e)})(\xi))^\sigma (u_\sigma)_{\alpha(i,e)}(\theta_{(i,e)}(\xi)) = (L_\sigma u_\sigma)_i(\xi) = r(L_\sigma)(u_\sigma)_i(\xi)$$

to obtain (4.13). \square

Now we are ready to prove the remaining inequality.

Theorem 4.16. *Assume that the hypotheses H4.1, H4.2, H4.3 and H4.4 are satisfied and let $C_j \subset S_j$, $1 \leq j \leq p$ be the unique invariant list such that*

$$C_i = \bigcup_{(j,e) \in \Gamma_i} \theta_{(j,e)}(C_j) \text{ for } 1 \leq i \leq p.$$

Also assume that $\theta_{(j,e)}|_{C_j}$ is one-to-one for all $(j,e) \in \Gamma$ and that $\theta_{(j,e)}(C_j) \cap \theta_{(j',e')}(C_{j'}) = \emptyset$ whenever $\alpha(j,e) = \alpha(j',e')$ and $(j,e) \neq (j',e')$. Let σ_0 be the unique nonnegative real number such that $r(L_{\sigma_0}) = 1$ and β_0 denote the common Hausdorff dimension of C_i for $1 \leq i \leq p$. Then $\beta_0 \geq \sigma_0$.

Proof. We make the same notational conventions as in Lemma 4.15. If B is any compact subset of C_j , $1 \leq j \leq p$, and $\epsilon > 0$, it is convenient to note that there is a positive, decreasing function $\phi(\epsilon)$, independent of j and B , such that $\tilde{\mathcal{H}}_{\epsilon, \eta}^\sigma(B; C_j) := \tilde{\mathcal{H}}_{\epsilon, \eta}^\sigma(B)$ is defined whenever $0 < \eta < \phi(\epsilon)$. The proof is left to the reader.

By Lemma 4.14 either (a) each C_j is a compact, perfect metric space or (b) each C_j is a single point. In case (b), our assumptions imply that for each i , $1 \leq i \leq p$, there is a unique $(j,e) \in \Gamma$ with $\alpha(j,e) = i$. The linear map L_σ then takes a simple form and the reader can verify that $\sigma_0 = \beta_0 = 0$ in case (b). Thus we shall assume that we are in case (a).

Suppose $\beta_0 < \sigma_0$. Then there exists a $\sigma < \sigma_0$ such that $\mathcal{H}^\sigma(C_i) = 0$ for $1 \leq i \leq p$. This implies for every $\epsilon > 0$, $\mathcal{H}_\epsilon^\sigma(C_i) = 0$ and using (4.7), we have

$$(4.21) \quad \tilde{\mathcal{H}}_\epsilon^\sigma(C_i) = 0 \quad \text{for } \epsilon > 0.$$

Let $(J, E) = [(j_1, e_1), \dots, (j_n, e_n)] \in \Gamma^{(n)}$ and $\theta_{(J,E)} = \theta_{(j_1, e_1)} \circ \dots \circ \theta_{(j_n, e_n)}$. Then $\text{diam}(\theta_{(J,E)}(C_{j_n})) \leq c^n \text{diam}(C_{j_n})$. So, given $\nu > 0$, we can choose $n = n(\nu)$ large enough such that

$$\text{diam}(\theta_{(J,E)}(C_{j_n})) < \nu \text{ for all } (J, E) \in \Gamma^{(n)}.$$

For $1 \leq i \leq p$, we have

$$(4.22) \quad C_i = \bigcup_{(J,E) \in \Gamma_i^{(n)}} \theta_{(J,E)}(C_{j_n})$$

with the union being pairwise disjoint because of the disjointness assumption and the assumption that $\theta_{(j,e)}|_{C_j}$ is one-to-one for $(j,e) \in \Gamma$. By Lemma 4.15, writing $\epsilon^* = \mu_1(\nu)\epsilon$,

$$\mu(\nu)r(L_\sigma)\tilde{\mathcal{H}}_{\epsilon^*,\eta}^\sigma(\theta_{(J,E)}(C_{j_n})) \leq \sum_{e \in E_i} \tilde{\mathcal{H}}_{m\epsilon,\eta}^\sigma(\theta_{(i,e)}(\theta_{(J,E)}(C_{j_n})))$$

where $\mu(\nu) \rightarrow 1$ as $\nu \rightarrow 0$. Since $\mathcal{H}^\sigma(C_j) = 0$ for $1 \leq j \leq p$, using Lemma 4.12, we get $\eta_0 > 0$ such that for $0 < \eta < \eta_0$

$$\tilde{\mathcal{H}}_{m\epsilon,\eta}^\sigma(\theta_{(i,e)}(\theta_{(J,E)}(C_{j_n}))) = \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(\theta_{(i,e)}(\theta_{(J,E)}(C_{j_n})))$$

and

$$\tilde{\mathcal{H}}_{\epsilon^*,\eta}^\sigma(\theta_{(J,E)}(C_{j_n})) = \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(\theta_{(J,E)}(C_{j_n}))$$

Therefore, the previous inequality becomes

$$(4.23) \quad \mu(\nu)r(L_\sigma)\tilde{\mathcal{H}}_{\epsilon^*,\eta}^\sigma(\theta_{(J,E)}(C_{j_n})) \leq \sum_{e \in E_i} \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(\theta_{(i,e)}(\theta_{(J,E)}(C_{j_n})))$$

Now from eq. (4.22), since the union is disjoint, we can choose $\epsilon > 0$ so small that

$$N_\epsilon(\theta_{(J,E)}(C_{j_n})) \cap N_\epsilon(\theta_{(J',E')}(C_{j_n})) = \emptyset$$

for all $(J,E), (J',E') \in \Gamma_i^{(n)}$, $(J,E) \neq (J',E')$. This implies that

$$\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(C_i) = \sum_{(J,E) \in \Gamma_i^{(n)}} \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(\theta_{(J,E)}(C_{j_n})).$$

Therefore, we can sum eq. (4.23) over all $(J,E) \in \Gamma_i^{(n)}$ to obtain

$$\mu(\nu)r(L_\sigma)\tilde{\mathcal{H}}_{\epsilon^*,\eta}^\sigma(C_i) \leq \sum_{(J,E) \in \Gamma_i^{(n)}} \sum_{e \in E_i} \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(\theta_{(i,e)}(\theta_{(J,E)}(C_{j_n}))).$$

Now we sum over $i = 1, 2, \dots, p$ to get

$$\mu(\nu)r(L_\sigma) \sum_{i=1}^p \tilde{\mathcal{H}}_{\epsilon^*,\eta}^\sigma(C_i) \leq \sum_{i=1}^p \sum_{(J,E) \in \Gamma_i^{(n)}} \sum_{e \in E_i} \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(\theta_{(i,e)}(\theta_{(J,E)}(C_{j_n}))).$$

Note that $\theta_{(i,e)}(\theta_{(J,E)}(C_{j_n})) \subset C_{\alpha(i,e)}$, so collecting the terms with $\alpha(i,e) = j$, $1 \leq j \leq p$, we get

$$\mu(\nu)r(L_\sigma) \sum_{i=1}^p \tilde{\mathcal{H}}_{\epsilon^*,\eta}^\sigma(C_i) \leq \sum_{j=1}^p \sum_{(i,e) \in \Gamma_j} \sum_{(J,E) \in \Gamma_i^{(n)}} \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(\theta_{(i,e)}(\theta_{(J,E)}(C_{j_n}))).$$

Since $C_j = \bigcup_{(i,e) \in \Gamma_j} \bigcup_{(J,E) \in \Gamma_i^{(n)}} \theta_{(i,e)}(\theta_{(J,E)}(C_{j_n}))$ with disjoint union, we get

$$(4.24) \quad \mu(\nu)r(L_\sigma) \sum_{i=1}^p \tilde{\mathcal{H}}_{\epsilon^*,\eta}^\sigma(C_i) \leq \sum_{j=1}^p \tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(C_j)$$

Since $\sigma < \sigma_0$, Lemma 4.6 implies $r(L_\sigma) > 1$, so we can choose $\nu > 0$ small enough so that $\mu(\nu)r(L_\sigma) > 1$. But, for $1 \leq i \leq p$, we have $\tilde{\mathcal{H}}_{\epsilon^*,\eta}^\sigma(C_i) > 0$ by the definition of $\tilde{\mathcal{H}}_{\epsilon^*,\eta}^\sigma$. Also, we know that $\tilde{\mathcal{H}}_{\epsilon,\eta}^\sigma(C_i)$ is defined and finite for η small enough. So,

(4.24) cannot be true. Therefore, our initial assumption must be wrong. Thus, $\beta_0 \geq \sigma_0$. \square

Combining Theorem 4.10 and Theorem 4.16, we have proved the following theorem.

Theorem 4.17. *Assume that the hypotheses H4.1, H4.2, H4.3 and H4.4 are satisfied and let $C_j \subset S_j$, $1 \leq j \leq p$ be the unique invariant list such that*

$$C_i = \bigcup_{(j,e) \in \Gamma_i} \theta_{(j,e)}(C_j) \text{ for } 1 \leq i \leq p.$$

Also assume that $\theta_{(j,e)}|_{C_j}$ is one-to-one for all $(j,e) \in \Gamma$ and that $\theta_{(j,e)}(C_j) \cap \theta_{(j',e')}(C_{j'}) = \emptyset$ whenever $\alpha(j,e) = \alpha(j',e')$ and $(j,e) \neq (j',e')$. Let σ_0 be the unique nonnegative real number such that $r(L_{\sigma_0}) = 1$. Then the Hausdorff dimension of each C_i for $1 \leq i \leq p$ is the same, and if β_0 denotes the common Hausdorff dimension of C_i , then $\beta_0 = \sigma_0$.

Remark 4.18. With the aid of Theorem 4.17 and elementary facts about positive linear operators, it is straightforward to prove that the Hausdorff dimension varies continuously with the functions $\theta_{(j,e)}$, $(j,e) \in \Gamma$, in Theorem 4.17. Details will be given elsewhere, but we sketch the basic idea here. For each integer $m \geq 1$ and $(j,e) \in \Gamma$ suppose that $\theta_{(j,e),m} : S_j \rightarrow S_{\alpha(j,e)}$. Assume that, for $m \geq 1$, $\{\theta_{(j,e),m} : (j,e) \in \Gamma\}$ satisfies the assumptions of Theorem 4.17 and let $\sigma_{0,m}$ denote the corresponding Hausdorff dimension. For $(j,e) \in \Gamma$ and $x \in S_j$ assume that $\lim_{m \rightarrow \infty} \theta_{(j,e),m}(x) = \theta_{(j,e)}(x)$ and $\lim_{m \rightarrow \infty} D\theta_{(j,e),m}(x) = D\theta_{(j,e)}(x)$, where these limits define $\theta_{(j,e)}(x)$ and we assume the limits are uniform in $x \in S_j$. Assume the $\theta_{(j,e)}$ satisfies H4.1-H4.4 and that $\theta_{(j,e)}|_{C_j}$ is one-to-one, where $\{C_j : 1 \leq j \leq p\}$ is as in Theorem 2.3. For $\sigma \geq 0$, we have, in the obvious notation, linear operators $L_{\sigma,m}$ corresponding to $\{\theta_{(j,e),m} : (j,e) \in \Gamma\}$ and L_σ corresponding to $\{\theta_{(j,e)} : (j,e) \in \Gamma\}$. If σ_0 denotes the unique value of σ for which $r(L_\sigma) = 1$, then $\lim_{m \rightarrow \infty} \sigma_{0,m} = \sigma_0$. If we assume, in addition, that $\theta_{(j,e)}(C_j) \cap \theta_{(j',e')}(C_{j'}) = \emptyset$ whenever $(j,e) \neq (j',e')$ and $\alpha(j,e) = \alpha(j',e')$, Theorem 4.17 implies that σ_0 is the Hausdorff dimension of C_j for $1 \leq j \leq p$.

5. INFINITE ITERATED FUNCTION SYSTEMS

Let (S,d) be a compact metric space and let I be a countable set. For $i \in I$, let $\theta_i : S \rightarrow S$ be a Lipschitz map with $\text{Lip}(\theta_i) \leq c < 1$. Let I^∞ be the collection of all infinite sequences $\omega = (i_1, i_2, \dots, i_n, \dots)$ such that $i_n \in I$ for each $n \in \mathbb{N}$. Fix $\omega \in I^\infty$. For $x \in S$ and $n \in \mathbb{N}$ define $x_n(\omega) = \theta_{i_1} \circ \theta_{i_2} \circ \dots \circ \theta_{i_n}(x)$. Since $\text{Lip}(\theta_i) \leq c$ for each $i \in I$, if $m \geq n$ we have

$$d(x_n(\omega), x_m(\omega)) \leq c^n \text{diam}(S).$$

It follows that $\{x_n(\omega) : n \geq 1\}$ is a Cauchy sequence. Since (S,d) is a complete metric space, $x_n(\omega)$ must converge to a point $\xi(x,\omega)$ in S . If x and y are any two points in S , then we have

$$d(x_n(\omega), y_n(\omega)) = d(\theta_{i_1} \circ \theta_{i_2} \circ \dots \circ \theta_{i_n}(x), \theta_{i_1} \circ \theta_{i_2} \circ \dots \circ \theta_{i_n}(y)) \leq c^n d(x,y) \leq c^n \text{diam}(S),$$

which goes to zero as n approaches ∞ . This implies that $\xi(x,\omega) = \xi(y,\omega)$. Thus given $\omega \in I^\infty$, we can define $\pi(\omega) = \lim_{n \rightarrow \infty} \theta_{i_1} \circ \theta_{i_2} \circ \dots \circ \theta_{i_n}(x)$, where x is any point in S . Let us consider the set $J := \{\pi(\omega) : \omega \in I^\infty\}$, which will be called the limit set associated to the system $\{\theta_i : i \in I\}$.

Lemma 5.1. *Let J be the limit set defined above. Then*

$$(5.1) \quad J = \bigcup_{i \in I} \theta_i(J).$$

Proof. Since for any $i \in I$, $\theta_i(\pi(\omega)) = \pi(i\omega)$, we see that $\bigcup_{i \in I} \theta_i(J) \subset J$. Also if $\omega = (i_1, i_2, i_3, \dots)$, then $\pi(\omega) = \theta_{i_1}(\pi(i_2, i_3, \dots)) \in \theta_{i_1}(J)$. Thus we have that $J \subset \bigcup_{i \in I} \theta_i(J)$, and the lemma is proved. \square

Remark 5.2. Notice that if I is finite, then the limit set J is compact. Therefore, J is the unique nonempty, compact set satisfying (5.1). If I is infinite, J need not be compact.

Let $I = \mathbb{N}$ and $\theta_i : S \rightarrow S$ be a Lipschitz map with $\text{Lip}(\theta_i) \leq c < 1$ for each $i \in I$. Let J be the limit set associated with this infinite iterated function system. We wish to find a formula for the Hausdorff dimension of J . As in the case of finite iterated function systems, we study a Perron-Frobenius operator $L : C(S) \rightarrow C(S)$ of the form

$$(5.2) \quad (Lf)(t) = \sum_{i=1}^{\infty} b_i(t) f(\theta_i(t))$$

We shall make the following assumptions.

H5.1 For $1 \leq i < \infty$, the function $b_i : S \rightarrow \mathbb{R}$ is nonnegative and continuous. Furthermore, for each $t \in S$, $b(t) = \sum_{i=1}^{\infty} b_i(t) < \infty$ and $b : S \rightarrow \mathbb{R}$ is continuous.

H5.2 For $1 \leq i < \infty$, $\theta_i : S \rightarrow S$ is a Lipschitz map with $\text{Lip}(\theta_i) \leq c < 1$.

If H5.1 is satisfied and the maps θ_i are all continuous, it is easy to verify that L defines a bounded linear map on $C(S)$. We refer the reader to Section 5 of [19] for a detailed discussion of such operators.

Let $K(M, \lambda) = \{f \in C(S) : 0 \leq f(s) \leq f(t) \exp(Md(s, t)^\lambda) \text{ for all } s, t \in S\}$.

Lemma 5.3. *Let (S, d) be a compact metric space and let $L : C(S) \rightarrow C(S)$ be defined by eq. (5.2). Assume that H5.1 and H5.2 are satisfied and assume that there exist $M_0 > 0$ and $\lambda > 0$ such that for each $i \geq 1$, $b_i \in K(M_0, \lambda)$ and no b_i is identically zero. Then there exists $M > 0$ such that $L(K(M, \lambda)) \subset K(M, \lambda)$ and L has a strictly positive eigenvector $u \in K(M, \lambda)$ with eigenvalue $r(L) > 0$.*

Proof. Since $c < 1$ and $\lambda > 0$, we can choose $M > 0$ so that $M_0 + Mc^\lambda \leq M$. We claim that $L(K) \subset K$, where $K := K(M, \lambda)$. Let $f \in K$, $i \geq 1$ and $s, t \in S$. Then we have

$$f(\theta_i(s)) \leq f(\theta_i(t)) \exp(Md(\theta_i(s), \theta_i(t))^\lambda) \leq f(\theta_i(t)) \exp(Mc^\lambda d(s, t)^\lambda) \text{ and} \\ b_i(s) \leq b_i(t) \exp(M_0 d(s, t)^\lambda).$$

Thus

$$(Lf)(s) = \sum_{i=1}^{\infty} b_i(s) f(\theta_i(s)) \\ \leq \sum_{i=1}^{\infty} b_i(t) f(\theta_i(t)) \exp((M_0 + Mc^\lambda) d(s, t)^\lambda) \\ \leq (Lf)(t) \exp(Md(s, t)^\lambda).$$

This proves that $L(K) \subset K$.

By Lemma 3.2, $\{f \in K : \|f\| \leq 1\}$ is equicontinuous, and hence it is compact by Ascoli's theorem. It follows that $L|_K$ is a compact map. The constant function $e = 1$ is in K , so

$$r_K(L) \geq \lim_{n \rightarrow \infty} \|L^n(e)\|^{\frac{1}{n}} = r(L)$$

The opposite inequality is obviously true, so $r_K(L) = r(L)$. By assumption $b(t) = \sum_{i=1}^{\infty} b_i(t) > 0$ for all $t \in S$, and $b : S \rightarrow \mathbb{R}$ is continuous by H5.1. Therefore, there exists $\delta > 0$ with $\sum_{i=1}^{\infty} b_i(t) \geq \delta$. So $L(e) \geq \delta e$ from which it follows that $r(L) \geq \delta > 0$. Since $L|_K$ is a compact map and $r_K(L) = r(L) > 0$, a theorem of Bonsall [1] implies that there exists $u \in K$, $\|u\| = 1$, with $L(u) = ru$, $r = r(L)$. From the definition of $K(M, \lambda)$ it is clear that $u \in K \setminus \{0\}$ implies $u(t) > 0$ for all $t \in S$. \square

Throughout the remainder of this section we shall make the following assumption.

H5.3 Let (S, d) be a compact, perfect metric space and assume that for $1 \leq i < \infty$, $\theta_i : S \rightarrow S$ is an infinitesimal similitude on S and is a Lipschitz map with $\text{Lip}(\theta_i) \leq c < 1$. Assume that there exist $M_0 > 0$ and $\lambda > 0$ such that for all $i \geq 1$, $D\theta_i \in K(M_0, \lambda)$ and $D\theta_i(t) > 0$ for all $t \in S$. Assume, also, that there exist $\sigma > 0$ and $t \in S$ with $\sum_{i=1}^{\infty} (D\theta_i(t))^\sigma < \infty$.

Lemma 5.4. *Assume H5.3 is satisfied. If $\sum_{i=1}^{\infty} (D\theta_i(t_*))^s < \infty$ for some $t_* \in S$ and $s > 0$, then for any $\sigma \geq s$, $\sum_{i=1}^{\infty} (D\theta_i(t))^\sigma < \infty$ for all $t \in S$.*

Proof. Since $D\theta_i(t) < 1$ for all $t \in S$, $(D\theta_i(t))^\sigma \leq (D\theta_i(t))^s$ for $\sigma \geq s$. So $\sum_{i=1}^{\infty} (D\theta_i(t_*))^s < \infty$ implies that $\sum_{i=1}^{\infty} (D\theta_i(t_*))^\sigma < \infty$ for $\sigma \geq s$. Thus we only have to prove that for any $t \in S$, $\sum_{i=1}^{\infty} (D\theta_i(t))^s < \infty$. Let $t \in S$. Since $D\theta_i \in K(M_0, \lambda)$, $D\theta_i(t) \leq D\theta_i(t_*) \exp(M_0 d(t, t_*))^\lambda$. Therefore,

$$(D\theta_i(t))^s \leq (D\theta_i(t_*))^s \exp(sM_0 d(t, t_*)^\lambda)$$

from which the result follows. \square

Assume H5.3 is satisfied. Define for $\sigma > 0$ with $\sum_{i=1}^{\infty} (D\theta_i(t))^\sigma < \infty$,

$$(L_\sigma f)(t) = \sum_{i=1}^{\infty} (D\theta_i(t))^\sigma f(\theta_i(t)).$$

Let $\sigma > 0$ satisfying the above condition be fixed. If H5.1 is satisfied with $b_i(t) := (D\theta_i(t))^\sigma$ for $i \geq 1$, we know that L_σ defines a bounded linear map on $C(S)$. By H5.3, $D\theta_i \in K(M_0, \lambda)$ for some $M_0 > 0$, $\lambda > 0$. This implies that $b_i \in K(\sigma M_0, \lambda)$ for all $i \geq 1$. Therefore, if we choose $0 < M < \infty$ such that $\sigma M_0 + M c^\lambda \leq M$, L_σ maps $K(M, \lambda)$ into itself and has an eigenvector $u_\sigma \in K(M, \lambda)$ with eigenvalue $r(L_\sigma)$.

Let $\sigma_0 = \inf\{\sigma > 0 : \sum_{i=1}^{\infty} (D\theta_i(t))^\sigma < \infty \text{ for all } t \in S\}$. If H5.3 holds, note that if $\sum_{i=1}^{\infty} (D\theta_i(t_*))^\sigma < \infty$ for some $t_* \in S$, then $\sum_{i=1}^{\infty} (D\theta_i(t))^s < \infty$ for all $s \geq \sigma$ and $t \in S$. Assuming that H5.3 holds, there are two possible cases:

Case I: $\sum_{i=1}^{\infty} (D\theta_i(t))^\sigma < \infty$ for all $\sigma > \sigma_0$ and all $t \in S$ but $\sum_{i=1}^{\infty} (D\theta_i(t))^{\sigma_0} = \infty$ for all $t \in S$.

Case II: $\sum_{i=1}^{\infty} (D\theta_i(t))^\sigma < \infty$ for all $\sigma \geq \sigma_0$ and all $t \in S$ but $\sum_{i=1}^{\infty} (D\theta_i(t))^\sigma = \infty$ for all $\sigma < \sigma_0$ and all $t \in S$.

Note that L_σ is defined for $\sigma > \sigma_0$ in case I and for $\sigma \geq \sigma_0$ in case II.

Lemma 5.5. *The map $\sigma \mapsto r(L_\sigma)$ is strictly decreasing and continuous for $\sigma > \sigma_0$ in case I and for $\sigma \geq \sigma_0$ in case II. Also, $r(L_\sigma) \rightarrow 0$ as $\sigma \rightarrow \infty$.*

Proof. The proof that $\sigma \mapsto r(L_\sigma)$ is strictly decreasing and continuous for $\sigma > \sigma_0$ (Case I) or $\sigma \geq \sigma_0$ (Case II) is exactly the same as in the proof of Lemma 4.6, and we omit the details. To prove the last part of the lemma, fix $s > 0$ with $\sup_{t \in S} \sum_{i=1}^{\infty} (D\theta_i(t))^s \leq K < \infty$. Let $\sigma > s$. Then, since $D\theta_i(t) \leq c$,

$$\sum_{i=1}^{\infty} (D\theta_i(t))^\sigma \leq c^{\sigma-s} \sum_{i=1}^{\infty} (D\theta_i(t))^s$$

Therefore, $\|L_\sigma\| \leq \sup_{t \in S} \sum_{i=1}^{\infty} (D\theta_i(t))^\sigma \leq c^{\sigma-s} K$, which implies $\|L_\sigma\| \rightarrow 0$ as $\sigma \rightarrow \infty$ because $c < 1$. Since $r(L_\sigma) \leq \|L_\sigma\|$, the result follows. \square

We should note that in the case of infinite iterated function systems there need not be a value of σ for which $r(L_\sigma) = 1$ because we cannot guarantee a σ for which $r(L_\sigma) \geq 1$. It is possible that $r(L_\sigma) < 1$ for all the values of σ for which L_σ is defined. Let

$$(5.3) \quad \sigma_\infty = \inf\{\sigma > 0 : r(L_\sigma) < 1\}$$

We claim that, under natural further assumptions, the Hausdorff dimension of the invariant set J is equal to σ_∞ .

By Lemma 4.3 and Remark 4.4 we know that given $\epsilon > 0$, there exists a $\mu_i(\epsilon)$ such that for every $t, s \in S$ with $0 < d(s, t) < \epsilon$,

$$\mu_i(\epsilon)^{-1} (D\theta_i)(t) \leq \frac{d(\theta_i(s), \theta_i(t))}{d(s, t)} \leq \mu_i(\epsilon) (D\theta_i)(t)$$

and $\lim_{\epsilon \rightarrow 0^+} \mu_i(\epsilon) = 1$. In the case of finitely many θ_i 's a uniform $\mu(\epsilon)$ satisfying the above property could be chosen by taking the maximum over i . But for the infinite case we cannot guarantee a uniform $\mu(\epsilon)$ which would work for each θ_i . So instead we shall make the assumption that a uniform $\mu(\epsilon)$ can be chosen. For a specific problem we would have to check that this condition is indeed satisfied. For some important examples like complex continued fractions, which have been studied by other authors (see Section 6 of [12]), this condition can be easily verified.

H5.4 Given $\epsilon > 0$, there exists a $\mu(\epsilon) > 1$ such that for any $i \geq 1$ and for every $t, s \in S$ with $0 < d(s, t) < \epsilon$,

$$\mu(\epsilon)^{-1} (D\theta_i)(t) \leq \frac{d(\theta_i(s), \theta_i(t))}{d(s, t)} \leq \mu(\epsilon) (D\theta_i)(t)$$

and $\lim_{\epsilon \rightarrow 0^+} \mu(\epsilon) = 1$.

Now we are ready to prove the upper bound for the Hausdorff dimension of K . The proof is very similar to the proof of Theorem 4.10 but we provide the proof for the sake of completeness.

Theorem 5.6. *Assume that H5.3 and H5.4 are satisfied. Let $\dim(J)$ denote the Hausdorff dimension of the invariant set J and let σ_∞ be as defined in eq. (5.3). Then $\dim(J) \leq \sigma_\infty$.*

Proof. Fix $\epsilon > 0$. Take $\delta > 0$ and $\sigma > 0$. We can choose a covering $\{A_k\}_{k=1}^{\infty}$ of J and points $\xi_k \in A_k$ such that $\text{diam}(A_k) < \epsilon$ and

$$(5.4) \quad \sum_{k=1}^{\infty} u_\sigma(\xi_k) (\text{diam}(A_k))^\sigma \leq \tilde{\mathcal{H}}_\epsilon^\sigma(J) + \delta.$$

Since $J = \bigcup_{i=1}^{\infty} \theta_i(J)$, we have that $\{\theta_i(A_k) : 1 \leq k < \infty, 1 \leq i < \infty\}$ is a covering of J with

$$\text{diam}(\theta_i(A_k)) \leq c \text{diam}(A_k) < c\epsilon.$$

Furthermore, by H5.4 there exists $\mu(\epsilon) > 1$ with $\mu(\epsilon) \rightarrow 1$ as $\epsilon \rightarrow 0+$ such that

$$(5.5) \quad \text{diam}(\theta_i(A_k)) \leq \mu(\epsilon)(D\theta_i)(\xi_k)\text{diam}(A_k)$$

for all $i \geq 1$ and $k \geq 1$.

$$\begin{aligned} \tilde{\mathcal{H}}_{c\epsilon}^{\sigma}(J) &\leq \sum_{k=1}^{\infty} \sum_{i=1}^{\infty} u_{\sigma}(\theta_i(\xi_k))(\text{diam}(\theta_i(A_k)))^{\sigma} \\ &\leq \sum_{k=1}^{\infty} \sum_{i=1}^{\infty} u_{\sigma}(\theta_i(\xi_k))(\mu(\epsilon))^{\sigma}((D\theta_i)(\xi_k))^{\sigma}(\text{diam}(A_k))^{\sigma} \\ &= (\mu(\epsilon))^{\sigma} \sum_{k=1}^{\infty} \left\{ \sum_{i=1}^{\infty} ((D\theta_i)(\xi_k))^{\sigma} u_{\sigma}(\theta_i(\xi_k)) \right\} (\text{diam}(A_k))^{\sigma} \\ &= (\mu(\epsilon))^{\sigma} \sum_{k=1}^{\infty} (L_{\sigma} u_{\sigma})(\xi_k) (\text{diam}(A_k))^{\sigma} \\ &= (\mu(\epsilon))^{\sigma} r(L_{\sigma}) \sum_{k=1}^{\infty} u_{\sigma}(\xi_k) (\text{diam}(A_k))^{\sigma}. \end{aligned}$$

Thus, using (5.4), we get

$$\tilde{\mathcal{H}}_{c\epsilon}^{\sigma}(J) \leq (\mu(\epsilon))^{\sigma} r(L_{\sigma}) \left(\tilde{\mathcal{H}}_{\epsilon}^{\sigma}(J) + \delta \right).$$

Since $c < 1$, $\tilde{\mathcal{H}}_{\epsilon}^{\sigma}(J) \leq \tilde{\mathcal{H}}_{c\epsilon}^{\sigma}(J)$. Also $\delta > 0$ was arbitrary. Therefore,

$$(5.6) \quad \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(J) \leq (\mu(\epsilon))^{\sigma} r(L_{\sigma}) \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(J).$$

For $\sigma > \sigma_{\infty}$, $r(L_{\sigma}) < 1$. Since $\mu(\epsilon) \rightarrow 1$ as $\epsilon \rightarrow 0$, given $\sigma > \sigma_{\infty}$, we can choose $\epsilon > 0$ small so that $(\mu(\epsilon))^{\sigma} r(L_{\sigma}) < 1$. By the definition, $\tilde{\mathcal{H}}_{\epsilon}^{\sigma}(J) < \infty$ because we can take a finite ϵ -cover of the totally bounded set J as it is contained in the compact set S . Thus, if $\sigma > \sigma_{\infty}$, eq.(5.6) can hold only if

$$\lim_{\epsilon \rightarrow 0+} \tilde{\mathcal{H}}_{\epsilon}^{\sigma}(J) = 0.$$

Hence using (4.7), $\lim_{\epsilon \rightarrow 0+} \mathcal{H}_{\epsilon}^{\sigma}(J) = 0$, i.e., $\mathcal{H}^{\sigma}(J) = 0$ for all $\sigma > \sigma_{\infty}$. Thus, by the definition of Hausdorff dimension, $\dim(J) \leq \sigma_{\infty}$. \square

To prove the other half, $\dim(J) \geq \sigma_{\infty}$, we shall consider the infinite iterated system as the limit of finite iterated systems and use the result that we have for the finite case.

Define for $N \geq 1$ and $f \in C(S)$,

$$(L_{\sigma, N} f)(t) = \sum_{i=1}^N (D\theta_i(t))^{\sigma} f(\theta_i(t)).$$

For $N \geq 1$, let J_N be the unique nonempty, compact invariant set satisfying

$$J_N = \bigcup_{i=1}^N \theta_i(J_N)$$

and let σ_N be the unique positive real number such $r(L_{\sigma_N, N}) = 1$. We shall assume the following hypothesis.

H5.5 For each $N \geq 1$, $\theta_i(J_N) \cap \theta_j(J_N) = \emptyset$ for $1 \leq i < j \leq N$, and $\theta_i|_{J_N}$ is one-to-one for $1 \leq i \leq N$.

By Theorem 4.17 we know that, assuming H5.3 and H5.5, $\dim(J_N) = \sigma_N$.

Lemma 5.7. *For each $N \geq 1$, $J_N \subset J_{N+1}$ and $J_N \subset J$. Hence, $\dim(J) \geq \dim(J_N) = \sigma_N$ for all $N \geq 1$.*

Proof. By Remark 5.2, we know that $J_N = \{\pi(\omega) : \omega \in \{1, 2, \dots, N\}^\infty\}$ for any $N \geq 1$. It follows that $J_N \subset J_{N+1}$ for all $N \geq 1$. Also since $J = \{\pi(\omega) : \omega \in \mathbb{N}^\infty\}$, we have $J_N \subset J$ for all $N \geq 1$. \square

It is easy to see that $\|L_\sigma - L_{\sigma, N}\| \rightarrow 0$ as $N \rightarrow \infty$ for $\sigma > \sigma_0$ in case I and for $\sigma \geq \sigma_0$ in case II.

Remark 5.8. Let X be a real or complex Banach space and $L : X \rightarrow X$, $L_k : X \rightarrow X$, $k \geq 1$ be bounded linear operators. Assume that $\lim_{k \rightarrow \infty} \|L_k - L\| = 0$. Then we have that $\limsup_{k \rightarrow \infty} r(L_k) \leq r(L)$. But, in general, it is not true that $\lim_{k \rightarrow \infty} r(L_k) \rightarrow r(L)$. In fact, Kakutani has given an example of a sequence of bounded linear operators L_k on a Hilbert space which converges in the operator norm to an operator L and satisfies $r(L_k) = 0$ for all $k \geq 1$ and $r(L) > 0$. The example can be found on pages 282-283 of [20]. If, in addition, we know that $\rho(L) < r(L)$, where $\rho(L)$ is the essential spectral radius of L , then it is true that $r(L_k) \rightarrow r(L)$. To see this, note that by using the natural extension of L to the complexification of X , we can assume that X is a complex Banach space. If $\sigma(L)$ denotes the spectrum of L , recall that $\sigma(L) \cap \{z \in \mathbb{C} : |z| > \rho(L)\}$ consists of isolated points each of which is an eigenvalue of L of finite algebraic multiplicity. Then exactly the argument on pages 227-228 of [17] proves that $r(L_k) \rightarrow r(L)$.

The following lemma is known. The proof is included for the reader's convenience.

Lemma 5.9. *Let (S, d) be a compact metric space and suppose that $L : X = C(S) \rightarrow X$ is a positive bounded linear map, i.e., $f(t) \geq 0$ for all $t \in S$ implies that $(Lf)(t) \geq 0$ for all $t \in S$. Let e denotes the function identically equal to 1. If $r(L)$ denotes the spectral radius of L , we have $r(L) = \lim_{n \rightarrow \infty} \|L^n e\|^{\frac{1}{n}}$. Furthermore, if $u \in X$ is such that $u(t) > 0$ for all $t \in S$, then $r(L) = \lim_{n \rightarrow \infty} \|L^n u\|^{\frac{1}{n}}$. Finally, if $Lu = ru$ with $u(t) > 0$ for all $t \in S$, then $r(L) = r$.*

Proof. We shall write $f \leq g$ to mean $f(t) \leq g(t)$ for all $t \in S$. Since L is linear and maps nonnegative functions to nonnegative functions, it follows that $Lf \leq Lg$ whenever $f \leq g$. If $f \in X$ with $\|f\| \leq 1$, we have $-e \leq f \leq e$. So, $-L^n e \leq L^n f \leq L^n e$ which implies that $|(L^n f)(t)| \leq |L^n e(t)|$ for all $t \in S$. Thus $\|L^n f\| \leq \|L^n e\|$ whenever $\|f\| \leq 1$ which gives $\|L^n\| = \|L^n e\|$. Taking the n th root and taking the limit as n goes to ∞ , we get $r(L) = \lim_{n \rightarrow \infty} \|L^n e\|^{\frac{1}{n}}$.

Now let $u \in X$ such that $u(t) > 0$ for all $t \in S$. Since S is compact, there exist $0 < m < M < \infty$ such that $me \leq u \leq Me$. This implies $mL^n e \leq L^n u \leq ML^n e$, so $m\|L^n e\| \leq \|L^n u\| \leq M\|L^n e\|$. Taking the n th root and taking the limit, we get $\lim_{n \rightarrow \infty} \|L^n u\|^{\frac{1}{n}} = \lim_{n \rightarrow \infty} \|L^n e\|^{\frac{1}{n}} = r(L)$. To see the last part, note that $Lu = ru$ implies $L^n u = r^n u$. So, $\|L^n u\|^{\frac{1}{n}} = r\|u\|^{\frac{1}{n}}$. Since $\|u\| > 0$, $\|u\|^{\frac{1}{n}} \rightarrow 1$. Thus we get $r(L) = r$. \square

Corollary 5.10. *Let (S, d) be a compact metric space and suppose that $L : X = C(S) \rightarrow X$ and $L_k : X \rightarrow X$, $k \geq 1$, are positive bounded linear maps. Assume that $\|L_k - L\| \rightarrow 0$ as $k \rightarrow \infty$. Suppose $Lu = ru$ with $u(t) > 0$ for all $t \in S$. Then $r(L_k) \rightarrow r(L) = r$.*

Proof. First we know that $r = r(L)$ by Lemma 5.9. Now we have that $\|L_k u - Lu\| \rightarrow 0$ as $k \rightarrow \infty$. Because u is strictly positive, given $\delta > 0$, there exists k_0 such that $L_k u \geq (1 - \delta)ru$ for all $k \geq k_0$. This implies for any $n \geq 1$, $\|L_k^n u\| \geq (1 - \delta)^n r^n \|u\|$ for $k \geq k_0$. By Lemma 5.9, $r(L_k) = \lim_{n \rightarrow \infty} \|L_k^n u\|^{\frac{1}{n}} \geq (1 - \delta)r$ for $k \geq k_0$. Since $\delta > 0$ was arbitrary, $\liminf_{k \rightarrow \infty} r(L_k) \geq r$. Thus we are done because we always have $\limsup_{k \rightarrow \infty} r(L_k) \leq r(L)$. \square

Lemma 5.11. *Assume that H5.3 and H5.4 are satisfied. Then we have that $r(L_{\sigma, N}) \uparrow r(L_\sigma)$ as $N \rightarrow \infty$ for $\sigma > \sigma_0$ in case I and for $\sigma \geq \sigma_0$ in case II.*

Proof. Let $\sigma > \sigma_0$ if we are in case I or $\sigma \geq \sigma_0$ if we are in case II. If $b_i(t) = (D\theta_i(t))^\sigma$, then H5.3 implies that $b_i \in K(\sigma M_0, \lambda)$, and using this fact and H5.3, the reader can verify that $b(t) := \sum_{i=1}^{\infty} b_i(t) > 0$ for all $t \in S$ and b is continuous. It follows that L_σ defines a positive bounded linear operator on $C(S)$. Clearly $L_{\sigma, N} e \leq L_{\sigma, N+1} e$, so $\|L_{\sigma, N} e\| \leq \|L_{\sigma, N+1} e\|$. By Lemma 5.9, we get that $r(L_{\sigma, N}) \leq r(L_{\sigma, N+1})$. It is easy to see, using H5.3, that $\|L_\sigma - L_{\sigma, N}\| \rightarrow 0$ as $N \rightarrow \infty$. Since we know that L_σ has a strictly positive eigenvector with eigenvalue $r(L_\sigma)$, by Corollary 5.10, we must have $r(L_{\sigma, N}) \rightarrow r(L_\sigma)$. \square

Now we can prove the lower bound for the Hausdorff dimension of J .

Theorem 5.12. *Assume that H5.3, H5.4 and H5.5 are satisfied. Let $\dim(J)$ denote the Hausdorff dimension of J and σ_∞ be as defined in eq. (5.3). Then $\dim(J) = \sigma_\infty$.*

Proof. By Theorem 5.6, it suffices to prove that $\dim(J) \geq \sigma_\infty$. If $\sigma_0 < \sigma_\infty$ then for $\sigma_0 < \sigma < \sigma_\infty$, L_σ is defined and $r(L_\sigma) > 1$ by the definition of σ_∞ and the fact that $\sigma \mapsto r(L_\sigma)$ is strictly decreasing. So, by Lemma 5.11, there exists N_σ such that $r(L_{\sigma, N}) > 1$ for all $N \geq N_\sigma$. Since $r(L_{\sigma_N, N}) = 1$, $\sigma_N > \sigma$ for all $N \geq N_\sigma$. Therefore, by Lemma 5.7, we have that $\dim(J) \geq \dim(J_{N_\sigma}) = \sigma_{N_\sigma} > \sigma$. Since this is true for any σ with $\sigma_0 < \sigma < \sigma_\infty$, it follows that $\dim(J) \geq \sigma_\infty$. If $\sigma_0 = \sigma_\infty$ then for $\sigma < \sigma_\infty$, $\sum_{i=1}^{\infty} (D\theta_i(t))^\sigma = \infty$, so for large N (depending on σ), $\sum_{i=1}^N (D\theta_i(t))^\sigma > 1$ for all $t \in S$. This implies that $r(L_{\sigma, N}) > 1$, so $\sigma_N > \sigma$. Thus $\dim(J) > \sigma$ for all $\sigma < \sigma_\infty$, and hence $\dim(J) \geq \sigma_\infty$. This completes the proof of the theorem. \square

To illustrate Theorem 5.12, we discuss a special infinite iterated function system that is generated by complex continued fractions. This has been studied in Section 6 of [12]. We show that our theory is applicable to this particular example.

Example 5.13. (*Complex continued fractions*) Let I be an infinite subset of $\{m + ni : m \in \mathbb{N}, n \in \mathbb{Z}\}$, where \mathbb{Z} is the set of integers and \mathbb{N} is the set of positive integers. Let $X \subset \mathbb{C}$ be the closed disc centered at the point $\frac{1}{2}$ with radius $\frac{1}{2}$. For $b \in I$ one can easily verify that $\theta_b(X) \subset X$, where

$$\theta_b(z) = \frac{1}{b+z}.$$

The mappings θ_b , $b \in I$, may not all be strict contractions in the Euclidean metric; θ_1 is not a strict contraction because $|\theta'_1(0)| = 1$. Therefore we consider the system

$\{\theta_b \circ \theta_c : b, c \in I\}$. It is easy to verify that $\theta_b \circ \theta_c$ is a strict contraction for each $b, c \in I$ with a uniform Lipschitz constant $\kappa < 1$. Let J be the limit set for this system. First note that $\theta_b(z) = \theta_c(w)$ implies that $|z - w| = |b - c|$. So, if $|b - c| > 1$ then $\theta_b(X) \cap \theta_c(X) = \emptyset$. Furthermore, if $|b - c| = 1$ then $\theta_b(z) = \theta_c(w)$ implies that z and w belong to the boundary of X and $|z - w| = 1$.

Lemma 5.14. $\theta_b \circ \theta_c(X)$ is contained in the interior of X .

Proof. First we claim that $\theta_b(z) \in \partial X$ implies $z = 0$. Let $b = m + ni$, $m \in \mathbb{N}$, $n \in \mathbb{Z}$ and $z = x + yi \in X$. Then $\theta_b(z) \in \partial X$ implies $|\frac{1}{b+z} - \frac{1}{2}| = \frac{1}{2}$ which implies $|2 - b - z| = |b + z|$. Therefore, $(2 - m - x)^2 + (n + y)^2 = (m + x)^2 + (n + y)^2$ which implies $m + x = 1$, i.e., $x = 1 - m$. Since $m \geq 1$ and $x \geq 0$ for $z \in X$, it follows that $m = 1$ and $x = 0$. But $x = 0$ implies that $z = 0$.

Now suppose that $\theta_b \circ \theta_c(z) \in \partial X$ for some $z \in X$. Then, by the above claim, $\theta_c(z) = 0$, which is impossible by the definition of $\theta_c(z)$. \square

Let us verify H5.5. By the previous lemma, we know that for any $b, c \in I$, $\theta_b \circ \theta_c(X)$ is a compact set contained in the interior of X . So if we take finitely many b_i, c_i , the union of the images would still be a compact subset of the interior of X . So, for any N , J_N is a compact subset of the interior of X which means $\text{diam}(K_N) < 1$. We claim that if $\hat{X} \subset \text{int}(X)$ with $\text{diam}(\hat{X}) < 1$ then $\theta_{b_1} \circ \theta_{c_1}(\hat{X}) \cap \theta_{b_2} \circ \theta_{c_2}(\hat{X}) = \emptyset$ for any $(b_1, c_1) \neq (b_2, c_2)$. Suppose instead that $\theta_{b_1}(\theta_{c_1}(z)) = \theta_{b_2}(\theta_{c_2}(w))$ with $z, w \in \hat{X}$. This implies that $b_1 + \theta_{c_1}(z) = b_2 + \theta_{c_2}(w)$. If $b_1 = b_2$, this would imply that $\theta_{c_1}(z) = \theta_{c_2}(w)$, i.e., $c_1 + z = c_2 + w$, which is impossible because $|z - w| < 1$ and $|c_1 - c_2| \geq 1$. If $b_1 \neq b_2$, we must have $|\theta_{c_1}(z) - \theta_{c_2}(w)| = 1$ which is possible only if both $\theta_{c_1}(z)$ and $\theta_{c_2}(w)$ belong to the boundary of X , which is possible only if $z = w = 0$. This is a contradiction to the fact that \hat{X} is in the interior of X . Thus the disjointness condition in H5.5 is satisfied. Also for any $b \in I$, the map θ_b is clearly one-to-one.

For $b \in I$, we have $D\theta_b(z) = |\theta'_b(z)| = \frac{1}{|z+b|^2}$. So, $D\theta_b(z) > 0$ for all $z \in X$. We claim that there exists $0 < M_0 < \infty$ such that $D\theta_b \in K(M_0, \lambda)$ with $\lambda = 1$. Let $z, w \in X$. We have

$$\begin{aligned} (D\theta_b)(z) &\leq (D\theta_b)(w) \exp(M_0|z - w|) \\ \Leftrightarrow \frac{1}{|z + b|^2} &\leq \frac{1}{|w + b|^2} \exp(M_0|z - w|) \\ &\Leftrightarrow M_0 \geq 2 \frac{\ln \left| \frac{w+b}{z+b} \right|}{|w - z|} \end{aligned}$$

But $\ln \left| \frac{w+b}{z+b} \right| = \ln \left| 1 + \frac{w-z}{z+b} \right| \leq \ln \left(1 + \frac{|w-z|}{|z+b|} \right) \leq \frac{|w-z|}{|z+b|}$. Therefore, $2 \frac{\ln \left| \frac{w+b}{z+b} \right|}{|w-z|} \leq \frac{2}{|z+b|} \leq 2$. So, we can choose M_0 independent of $b \in I$.

To complete the verification of H5.3 note that

$$\sum_{b \in I} (D\theta_b)^\sigma(0) = \sum_{b \in I} \frac{1}{|b|^{2\sigma}} \leq \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{N}} \frac{1}{(m^2 + n^2)^{2\sigma}}$$

which converges for $\sigma > 1$.

To verify H5.4 it is enough to show that given $\epsilon > 0$ there exist $\mu_1(\epsilon)$ and $\mu_2(\epsilon)$ such that $\lim_{\epsilon \rightarrow 0+} \mu_1(\epsilon) = \lim_{\epsilon \rightarrow 0+} \mu_2(\epsilon) = 1$ and if $0 < |z - w| < \epsilon$ then $\mu_1(\epsilon) \leq \frac{1}{(D\theta_b)(z)} \frac{|\theta_b(z) - \theta_b(w)|}{|z - w|} \leq \mu_2(\epsilon)$ for all $b \in I$. But using $\theta_b(z) = \frac{1}{z+b}$, we get

$\frac{1}{(D\theta_b)(z)} \frac{|\theta_b(z) - \theta_b(w)|}{|z-w|} = \left| \frac{z+b}{w+b} \right| = \left| 1 + \frac{z-w}{w+b} \right|$, which is bounded between $1 - \frac{|z-w|}{|w+b|}$ and $1 + \frac{|z-w|}{|w+b|}$. Since for any $w \in X$ $|w+b|^{-1} = |\theta_b(w)| \leq 1$, taking $\mu_1(\epsilon) = 1 - \epsilon$ and $\mu_2(\epsilon) = 1 + \epsilon$ does the job.

Thus we see that all the hypotheses of the theorem are satisfied for this particular example and hence the Hausdorff dimension of the limit set J is given by the value of σ_∞ . Simpler versions of these arguments show that the results of Section 4 are applicable if I is a finite set.

In the next section we shall see that sometimes it is important to look at another metric rather than the Euclidean metric.

6. CHOICE OF APPROPRIATE METRIC

We need to recall the definition of the Carathéodory-Reiffen-Finsler (CRF) metric on bounded domains in Banach spaces. Let G be a bounded domain in a complex Banach space X and let U denote the open unit disc in \mathbb{C} . Let $Hol(G, U)$ be the family of all holomorphic functions $f : G \rightarrow U$. Define $\alpha : G \times X \rightarrow \mathbb{R}$ by

$$\alpha(x, v) = \sup\{|Dg(x)v| : g \in Hol(G, U)\}$$

where $Dg(x)$ denotes the Fréchet derivative of g at x . Given any two points x and y in G , consider the family of curves $\gamma : [0, 1] \rightarrow G$ that have piecewise continuous derivatives and $\gamma(0) = x$, $\gamma(1) = y$. Call such a curve admissible and define its length by

$$L(\gamma) = \int_0^1 \alpha(\gamma(t), \gamma'(t)) dt.$$

We now define the distance between x and y by

$$\rho(x, y) = \inf\{L(\gamma) : \gamma \text{ is admissible with } \gamma(0) = x \text{ and } \gamma(1) = y\}.$$

ρ is called the CRF metric on G . For a detailed discussion of the CRF metric we refer the reader to [6].

Let G be a bounded open set in \mathbb{C} and let $\theta : G \rightarrow G$ be a holomorphic map such that $\overline{\theta(G)}$ is a compact subset of G . If ρ denotes the CRF metric on G then it is known (see Theorem 13.1 in [6]) that θ is a strict contraction on G with respect to the CRF metric ρ . Also, on a compact subset C of G , ρ is a complete metric and is equivalent to the standard Euclidean metric, i.e., there exist positive constants m and M such that

$$m|z - w| \leq \rho(z, w) \leq M|z - w| \text{ for all } z, w \text{ in } C.$$

Let G be a bounded open set in \mathbb{C} and assume for $1 \leq i \leq N$ that $\theta_i : G \rightarrow G$ is a holomorphic map such that $C_i = \overline{\theta_i(G)}$ is a compact subset of G and $\theta_i'(z) \neq 0$ for all $z \in G$. Define $C := \bigcup_{i=1}^n C_i$. Then, by Theorem 2.3 and the above remarks, there exists a unique nonempty compact set K with $K = \bigcup_{i=1}^N \theta_i(K)$. For $k \geq 1$, define $\mathcal{I}_k = \{I = (i_1, i_2, \dots, i_k) : 1 \leq i_j \leq N \text{ for } 1 \leq j \leq k\}$. For $I = (i_1, i_2, \dots, i_k) \in \mathcal{I}_k$, define $\theta_I = \theta_{i_k} \circ \dots \circ \theta_{i_2} \circ \theta_{i_1}$. It is easy to see that $K = \bigcup_{I \in \mathcal{I}_k} \theta_I(K)$. We claim that, for large k , θ_I is a strict contraction map with respect to the Euclidean metric. Suppose $z, w \in C$, $z \neq w$. Then

$$\frac{|\theta_I(z) - \theta_I(w)|}{|z - w|} \leq \frac{\frac{1}{m}\rho(\theta_I(z), \theta_I(w))}{\frac{1}{M}\rho(z, w)} \leq \frac{M}{m}c^k,$$

where $c < 1$ is the maximum of the contraction ratios of the maps θ_i , $1 \leq i \leq N$, with respect to the metric ρ . If we choose k large enough so that $\frac{M}{m}c^k < 1$, then it follows that θ_I is a contraction map for all $I \in \mathcal{I}_k$ with respect to the Euclidean metric. Thus, if $\theta_I(K) \cap \theta_J(K) = \emptyset$ for $I, J \in \mathcal{I}_k$, $I \neq J$ (which is certainly true if $\{\theta_i(K)\}_{i=1}^N$ are pairwise disjoint and $\theta_i|_K$ is one-to-one for $1 \leq i \leq N$), the Hausdorff dimension of the invariant set K is given by Theorem 1.2 by considering the iterated function system given by the maps $\{\theta_I\}_{I \in \mathcal{I}_k}$ and the standard Euclidean metric. Note that in this case $(D\theta_I)(z)$ is nothing but $|\theta'_I(z)|$. If we write, for $\sigma \geq 0$, $(L_\sigma f)(z) = \sum_{i=1}^N |\theta'_i(z)|^\sigma f(\theta_i(z))$ and $(\tilde{L}_\sigma f)(z) = \sum_{I \in \mathcal{I}_k} |\theta'_I(z)|^\sigma f(\theta_I(z))$, where k is as chosen above, then it is easy to see that $\tilde{L}_\sigma = L_\sigma^k$. It follows from Lemma 6.1 below that $r(\tilde{L}_{\sigma_0}) = 1$ if and only if $r(L_{\sigma_0}) = 1$. Thus, to find the Hausdorff dimension of the invariant set K it is enough to find σ_0 such that $r(L_{\sigma_0}) = 1$.

The following lemma is well known, and the proof is given only for completeness.

Lemma 6.1. *Let X be a Banach space, $L : X \rightarrow X$ be a bounded linear map and $k \geq 1$ be a positive integer. Then $r(L^k) = (r(L))^k$, where $r(L)$ denotes the spectral radius of L .*

Proof. We have that

$$\begin{aligned} r(L^k) &= \lim_{n \rightarrow \infty} \|(L^k)^n\|^{1/n} \\ &= \lim_{n \rightarrow \infty} (\|L^{kn}\|^{1/kn})^k \\ &= \left(\lim_{n \rightarrow \infty} \|L^{kn}\|^{1/kn} \right)^k \\ &= (r(L))^k. \end{aligned}$$

□

REFERENCES

1. F. F. Bonsall, *Linear operators in complete positive cones*, Proc. London Math. Soc. **8** (1958), 53–75.
2. R. T. Bumby, *Hausdorff dimensions of Cantor sets*, J. Reine Angew. Math. **331** (1982), 192–206.
3. ———, *Hausdorff dimension of sets arising in number theory*, Number theory (New York, 1983–84), Lecture Notes in Math., vol. 1135, Springer, Berlin, 1985, pp. 1–8.
4. G. Edgar, *Measure, Topology and Fractal Geometry*, second ed., Undergraduate Texts in Mathematics, Springer, New York, 2008.
5. K. J. Falconer, *The Geometry of Fractal Sets*, Cambridge Univ. Press, 1985.
6. K. Goebel and S. Reich, *Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 83, Marcel Dekker Inc., New York, 1984.
7. J. E. Hutchinson, *Fractals and self-similarity*, Indiana Univ. Math. J. **30** (1981), no. 5, 713–747.
8. M. G. Krein and M. A. Rutman, *Linear operators leaving invariant a cone in a Banach space (in Russian)*, Uspehi Matem. Nauk **3** (1948), no. 1(23), 3–95.
9. J. Mallet-Paret and R. D. Nussbaum, *Eigenvalues for a class of homogeneous cone maps arising from max-plus operators*, Discrete Contin. Dyn. Syst. **8** (2002), no. 3, 519–562.
10. ———, *Generalizing the Krein-Rutman theorem, measures of noncompactness and the fixed point index*, J. Fixed Point Theory and Applications **7** (2010), 103–143.
11. P. Mattila, *Geometry of Sets and Measures in Euclidean Spaces*, Cambridge Univ. Press, 1995.
12. R. D. Mauldin and M. Urbański, *Dimensions and measures in infinite iterated function systems*, Proc. London Math. Soc. (3) (1996), no. 1, 105–154.

13. R. D. Mauldin and S. C. Williams, *Hausdorff dimension in graph directed constructions*, Trans. Amer. Math. Soc. **309** (1988), no. 2, 811–829.
14. S. Mazur and S. Ulam, *Sur les transformations isométriques d'espaces vectoriels normés*, C. R. Acad. Sci. Paris **194** (1932), 946–948.
15. P. A. P. Moran, *Additive functions of intervals and Hausdorff measure*, Proc. Cambridge Philos. Soc. **42** (1946), 15–23.
16. J. R. Munkres, *Topology*, second ed., Prentice Hall, 2000.
17. R. D. Nussbaum, *Periodic solutions of some nonlinear integral equations*, Dynamical systems (Proc. Internat. Sympos., Univ. Florida, Gainesville, Fla., 1976), Academic Press, New York, 1977, pp. 221–249.
18. ———, *Eigenvectors of nonlinear positive operators and the linear Krein-Rutman theorem*, Fixed point theory, Lecture Notes in Math., vol. 886, Springer, Berlin, 1981, pp. 309–330.
19. ———, *Periodic points of positive linear operators and Perron-Frobenius operators*, Integral Equations Operator Theory **39** (2001), no. 1, 41–97.
20. C. E. Rickart, *General Theory of Banach Algebras*, Robert E. Krieger Publishing Co., New York, 1974.
21. H. H. Schaefer, *Banach Lattices and Positive Operators*, Springer Verlag, New York, 1974.
22. A. Schief, *Self-similar sets in complete metric spaces*, Proc. Amer. Math. Soc. **124** (1996), no. 2, 481–490.
23. J. Väisälä, *A proof of the Mazur-Ulam theorem*, Amer. Math. Monthly **110** (2003), no. 7, 633–635.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW JERSEY 08854
E-mail address: `nussbaum@math.rutgers.edu`

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW JERSEY 08854
E-mail address: `amitpriy@math.rutgers.edu`

MATHEMATISCH INSTITUUT, UNIVERSITEIT LEIDEN, LEIDEN, THE NETHERLANDS
E-mail address: `verduyn@math.leidenuniv.nl`