

The first problem sketches a proof of a result you are probably familiar with: the binomial theorem.

#1(a) If  $n > i \geq 0$  are integers, define  $\binom{n}{i} = \frac{n!}{i!(n-i)!}$ . (Recall that  $n! = (n)(n-1)\dots(3)(2)(1)$  if  $n \geq 1$  and that  $0! = 1$ .) Prove that for  $n \geq i \geq 1$  we have

$$\binom{n}{i} = \binom{n-1}{i} + \binom{n-1}{i-1}$$

and that  $\binom{n}{i}$  is an integer.

(b) Let  $a, b, n \in \mathbf{Z}, n \geq 0$ . Prove that

$$(a+b)^n = \sum_{i=0}^n \binom{n}{i} a^i b^{n-i}.$$

The next problem uses the binomial theorem to derive a celebrated result. We will see another way to prove this result later in the course (using material of Chapter 7).

#2(a) Let  $p$  be a prime and  $0 < i < p$ . Prove that  $p$  divides  $\binom{p}{i}$ .

(b) Let  $p$  be a prime and  $a, b \in \mathbf{Z}$ . Prove that  $p$  divides  $(a+b)^p - a^p - b^p$ .

(c) Prove:

**Fermat's Little Theorem:** Let  $p$  be a prime and  $a \in \mathbf{Z}$ . Then  $p$  divides  $a^p - a$ .

It is frequently interesting to investigate whether all the hypotheses of a result are really needed and, if so, why. This explains the next problem.

#3 Does the Fermat's Little Theorem continue to hold if the hypothesis that  $p$  is prime is omitted? Justify your answer by giving a proof or an example.

Now some unrelated problems:

#4 Let  $n$  be an integer that is not divisible by 2 or 5. Let  $J_m$  denote the integer  $11\dots1$  where there are  $m$  1's (thus  $J_m = 1 + 10 + (10)^2 + \dots + (10)^{m-1}$ ). Prove that  $n$  divides  $J_m$  for some  $m$ .

#5 Let  $r, s, t \in \mathbf{Z}$  be such that the only positive integer dividing  $r, s,$  and  $t$  is 1. Prove that there are integers  $a, b, c$  such that  $ar + bs + ct = 1$ . (Note that this generalizes the result for two integers that was proved in class.)