

# ON THE PRESCRIBING $\sigma_2$ CURVATURE EQUATION ON $\mathbb{S}^4$

S.-Y. ALICE CHANG, ZHENG-CHAO HAN, AND PAUL YANG

ABSTRACT. Prescribing  $\sigma_k$  curvature equations are fully nonlinear generalizations of the prescribing Gaussian or scalar curvature equations. Given a positive function  $K$  to be prescribed on the 4-dimensional round sphere. We obtain asymptotic profile analysis for potentially blowing up solutions to the  $\sigma_2$  curvature equation with the given  $K$ ; and rule out the possibility of blowing up solutions when  $K$  satisfies a non-degeneracy condition. We also prove uniform a priori estimates for solutions to a family of  $\sigma_2$  curvature equations deforming  $K$  to a positive constant under the same non-degeneracy condition on  $K$ , and prove the existence of a solution using degree argument to this deformation involving fully nonlinear elliptic operators under an additional, natural degree condition on a finite dimensional map associated with  $K$ .

## 1. DESCRIPTION OF MAIN RESULTS

Our main results in this paper are (potential) blow up profile analysis, a priori estimates and existence of admissible solutions  $w$  to the  $\sigma_2$  curvature equation

$$(1) \quad \sigma_2(g^{-1} \circ A_g) = K(x),$$

on  $\mathbb{S}^4$ , where  $g = e^{2w(x)}g_c$  is a metric conformal to  $g_c$ , with  $g_c$  being the canonical background metric on the round sphere  $\mathbb{S}^4$ ,  $A_g$  is the the Weyl-Schouten tensor of the metric  $g$ ,

$$(2) \quad \begin{aligned} A_g &= \frac{1}{n-2} \left\{ Ric - \frac{R}{2(n-1)}g \right\} \\ &= A_{g_c} - \left[ \nabla^2 w - dw \otimes dw + \frac{1}{2} |\nabla w|^2 g_c \right], \end{aligned}$$

and  $\sigma_k(\Lambda)$ , for any 1–1 tensor  $\Lambda$  on an  $n$ –dimensional vector space and  $k \in \mathbb{N}$ ,  $0 \leq k \leq n$ , is the  $k$ -th elementary symmetric function of the eigenvalues of  $\Lambda$ ;  $K(x)$  is a given function

---

The research of the first and third author is partially supported by NSF through grant DMS-0758601; the first author also gratefully acknowledges partial support from the Minerva Research Foundation and The Charles Simonyi Endowment fund during the academic year 08-09 while visiting Institute of Advanced Study, Princeton. The research of the second author is partially supported by NSF through grant DMS-0103888 and by a Rutgers University Research Council Grant(202132).

on  $\mathbb{S}^4$  with some appropriate assumptions, and an *admissible* solution is defined to be a  $C^2(M)$  solution  $w$  to (1) such that for all  $x \in \mathbb{S}^4$ ,  $A_g(x) \in \Gamma_k^+$ , namely,  $\sigma_j(g^{-1} \circ A_g) > 0$  for  $1 \leq j \leq k$ . Note that  $\sigma_1(A_g)$  is simply a positive constant multiple of the scalar

curvature of  $g$ , so  $A_g$  in the  $\Gamma_k^+$  class is a generalization of the notion that the scalar curvature  $R_g$  of  $g$  having a fixed  $+$  sign.

Note that, since

$$\sigma_2(g^{-1} \circ A_g) = e^{-4w} \sigma_2(g_c^{-1} \circ A_g),$$

so (1) is equivalent to

$$(3) \quad \sigma_2(g_c^{-1} \circ \left[ A_{g_c} - \nabla^2 w + dw \otimes dw - \frac{1}{2} |\nabla w|^2 g_c \right]) = K(x) e^{4w(x)}.$$

It is well known that (3) is *elliptic* at an admissible solution; and in fact, any solution  $w$  to (3) on  $\mathbb{S}^4$  is admissible. There have been a large number of papers on problems related to the  $\sigma_k$  curvature since the work [V00a] of Viaclovsky a decade ago. It is inadequate to do even a short survey of recent work in the introductory remarks here. We will instead refer the reader to recent surveys [V06] by Viaclovsky and [CC07] by Chang and Chen.

As alluded to above, a similar problem to (3) for the  $\sigma_1$  curvature was a predecessor to (3). More specifically, if we prescribe a function  $K(x)$  on the round  $n$ -dimensional sphere  $(\mathbb{S}^n, g_c)$  to be the scalar curvature of a metric  $g = e^{2w} g_c$  pointwise conformal to  $g_c$ , then  $w$  satisfies

$$(4) \quad 2(n-1)\Delta_{g_c} w + (n-1)(n-2)|\nabla w|^2 = R_{g_c} - K(x)e^{2w}.$$

Similar equation can be formulated for a general manifold. One difference between (4) and (3) is that (4) is semilinear in  $w$ , while (3) is fully nonlinear in  $w$ . (4) takes on the familiar form

$$(5) \quad 2\Delta_{g_c} w = R_{g_c} - K(x)e^{2w},$$

when  $n = 2$ , which is the Nirenberg problem. The  $n \geq 3$  case of (4) is often written

in terms of a different variable  $u = e^{(n-2)w/2}$ , which would render the equation in the familiar form

$$(6) \quad -4\frac{n-1}{n-2}\Delta_{g_c} u + R_{g_c} u = K(x)u^{\frac{n+2}{n-2}}.$$

The  $K(x) \equiv \text{const.}$  case of (6) on a general compact manifold is the famous Yamabe problem. (5) and (6) have attracted enormous attention in the last several decades. A large collection of phenomena on the possible behavior of solutions to these equations,

and methods and techniques of attacking these problems have been accumulated, which have tremendously enriched our understanding in solving a large class of nonlinear (elliptic) PDEs, and provided guidance in attacking seemingly unrelated problems. It is impossible in the space here to provide even a partial list of references. Please see [K85], [LP87], [S91], [L98], [CY02], [KMS09], and the references therein to get a glimpse of the results and techniques in this area.

Directly related to our current work are some work on the (potential) blow up analysis, a priori estimates, and existence of solutions to (5) or (6). It is proved in [H90] and [CGY93] that when  $K$  is a positive function on  $\mathbb{S}^2$ , a sequence of blowing up solutions to (5) has only one point blow up and has a well-defined blow up profile, and that when  $K$  is a positive  $C^2$  function on  $\mathbb{S}^2$  such that  $\Delta_{g_c} K(x) \neq 0$  at any of its critical points, no blow up can happen, more precisely, there is an a priori bound on the set of solutions to (5) which depends on the  $C^2$  norm of  $K$ , the positive lower bound of  $K$  and  $|\Delta_{g_c} K(x)|$  near the critical points of  $K$ , and the modulus of continuity of the second derivatives of  $K$ . Similar results for (6) in the case  $n = 3$  were proved in [Z90] and [CGY93], and for (6) in the case  $n \geq 4$  in [L95] under a flatness condition of  $K$  at its critical points. When  $K$  is a Morse function, say, this flatness condition fails when  $n \geq 4$ . In fact it is proved in [L96] that in such cases on  $\mathbb{S}^4$ , there can be a sequence of solutions to (6) blowing up at more than one points. Later on [CL99] constructed solutions blowing up on  $\mathbb{S}^n$ ,  $n \geq 7$ , with unbounded layers of “energy concentration” for certain non-degenerate  $K$ . A natural question concerning the  $\sigma_k$  curvature equations such as (3) is: which kind of behavior does its solution exhibit?

In the following we will often transform (3) through a conformal automorphism  $\varphi$  of  $\mathbb{S}^4$  as follows. Let  $|d\varphi(P)|$  denote the factor such that  $|d\varphi(P)[X]| = |d\varphi(P)||X|$  for any tangent vector  $X \in T_P(\mathbb{S}^4)$ , and

$$(7) \quad w_\varphi(P) = w \circ \varphi(P) + \ln |d\varphi(P)|.$$

Then  $w$  is a solution to (3) iff  $w_\varphi$  is a solution to

$$(8) \quad \sigma_2(g_c^{-1} \circ \left[ A_{g_c} - \nabla^2 w_\varphi + dw_\varphi \otimes dw_\varphi - \frac{1}{2} |\nabla w_\varphi|^2 g_c \right]) = K \circ \varphi(x) e^{4w_\varphi(x)}.$$

Our first results show that solutions to (3) on  $\mathbb{S}^4$  exhibits similar behavior as those to (5) on  $\mathbb{S}^2$  or (6) on  $\mathbb{S}^3$ .

**Theorem 1.** *Consider a family of admissible conformal metrics  $g_j = e^{2w_j} g_c$  on  $\mathbb{S}^4$  with*

$$\sigma_2(g_j^{-1} \circ A_{g_j}) = K(x), \text{ where } g_c \text{ denotes the canonical round metric on } \mathbb{S}^4 \text{ and } K(x)$$

*denotes a  $C^2$  positive function on  $\mathbb{S}^4$ . Then there exists at most one isolated simple blow up point in the sense that, if  $\max w_j = w_j(P_j) \rightarrow \infty$ , then there exists conformal*

automorphism  $\varphi_j$  of  $\mathbb{S}^4$  such that, if we define  $v_j(P) = w_j \circ \varphi_j(P) + \ln |d\varphi_j(P)|$ , we have

$$(9) \quad v_j(P) - \frac{1}{4} \ln \frac{6}{K(P_j)} \rightarrow 0 \quad \text{in } L^\infty(\mathbb{S}^4),$$

and

$$(10) \quad \int_{\mathbb{S}^4} |\nabla v_j|^4 \rightarrow 0.$$

In fact, we have the stronger conclusion that the  $W^{2,6}$  norm of  $v_j$  stays bounded and

$$v_j - \frac{1}{4} \ln \frac{6}{K(P_j)} \rightarrow 0 \quad \text{in } C^{1,\alpha}(\mathbb{S}^4) \quad \text{for any } 0 < \alpha < 1/3.$$

We also have

**Theorem 2.** *Let  $K(x)$  be a  $C^2$  positive function on  $\mathbb{S}^4$  satisfying a non-degeneracy condition*

$$(11) \quad \Delta K(P) \neq 0 \quad \text{whenever } \nabla K(P) = 0,$$

and we consider solutions  $w(x)$  to (3), with  $K(x)$  replaced by

$$K^{[s]}(x) := (1-s)6 + sK(x),$$

for  $0 < s \leq 1$ , namely,

$$(3') \quad \sigma_2(g_c^{-1} \circ \left[ A_{g_c} - \nabla^2 w + dw \otimes dw - \frac{1}{2} |\nabla w|^2 g_c \right]) = K^{[s]}(x) e^{4w(x)}.$$

Then there exist a priori  $C^{2,\alpha}$  estimates on  $w$ , uniform in  $0 < s \leq 1$ , which depend on the  $C^2$  norm of  $K$ , the modulus of continuity of  $\nabla^2 K$ , positive lower bound of  $\min K$  and positive lower bound of  $|\Delta K(x)|$  in a neighborhood of the critical points of  $K$ .

**Remark 1.** *Theorems 1 and 2, with the uniform estimates in Theorem 2 for solutions to (3') only for  $0 < s_0 \leq s \leq 1$  and the estimates possibly depending on  $0 < s_0 < 1$ , were obtained several years ago and were announced in [H04]. The details were written up in [CHY04] and presented by the second author on several occasions, including at the 2006 Banff workshop “Geometric and Nonlinear Analysis”. The current work can be considered as a completion of [CHY04]. As mentioned above similar statements for (5) and (6) were obtained earlier in [H90], [CGY93], [L95], [Z90], among others. When applying these estimates to the corresponding equation such as (5) and (6) with  $K(x)$*

replaced by  $K^{[s]}(x)$ , all previous work stated and proved that the a priori estimates on the solutions remain uniform as long as  $0 < s_0 \leq s \leq 1$ , for any fixed  $s_0$ . This stems from the dependence of the a priori estimates on a positive lower bound of  $|\Delta K(x)|$  near the

critical points of  $K$ , among other things. Since  $\Delta K^{[s]} = s\Delta K(x)$  becomes small when  $s > 0$  is small, previous work in this area assumed that the a priori estimates could deteriorate as  $s > 0$  becomes small. In these previous work, one has to devise a way to study the problem when  $s > 0$  becomes small. [CGY93] and [L95] used some kind of “center of mass” analysis via conformal transformations of the round sphere. Technically [CGY93] and [L95] used a constrained variational problem to study the “centered problem”. In essence the success of these methods was due to the semilinear nature of the relevant equations, so one could still have control on the “centered solution” in some norm weaker than  $C^{2,\alpha}$  norm, say,  $W^{2,p}$  norm, when  $s > 0$  is small, and used these estimates to prove existence of solutions under natural geometric/topological assumptions on  $K$ . This approach was problematic for our fully nonlinear equation (3). Due to this

difficulty, until recently we have not been successful in using our preliminary version of **Theorem 2** (for estimates in the range  $0 < s_0 \leq s \leq s$  which may depend on  $s_0$ ) and

the deformation  $K^{[s]}$  above to the equation to establish solutions to (3), under natural geometric/topological assumptions on  $K$ . It was our recent realization that in our setting, as well as in those of [CGY93] and [L95], the a priori estimates of solutions, under conditions like those in **Theorem 2**, remain uniform for all  $1 \geq s > 0$ ! After we completed this work and were compiling the bibliography for this paper, we noticed that M. Ji made a similar observation in her work on (5) in [J04]. This uniform a prior estimates for all  $1 \geq s > 0$  leads to our next Theorem.

**Theorem 3.** Suppose  $K(x)$  is a  $C^2$  positive function on  $\mathbb{S}^4$  satisfying (11). Then the map

$$G(P, t) = |\mathbb{S}^4|^{-1} \int_{\mathbb{S}^4} K \circ \varphi_{P,t}(x) x \, dvol_{g_c} \in \mathbb{R}^5$$

does not have a zero for  $(P, t) \in \mathbb{S}^4 \times [t_1, \infty)$ , for  $t_1$  large.

Furthermore, consider  $G$  as a map defined on  $(t-1)P/t \in B_r(O)$  for  $r > (t-1)/t$  and if

$$(12) \quad \deg(G, B_r(O), O) \neq 0, \quad \text{for } r \geq r_1 = (t-1)/t,$$

then (3) has a solution.

In particular, if  $K$  has only isolated critical points in the region  $\{x \in \mathbb{S}^4 : \Delta K(x) < 0\}$  and

$$\sum_{x \in \mathbb{S}^4 : \Delta K(x) < 0, \nabla K(x) = 0} \text{ind}(\nabla K(x)) \neq 1,$$

where  $\text{ind}(\nabla K(x))$  stands for the index of the vector field  $\nabla K(x)$  at its isolated zero  $x$ , then (12) holds, therefore, (3) has a solution.

A corollary of the proof for the  $W^{2,p}$  estimate in Theorem 1 is a bound on a functional determinant whose critical points are solutions to (3). We recall that the relevant functional determinant is defined, similar to [CGY02a], through

$$II[w] = \int_{\mathbb{S}^4} ((\Delta_0 w)^2 + 2|\nabla_0 w|^2 + 12w) \, d\text{vol}_{g_c},$$

$$C_K[w] = 3 \log \left( \int_{\mathbb{S}^4} K e^{4w} \, d\text{vol}_{g_c} \right),$$

and

$$\begin{aligned} Y[w] &= \frac{1}{36} \left( \int_{\mathbb{S}^4} R^2 \, d\text{vol}_{g_c} - \int_{\mathbb{S}^4} R_0^2 \, d\text{vol}_{g_c} \right) \\ &= \int_{\mathbb{S}^4} (\Delta_0 w + |\nabla_0 w|^2)^2 \, d\text{vol}_{g_c} - 4 \int_{\mathbb{S}^4} |\nabla_0 w|^2 \, d\text{vol}_{g_c}. \end{aligned}$$

$F[w] = Y[w] - II[w] + C_K[w]$  is the relevant functional determinant and a critical point of  $F[w]$  is a solution of (3). It is known that, for any conformal transformation  $\varphi$  of  $(\mathbb{S}^4, g_c)$ ,  $Y[w_\varphi] = Y[w]$ , and  $II[w_\varphi] = II[w]$ .

There is a similar functional determinant and a variational characterization for solutions to the prescribing Gaussian curvature problem on  $\mathbb{S}^2$ . Chang, Gursky and Yang proved in [CGY93] that this functional is bounded on the set of solutions to the prescribing Gaussian curvature problem on  $\mathbb{S}^2$  for any positive function  $K$  on  $\mathbb{S}^2$  to be prescribed. Our corollary is in the same spirit.

**Corollary 1.** *Let  $K(x)$  be a given positive  $C^2$  function on  $\mathbb{S}^4$ . Then there is a bound  $C$  depending on  $K$  only through the  $C^2$  norm of  $K$ , a positive upper and lower bound of  $K$  on  $\mathbb{S}^4$ , such that*

$$|F[w]| \leq C$$

for all admissible solutions  $w$  to (3).

**Theorem 1** will be established using blow up analysis, Liouville type classification results of entire solutions, and integral type estimates for such fully nonlinear equations from [CGY02a] and [H04]. **Theorem 2** will be established using a weaker version of

**Theorem 1** and a Kazdan-Warner type identity satisfied by the solutions. The weaker version of **Theorem 1** only needs to establish

$$(9') \quad v_j(P) - \frac{1}{4} \frac{6}{K(P_j)} \rightarrow 0 \quad \text{pointwise on } \mathbb{S}^4 \setminus \{-P_j\}, \text{ and bounded in } L^\infty(\mathbb{S}^4),$$

instead of (9), (10) and the  $W^{2,6}$  norm estimates. A degree argument for a fully nonlinear operator associated with (3) and **Theorem 2** will be used to establish **Theorem 3**. To streamline our presentation, we will first outline the main steps for proving **Theorem 3**, assuming **Theorem 2** and all the other needed ingredients. In the remaining sections, we will first provide a proof for (9') in **Theorem 1** and for **Theorem 2**, before finally providing a proof for the  $W^{2,6}$  norm estimates in **Theorem 1** and for **Corollary 1**.

## 2. PROOF OF THEOREM 3

The first and third parts of **Theorem 3** is contained in [CY91] and [CGY93]. We will establish the second part of **Theorem 3** by formulating the existence of a solution to (3) as a degree problem for a nonlinear map and linking the degree of this map to that of  $G$ .

By a fibration result from [CY87], [CY91], [CL93] and [L95], see also [A79], [O82] for early genesis of these ideas, if we define

$$\mathcal{S}_0 = \left\{ v \in C^{2,\alpha}(\mathbb{S}^4) : \int_{\mathbb{S}^4} e^{4v(x)} x \, d\text{vol}_{g_c} = 0 \right\},$$

then the map  $\pi : (v, \xi) \in \mathcal{S}_0 \times B \mapsto C^{2,\alpha}(\mathbb{S}^4)$  defined by

$$\pi(v, \xi) = v \circ \varphi_{P,t}^{-1} + \ln |d\varphi_{P,t}^{-1}|,$$

with  $B$  denoting the open unit ball in  $\mathbb{R}^5$  and  $\xi = rP$ ,  $P \in \mathbb{S}^4$ ,  $r = (t-1)/t$ ,  $t \geq 1$ , is a  $C^2$  diffeomorphism from  $\mathcal{S}_0 \times B$  onto  $C^{2,\alpha}(\mathbb{S}^4)$ . Thus  $(v, P, t) \in \mathcal{S}_0 \times \mathbb{S}^4 \times [1, \infty)$  provide global coordinates for  $C^{2,\alpha}(\mathbb{S}^4)$  (with a coordinate singularity at  $t = 1$ , similar to the coordinate singularity of polar coordinates at  $r = 0$ ) through

$$w = v \circ \varphi_{P,t}^{-1} + \ln |d\varphi_{P,t}^{-1}|.$$

$w$  solves (3) with  $K$  replaced by  $K^{[s]}$  iff  $v$  solves

$$(13) \quad \sigma_2(A_v) = K^{[s]} \circ \varphi_{P,t} e^{4v}.$$

Then the estimates for  $w$  in **Theorem 2** turn into the following estimates for  $v$  and  $t$ .

**Proposition 1.** *Assume that  $K$  is a positive  $C^2$  function on  $\mathbb{S}^4$  satisfying the non-degeneracy condition (11), and let  $w$  be a solution to (3) with  $K$  replaced by  $K^{[s]}$  and  $(v, P, t) \in \mathcal{S}_0 \times \mathbb{S}^4 \times [1, \infty)$  be the coordinates of  $w$  defined in the paragraph above. Then there exist  $t_0$  and  $\epsilon(s) > 0$  with  $\lim_{s \rightarrow 0} \epsilon(s) = 0$ , such that*

$$(14) \quad t \leq t_0 \quad \text{and} \quad \|v\|_{C^{2,\alpha}(\mathbb{S}^4)} < \epsilon(s).$$

A proof for **Proposition 1** will be postponed to the end of the next section.

We treat (13) as a nonlinear map

$$F^{[s]}[v, \xi] := e^{-4v(x)} \sigma_2(g_c^{-1} \circ \left[ A_{g_c} - \nabla^2 v + dv \otimes dv - \frac{1}{2} |\nabla v|^2 g_c \right]) - K^{[s]} \circ \varphi_{P,t},$$

from  $\mathcal{S}_0 \times B$  into  $C^\alpha(\mathbb{S}^4)$ , for  $0 < s \leq 1$ , where  $\xi = (t-1)P/t \in B$ . **Proposition 1** implies that there is a neighborhood  $\mathcal{N} \subset \mathcal{S}_0$  of  $0 \in \mathcal{S}_0$  and  $0 < r_0 = (t_0 - 1)/t_0 < 1$  such

that  $F^{[s]}$  does not have a zero on  $\partial(\mathcal{N} \times B_r)$  for all  $r_0 \leq r < 1$  and  $0 < s \leq 1$ . According

to [L89], there is a well defined degree for  $F^{[s]}$  on  $\mathcal{N} \times B_{r_0}$  and it is independent of

$0 < s \leq 1$ . We will compute this degree of  $F^{[s]}$ , for  $s > 0$  small, through the degree of a finite dimensional map.

We first use the implicit function theorem to define this map and link the solutions to (13) to the zeros of this map. Note that  $F^{[0]}[0, (t-1)P/t] = 0$  and  $D_v F^{[0]}[0, (t-1)P/t](\eta) = -6\Delta\eta - 24\eta$ . If  $\Pi$  denotes the projection from  $C^\alpha(\mathbb{S}^4)$  into

$$Y := \{f \in C^\alpha(\mathbb{S}^4) : \int_{\mathbb{S}^4} f x_j dvol_{g_c} = 0 \quad \text{for } j = 1, \dots, 5\}$$

defined by  $\Pi(f) = f - 5|\mathbb{S}^4|^{-1} \sum_{j=1}^5 \left( \int_{\mathbb{S}^4} f x_j dvol_{g_c} \right) x_j$ , then we can apply the implicit

function theorem to  $\Pi \circ F^{[s]}$  at  $v = 0$  to conclude

**Proposition 2.** *There exist some neighborhood  $\mathcal{N}_\epsilon \subset \mathcal{N}$  of  $0 \in \mathcal{S}_0$  and  $s_0 > 0$ , such that for all  $0 < s < s_0$ ,  $(P, t) \in \mathbb{S}^4 \times [1, t_0]$ , there exists a unique  $v = v(x; P, t, s) \in \mathcal{N}_\epsilon$ ,*



depending differentiably on  $(P, t, s)$  such that

$$(15) \quad \Pi \circ F^{[s]}[v(x; P, t, s), (t-1)P/t] = 0.$$

Furthermore, there exists some  $C > 0$  such that, for  $0 < s \leq s_0$ ,  $1 \leq t \leq t_0$ ,

$$(16) \quad \|v(x; P, t, s)\|_{C^{2,\alpha}(\mathbb{S}^4)} \leq C \|K^{[s]} \circ \varphi_{P,t} - 6\|_{C^\alpha(\mathbb{S}^4)} = Cs \|K \circ \varphi_{P,t} - 6\|_{C^\alpha(\mathbb{S}^4)}.$$

(15) implies that

$$F^{[s]}[v(x; P, t, s), (t-1)P/t] = \sum_{j=1}^5 \Lambda_j(P, t, s)x_j,$$

for some Lagrange multipliers  $\Lambda_j(P, t, s)$ , which depend differentiably on  $(P, t, s)$ . Or, equivalently,

$$(17) \quad \sigma_2(A_{v(x;P,t,s)}) = \left( K^{[s]} \circ \varphi_{P,t} + \sum_{j=1}^5 \Lambda_j(P, t, s)x_j \right) e^{4v(x;P,t,s)}.$$

A zero of the map  $\Lambda^{[s]}(P, t) := (\Lambda_1(P, t, s), \dots, \Lambda_5(P, t, s))$  corresponds to a solution to (13). **Propositions 1** and **2** say that, for  $s_0 > 0$  small, all solutions  $v \in \mathcal{S}_0$  to (13), for

$0 < s \leq s_0$ , are in  $\mathcal{N}_\epsilon$ , thus correspond to the zeros of the map  $\Lambda^{[s]}(P, t)$ .

**Remark 2.**  $\|K \circ \varphi_{P,t} - 6\|_{C^\alpha(\mathbb{S}^4)}$  could become unbounded when  $t \rightarrow \infty$ ; yet thanks to

the bound  $1 \leq t \leq t_0$  from **Proposition 1**, it remains bounded in terms of  $\|K\|_{C^\alpha(\mathbb{S}^4)}$  in

the range  $1 \leq t \leq t_0$ . Note also that  $\|K \circ \varphi_{P,t} - 6\|_{L^p(\mathbb{S}^4)}$  remains bounded in terms of

$\|K\|_{L^\infty(\mathbb{S}^4)}$  even in the range  $1 \leq t < \infty$ . It is essentially this bound and the applicability

of  $W^{2,p}$  estimates in the semilinear setting of [CGY93] and [L95] which allowed them to handle their cases without using the bound  $1 \leq t \leq t_0$ .

**Remark 3.** The implicit function theorem procedure here works also in the setting of [CY91], [CGY93] and [L95] using  $W^{2,p}$  space, as does Proposition 1 in the setting of [CGY93] and [L95], and can be used to simplify the arguments there.

At this point, we need the following Kazdan-Warner type identity for solutions to (3).

**Proposition 3.** *Let  $w$  be a solution to (3). Then, for  $1 \leq j \leq 5$ ,*

$$(18) \quad \int_{\mathbb{S}^4} \langle \nabla K(x), \nabla x_j \rangle e^{4w(x)} dvol_{g_c} = 0.$$

**Proposition 3** is a special case of the results in [V00b] and [H06]. But in the special case of  $\mathbb{S}^4$ , it is a direct consequence of the variational characterization of the solution to (3), as given after the statement of Theorem 1. A solution  $w$  to (3) is a critical point of  $F[w] = Y[w] - II[w] + C_K[w]$  there, thus satisfies, for any one-parameter family of conformal diffeomorphisms  $\varphi_s$  of  $\mathbb{S}^4$  with  $\phi_0 = \text{Id}$ ,

$$\left. \frac{d}{ds} \right|_{s=0} F[w_{\varphi_s}] = 0,$$

with  $w_{\varphi_s} = w \circ \varphi_s + \log |d\varphi_s|$ . Since  $Y[w_{\varphi_s}] = Y[w]$  and  $II[w_{\varphi_s}] = II[w]$ , a solution  $w$  of (3) thus satisfies

$$\left. \frac{d}{ds} \right|_{s=0} C_K[w_{\varphi_s}] = \left. \frac{d}{ds} \right|_{s=0} \left( \int_{\mathbb{S}^4} K \circ \varphi_s^{-1} e^{4w} dvol_{g_c} \right) = 0,$$

which is (18).

Applying (18) to  $v(x; P, t, s)$ , a solution to (17), we obtain,

$$\int_{\mathbb{S}^4} \langle \nabla \left( K^{[s]} \circ \varphi_{P,t}(x) + \sum_{j=1}^5 \Lambda_j(P, t, s) x_j \right), \nabla x_k \rangle e^{4v(x; P, t, s)} dvol_{g_c} = 0, \quad \text{for } 1 \leq k \leq 5,$$

from which we obtain, for  $1 \leq k \leq 5$ ,

$$\begin{aligned} & - \sum_{j=1}^5 \Lambda_j(P, t, s) \int_{\mathbb{S}^4} \langle \nabla x_j, \nabla x_k \rangle e^{4v(x; P, t, s)} dvol_{g_c} \\ &= \int_{\mathbb{S}^4} \langle \nabla (K^{[s]} \circ \varphi_{P, t}(x)), \nabla x_k \rangle e^{4v(x; P, t, s)} dvol_{g_c} \\ &= s \int_{\mathbb{S}^4} \langle \nabla (K \circ \varphi_{P, t}(x)), \nabla x_k \rangle e^{4v(x; P, t, s)} dvol_{g_c}. \end{aligned}$$

As in [CY91], [CGY93] and [L95], we define

$$A^{[s]}(P, t) = (4|\mathbb{S}^4|)^{-1} \int_{\mathbb{S}^4} \langle \nabla (K \circ \varphi_{P, t}(x)), \nabla x \rangle e^{4v(x; P, t, s)} dvol_{g_c} \in \mathbb{R}^5.$$

Since

$$\left( \int_{\mathbb{S}^4} \langle \nabla x_j, \nabla x_k \rangle e^{4v(x; P, t, s)} dvol_{g_c} \right)$$

is positive definite, we conclude that

$$\deg(A^{[s]}, B_{r_0}, O) = -\deg(\Lambda^{[s]}, B_{r_0}, O),$$

for  $s_0 > s > 0$  provided that one of them is well defined.

Using  $v(x; P, t, s) \in \mathcal{S}_0$ , and  $\Delta x = -4x$  on  $\mathbb{S}^4$ , we have, as in [CY91], [CGY93] and [L95],

$$A^{[s]}(P, t) = G(P, t) + I + \Pi,$$

where

$$I = |\mathbb{S}^4|^{-1} \int_{\mathbb{S}^4} (K \circ \varphi_{P, t}(x) - K(P)) x (e^{4v(x; P, t, s)} - 1) dvol_{g_c},$$

and

$$\Pi = -(4|\mathbb{S}^4|)^{-1} \int_{\mathbb{S}^4} (K \circ \varphi_{P, t}(x) - K(P)) \langle \nabla x, \nabla e^{4v(x; P, t, s)} \rangle dvol_{g_c}.$$

We could have fixed  $t_0 \geq t_1$  such that  $G(P, t) \neq 0$  for  $t = t_0$ , and there will be a  $\delta > 0$  such that  $|G(P, t)| \geq \delta$  for  $t = t_0$ . Since (16) implies that

$$\|v(x; P, t, s)\|_{C^{2,\alpha}(\mathbb{S}^4)} = O(s), \quad \text{uniformly for } (P, t) \in \mathbb{S}^4 \times [1, t_0],$$

we find that, by fixing  $s_0 > 0$  small if necessary,

$$|I| + |\Pi| \leq \frac{1}{2}|G(P, t)|, \quad \text{for } 0 < s \leq s_0 \text{ and } t = t_0.$$

This implies that  $A^{[s]}(P, t) \cdot G(P, t) > 0$  for  $0 < s \leq s_0$  and  $t = t_0$ . Therefore

$$-\deg(\Lambda^{[s]}, B_{r_0}, O) = \deg(A^{[s]}, B_{r_0}, O) = \deg(G, B_{r_0}, O) \neq 0,$$

for  $0 < s \leq s_0$ . Finally, we now prove

$$(19) \quad \deg(F^{[s]}, \mathcal{N} \times B_{r_0}, O) = -\deg(\Lambda^{[s]}, B_{r_0}, O),$$

for  $0 < s \leq s_0$ , from which follows the existence of a solution to (3).

The verification of (19) is routine, but requires several steps. First, we may perturb  $K$ , if necessary, within the class of functions satisfying the conditions in **Theorem 3** such that the corresponding  $G(P, t)$  has only isolated and non-degenerate zeros in  $B_{r_0}(O)$ .

We will prove momentarily that for  $s > 0$  small, the zeros of  $\Lambda^{[s]}$  for  $s > 0$  small will be close to the zeros of  $G(P, t)$  and are isolated, non-degenerate. Therefore the zeros of

$F^{[s]}$  in  $\mathcal{N} \times B_{r_0}$  are isolated and non-degenerate. This can be argued as follows. First, it follows from (17) that

$$\Lambda^{[s]}(\xi) \cdot x = (\text{Id} - \Pi) \left( e^{-4v(x; \xi, s)} \sigma_2(A_{v(x; \xi, s)}) - K^{[s]} \circ \varphi_{P, t} \right).$$

Using (16), we can then write

$$e^{-4v(x; \xi, s)} \sigma_2(A_{v(x; \xi, s)}) = 6 - 6\Delta v(x; \xi, s) - 24v(x; \xi, s) + Q(v(x; \xi, s)),$$

with  $\|Q(v(x; \xi, s))\|_Y \lesssim \|v(x; \xi, s)\|_X^2 \lesssim s^2$ . Therefore, using  $(\text{Id} - \Pi)(1) = (\text{Id} - \Pi)(6\Delta v(x; \xi, s) + 24v(x; \xi, s)) = 0$ , and

$$\Lambda^{[s]}(\xi) = 5|\mathbb{S}^4|^{-1} \int_{\mathbb{S}^4} (\Lambda^{[s]}(\xi) \cdot x) x \, d\text{vol}_{g_c},$$

we have

$$\Lambda^{[s]}(\xi) = -5sG(P, t) + 5|\mathbb{S}^4|^{-1} \int_{\mathbb{S}^4} [(\text{Id} - \Pi)(Q(v(x; \xi, s)))] x \, d\text{vol}_{g_c},$$

with  $|(\text{Id} - \Pi)(Q(v(x; \xi, s)))| \lesssim s^2$ , so the zeros of  $\Lambda^{[s]}(\xi)$  for  $s > 0$  small are close to the zeros of  $G(P, t)$ . We can further use the implicit function theorem to prove that for

$s > 0$  small there is a (unique) non-degenerate zero of  $\Lambda^{[s]}(\xi)$  near each zero of  $G(P, t)$ .

**Remark 4.** *This argument shows that, for each non-degenerate zero of  $G(P, t)$ , if we associate  $(P, t)$  with the center of mass of  $\varphi_{P, t}$ ,*

$$C.M(\varphi_{P, t}) := |\mathbb{S}^4|^{-1} \int_{\mathbb{S}^4} \varphi_{P, t}(x) \, d\text{vol}_{g_c} \in B_1(O)$$

as a geometric representation of  $(P, t)$ , then for  $s > 0$  small, there is a unique solution  $w$  to (3) whose center of mass approaches  $C.M(\varphi_{P, t})$ , for our argument gives rise to a solution

$$w(x) = v(\cdot; P', t') \circ \varphi_{P', t'}^{-1}(x) + \ln |d\varphi_{P', t'}^{-1}(x)|$$

with  $(P', t')$  approaching  $(P, t)$ , and  $v(x; P', t')$  approaching 0 as  $s \rightarrow 0$ , thus the center of mass of  $w$  is

$$|\mathbb{S}^4|^{-1} \int_{\mathbb{S}^4} e^{4w(x)} x \, d\text{vol}_{g_c} = |\mathbb{S}^4|^{-1} \int_{\mathbb{S}^4} e^{4v(y)} \varphi_{P', t'}(y) \, d\text{vol}_{g_c} \rightarrow C.M(\varphi_{P, t})$$

as  $s \rightarrow 0$ .

Now  $\deg(F^{[s]}, \mathcal{N} \times B_{r_0}, O)$  is well defined in the manner of [L89], and according to Propositions 2.1–2.4 of [L89],

$$\deg(F^{[s]}, \mathcal{N} \times B_{r_0}, O) = \sum_{\xi \in B_{r_0}(O): \Lambda^{[s]}(\xi)=0} \text{ind}(DF^{[s]}[v(x; \xi, s), \xi]),$$

where  $\text{ind}(DF^{[s]}[v(x; \xi, s), \xi])$  refers to the index of the linear operator  $DF^{[s]}[v(x; \xi, s), \xi]$ , and is computed as  $(-1)^\beta$ , with  $\beta$  denoting the number of negative eigenvalues of

$DF^{[s]}[v(x; \xi, s), \xi]$ . We also have

$$\deg(\Lambda^{[s]}, B_{r_0}, O) = \sum_{\xi \in B_{r_0}(O): \Lambda^{[s]}(\xi)=0} \text{ind}(D\Lambda^{[s]}(\xi)).$$

To compute  $DF^{[s]}[v(x; \xi, s), \xi]$ , we identify  $\xi \in \mathbb{R}^5$  with  $\xi \cdot x \in \text{span}\{x_1, \dots, x_5\}$ , and write the differential of  $F^{[s]}$  in the direction of  $\dot{v}$  as  $D_v F^{[s]}[v(x; \xi, s), \xi](\dot{v})$ , or simply  $D_v F^{[s]}(\dot{v})$ , and the differential of  $F^{[s]}$  in the direction of  $\dot{\xi} \cdot x$  as  $D_\xi F^{[s]}[v(x; \xi, s), \xi](\dot{\xi})$ . Then

$$D_\xi F^{[s]}[v(x; \xi, s), \xi](\dot{\xi}) = -s\dot{\xi} \cdot \nabla_\xi (K \circ \varphi_{P,t}),$$

and

$$D_v F^{[s]}[v(x; \xi, s), \xi](\dot{v}) = M^{ij}[v(x; \xi, s)] \nabla_{ij}^{v(x; \xi, s)} \dot{v} - 4K^{[s]} \circ \varphi_{P,t} \dot{v},$$

where  $M^{ij}[v(x; \xi, s)]$  stands for the Newton tensor associated with  $\sigma_2(e^{-2v(x; \xi, s)} A_{v(x; \xi, s)})$ ,

and  $\nabla_{ij}^{v(x; \xi, s)}$  stands for the covariant differentiation in the metric  $e^{2v(x; \xi, s)} g_c$ . Thus,

$$DF^{[s]}[v(x; \xi, s), \xi](\dot{v} + \dot{\xi} \cdot x) = M^{ij}[v(x; \xi, s)] \nabla_{ij}^{v(x; \xi, s)} \dot{v} - 4K^{[s]} \circ \varphi_{P,t} \dot{v} - s\dot{\xi} \cdot \nabla_\xi (K \circ \varphi_{P,t}).$$

At a fixed zero  $\xi$  of  $\Lambda^{[s]}(\xi) = 0$ , we define a family of deformed linear operators  $L_{\tau, s}$  for  $0 \leq \tau \leq 1$  by

$$L_{\tau, s}(\dot{v} + \dot{\xi} \cdot x) = M^{ij}[v^{[\tau]}] \nabla_{ij}^{v^{[\tau]}} \dot{v} - 4K^{[s\tau]} \circ \varphi_{P,t} \dot{v} - s\dot{\xi} \cdot \nabla_\xi (K \circ \varphi_{P,t}),$$

where  $v^{[\tau]} = \tau v(x; \xi, s)$ , and  $\dot{v} \in X := \{\dot{v} \in C^{2,\alpha}(\mathbb{S}^4) : \int_{\mathbb{S}^4} \dot{v}(x) x_j = 0, j = 1, \dots, 5\}$ .

Then  $L_{\tau,s}$  defines self-adjoint operators with respect to the metric  $e^{2v^{[\tau]}} g_c$ , thus its eigenvalues are all real. We first assume the

**Claim.** *For  $s > 0$  small and  $0 \leq \tau \leq 1$ , the spectrum of  $L_{\tau,s}$  does not contain zero.*

Thus  $\text{ind}(L_{0,s}) = \text{ind}(L_{1,s}) = \text{ind}(DF^{[s]}[v(x; \xi, s), \xi])$ . We will next establish

$$(20) \quad \text{ind}(L_{0,s}) = (-1)^{1+\gamma}, \quad \text{for } s > 0 \text{ small,}$$

where  $\gamma$  is the number of positive eigenvalues of  $\nabla G(P, t)$  at  $\xi$ , and

$$(21) \quad \gamma = \text{the number of negative eigenvalues of } D_\xi \Lambda^{[s]}(\xi) \text{ for } s > 0 \text{ small.}$$

First note that

$$L_{0,s}(\dot{v} + \dot{\xi} \cdot x) = -6\Delta \dot{v} - 24\dot{v} - s\dot{\xi} \cdot \nabla_\xi (K \circ \varphi_{P,t}),$$

so if  $\dot{v} + \dot{\xi} \cdot x$  is an eigenfunction corresponding to a negative eigenvalue  $-\lambda$ , with  $\dot{v} \in X$ , then

$$-6\Delta \dot{v} - 24\dot{v} - s\dot{\xi} \cdot \nabla_\xi (K \circ \varphi_{P,t}) = -\lambda(\dot{v} + \dot{\xi} \cdot x).$$

Taking projection in  $\text{span}\{x_1, \dots, x_5\}$ , we find

$$-s\nabla G(P, t)\dot{\xi} = -\frac{\lambda}{5}\dot{\xi},$$

and taking projection in  $X$ , we find

$$(22) \quad -6\Delta \dot{v} - 24\dot{v} - s\Pi\left(\dot{\xi} \cdot \nabla_\xi (K \circ \varphi_{P,t})\right) = -\lambda\dot{v}.$$

If  $\dot{\xi} \neq 0$ , then  $\dot{\xi}$  is an eigenvector of  $\nabla G(P, t)$  with eigenvalue  $\frac{\lambda}{5s} > 0$ ; and if  $\dot{\xi} = 0$ , then  $\dot{v} \neq 0$  solves  $-6\Delta \dot{v} - 24\dot{v} = -\lambda\dot{v}$ , which is possible for some  $\lambda > 0$  iff  $-\lambda = -24$  and

$\dot{v} = \text{constant}$ . Conversely, for any eigenvector  $\dot{\xi} \neq 0$  of  $\nabla G(P, t)$  with eigenvalue  $\mu > 0$ , the operator  $-6\Delta - 24 + 5s\mu$  is an isomorphism from  $X$  to  $Y$  for  $s > 0$  small, so we

can solve (22) as  $(-6\Delta - 24 + 5s\mu)\dot{v} = s\Pi\left(\dot{\xi} \cdot \nabla_{\xi}(K \circ \varphi_{P,t})\right)$  for  $\dot{v} \in X$  and  $\dot{v} + \dot{\xi} \cdot x$  becomes an eigenfunction of  $L_{0,s}$  with eigenvalue  $-5s\mu$ . Therefore we conclude (20).

We now establish (21) to prove  $\text{ind}(DF^{[s]}[v(x; \xi, s), \xi]) = -\text{ind}(D_{\xi}\Lambda^{[s]}(\xi))$  for  $s > 0$  small. From (17), which can be written as  $\Lambda^{[s]}(\xi) \cdot x = F^{[s]}[v(x; \xi, s), \xi]$ , we obtain

$$D_v F^{[s]}(D_{\xi}v(x; \xi, s)(\dot{\xi})) + D_{\xi} F^{[s]}(\dot{\xi}) = D_{\xi}\Lambda^{[s]}(\xi)(\dot{\xi}) \cdot x.$$

Taking projections in  $X$  and  $\text{span}\{x_1, \dots, x_5\}$ , respectively, and using  $D_{\xi}F^{[s]}(\dot{\xi}) = -s\dot{\xi} \cdot \nabla_{\xi}(K \circ \varphi_{P,t})$ , we obtain

$$(23) \quad \Pi\left(D_v F^{[s]}(D_{\xi}v(x; \xi, s)(\dot{\xi}))\right) - s\Pi\left(\dot{\xi} \cdot \nabla_{\xi}(K \circ \varphi_{P,t})\right) = 0,$$

and

$$(24) \quad \int_{\mathbb{S}^4} (\text{Id} - \Pi)\left(D_v F^{[s]}(D_{\xi}v(x; \xi, s)(\dot{\xi}))\right) x \, d\text{vol}_{g_c} - s\nabla_{\xi}G(P, t)\dot{\xi} = \frac{1}{5}D_{\xi}\Lambda^{[s]}(\xi)(\dot{\xi}).$$

Writing

$$D_v F^{[s]}(D_{\xi}v(x; \xi, s)(\dot{\xi})) = (-6\Delta - 24)(D_{\xi}v(x; \xi, s)(\dot{\xi})) + \Theta(D_{\xi}v(x; \xi, s)(\dot{\xi})),$$

we find, using (16), that  $\|\Theta(D_{\xi}v(x; \xi, s)(\dot{\xi}))\|_Y \lesssim s\|D_{\xi}v(x; \xi, s)(\dot{\xi})\|_X$ . Thus  $\Pi(D_v F^{[s]}(\cdot)) : X \mapsto Y$  is an isomorphism for  $s > 0$  small and has an inverse  $\Psi$ , and we can solve

$D_{\xi}v(x; \xi, s)(\dot{\xi})$  in terms of  $\dot{\xi}$  from (23):

$$D_{\xi}v(x; \xi, s)(\dot{\xi}) = \Psi\left(s\Pi\left(\dot{\xi} \cdot \nabla_{\xi}(K \circ \varphi_{P,t})\right)\right) := s\Upsilon(\dot{\xi}).$$



Using this in (24), we find

$$\begin{aligned} \frac{1}{5} D_\xi \Lambda^{[s]}(\xi) &= -s \nabla_\xi G(P, t) + s \int_{\mathbb{S}^4} [(\text{Id} - \Pi) \circ \Theta \circ \Upsilon] x \, d\text{vol}_{g_c} \\ &= -s (\nabla_\xi G(P, t) + O(s)). \end{aligned}$$

Thus for  $s > 0$  small,  $\gamma$  matches the number of negative eigenvalues of  $D_\xi \Lambda^{[s]}(\xi)$ , and we can conclude that  $\text{ind}(DF^{[s]}(v(x; \xi, s), \xi)) = -\text{ind}(D_\xi \Lambda^{[s]}(\xi))$ .

In the remainder of this section, we provide proof for our **Claim** above, leaving the proof for **Proposition 1** to the end of the next section.

*Proof of Claim.* Suppose that for (a sequence of)  $s > 0$  small and some  $0 \leq \tau \leq 1$ ,  $L_{\tau, s}$

has  $\dot{v} + \dot{\xi} \cdot x$ , with  $\dot{v} \in X$ ,  $\dot{\xi} \in \mathbb{R}^5$ , as eigenfunction with zero eigenvalue. Then, taking projections in  $\text{span}\{x_1, \dots, x_5\}$  and  $X$ , respectively, we obtain

$$(25) \quad \Pi \left[ M^{ij}[v^{[\tau]}] \nabla_{ij}^{v^{[\tau]}} \dot{v} - 4K^{[s\tau]} \circ \varphi_{P,t} \dot{v} \right] - s \Pi \left[ \dot{\xi} \cdot \nabla_\xi (K \circ \varphi_{P,t}) \right] = 0,$$

and

$$(26) \quad \int_{\mathbb{S}^4} (\text{Id} - \Pi) \left[ M^{ij}[v^{[\tau]}] \nabla_{ij}^{v^{[\tau]}} \dot{v} - 4K^{[s\tau]} \circ \varphi_{P,t} \dot{v} \right] x \, d\text{vol}_{g_c} - s \nabla_\xi G(P, t) \dot{\xi} = 0.$$

Using (16) again, we find

$$M^{ij}[v^{[\tau]}] \nabla_{ij}^{v^{[\tau]}} \dot{v} - 4K^{[s\tau]} \circ \varphi_{P,t} \dot{v} = -6\Delta \dot{v} - 24\dot{v} + \Theta^{\tau, s}(\dot{v}),$$

with  $\|\Theta^{\tau, s}(\dot{v})\|_Y \lesssim s\tau \|\dot{v}\|_X$ . Thus, for  $s > 0$  small, we can solve  $\dot{v}$  from (25) to obtain

$$\dot{v} = \Psi \left( s \Pi \left[ \dot{\xi} \cdot \nabla_\xi (K \circ \varphi_{P,t}) \right] \right) = s \Upsilon(\dot{\xi}).$$

Thus  $\dot{\xi} \neq 0$  and we can normalize it so that  $|\dot{\xi}| = 1$ . Using this in (26), we find

$$s(\text{Id} - \Pi) \circ \Theta^{\tau, s} \circ \Upsilon(\dot{\xi}) - s\nabla_{\xi} G(P, t)\dot{\xi} = 0.$$

Using  $\|\Theta^{\tau, s}\| \lesssim s\tau$ , we find this impossible for  $s > 0$  small under our non-degeneracy assumption on the zeros of  $G(P, t)$ .  $\square$

### 3. PROOF OF (9'), (10), THEOREM 2 AND PROPOSITION 1

*Proof of (9') and (10).* The full strength of (9) is established as soon as the  $W^{2,3}$  estimates are established — the latter is a step in proving the  $W^{2,6}$  estimates.

If there is a sequence of solutions  $w_j$  to (3) such that  $\max w_j = w_j(P_j) \rightarrow \infty$ , then we choose conformal automorphism  $\phi_j = \phi_{P_j, t_j}$  of  $\mathbb{S}^4$ , such that the rescaled function

$$(27) \quad v_j(P) = w_j \circ \phi_j(P) + \ln |d\phi_j(P)|,$$

satisfies the normalization condition

$$(28) \quad v_j(P_j) = \frac{1}{4} \ln \frac{6}{K(P_j)}.$$

If we use stereographic coordinates for  $\mathbb{S}^4$ , with  $P_j$  as the north pole, then

$$y(\phi_j(P)) = t_j y(P), \text{ for } P \in \mathbb{S}^4,$$

and

$$v_j(P) = w_j \circ \phi_j(P) + \ln \frac{t_j(1 + |y(P)|^2)}{1 + t_j^2|y(P)|^2}.$$

$v_j$  would satisfy

$$(29) \quad \sigma_2(A_{v_j}) = K \circ \phi_j e^{4v_j}.$$

The normalization in (28) amounts to choosing  $t_j$  such that

$$w_j(P_j) - \ln t_j = \frac{1}{4} \ln \frac{6}{K(P_j)}.$$

Thus,  $t_j \rightarrow \infty$ , and for any  $P \in \mathbb{S}^4$ ,

$$(30) \quad v_j(P) \leq \frac{1}{4} \ln \frac{6}{K(P_j)} + \ln \frac{t_j^2 (1 + |y(P)|^2)}{1 + t_j^2 |y(P)|^2}.$$

(30) implies that, away from  $-P_j$ ,  $v_j$  has an upper bound independent of  $j$ . Together with (29), the local gradient and higher derivative estimates of [GW03], there exists a subsequence, still denoted as  $\{v_j\}$ , such that,  $P_j \rightarrow P_*$ , and for any  $\delta > 0$ ,

$$(31) \quad v_j \rightarrow v_\infty \text{ in } C^{2,\alpha}(\mathbb{S}^4 \setminus B_\delta(-P_*)), \quad \text{for some limit } v_\infty.$$

We also have

$$(32) \quad \sigma_2(A_{v_\infty}) = K(P_*)e^{4v_\infty} \quad \text{on } \mathbb{S}^4 \setminus \{-P_*\},$$

$$(33) \quad \int_{\mathbb{S}^4} K(P_*)e^{4v_\infty} d\text{vol}_{g_c} \leq \liminf_{j \rightarrow \infty} \int_{\mathbb{S}^4} K \circ \phi_j e^{4v_j} d\text{vol}_{g_c} = 16\pi^2,$$

$$(34) \quad v_\infty(P_*) = \frac{1}{4} \ln \frac{6}{K(P_*)}, \quad \nabla v_\infty(P_*) = 0,$$

$$(35) \quad v_\infty(P) \leq \frac{1}{4} \ln \frac{6}{K(P_*)} + \ln \frac{1 + |y(P)|^2}{|y(P)|^2}.$$

A Liouville type classification result in [CGY02b] and [LL03] says that

$$v_\infty - \frac{1}{4} \ln \frac{6}{K(P_*)} = \ln |d\phi|$$

for some conformal automorphism  $\phi$  of  $\mathbb{S}^4$ , which together with (34) implies that

$$(36) \quad v_\infty \equiv \frac{1}{4} \ln \frac{6}{K(P_*)}.$$

Thus for any  $\delta > 0$ ,

$$\lim_{j \rightarrow \infty} \int_{\mathbb{S}^4 \setminus B_\delta(-P_*)} K \circ \phi_j e^{4v_j} d\text{vol}_{g_c} = 6 |\mathbb{S}^4 \setminus B_\delta(-P_*)|.$$

Together with the Gauss-Bonnet formula

$$\int_{\mathbb{S}^4} K \circ \phi_j e^{4v_j} dvol_{g_c} = 6 |\mathbb{S}^4|,$$

we have

$$\lim_{j \rightarrow \infty} \int_{B_\delta(-P_*)} K \circ \phi_j e^{4v_j} dvol_{g_c} = 6 |B_\delta(-P_*)|.$$

This allows us to apply our Theorem in [H04] on  $B_\delta(-P_*)$  for small  $\delta > 0$  to conclude that  $\exists C > 0$ , such that

$$(37) \quad \max_{\mathbb{S}^4} v_j \leq C.$$

Next we declare the

**Claim.** *There exists  $C' > 0$  such that*

$$(38) \quad \min_{\mathbb{S}^4} v_j \geq -C'.$$

The Claim can be proved making use of the information that  $R_{v_j} = R_{v_j} \circ \phi_j \geq 0$ , which implies

$$(39) \quad 2 - \Delta v_j - |\nabla v_j|^2 \geq 0.$$

Thus

$$(40) \quad \begin{aligned} v_j(P) - \bar{v}_j &= \int_{\mathbb{S}^4} (-\Delta v_j(Q)) G(P, Q) dvol_{g_c}(Q) \\ &\geq -2 \int_{\mathbb{S}^4} G(P, Q) dvol_{g_c}(Q), \end{aligned}$$

where  $G(P, Q)$  is the Green's function of  $-\Delta$  on  $\mathbb{S}^4$ . Integrating (39) over  $\mathbb{S}^4$  implies that

$$(41) \quad 2 \geq \int_{\mathbb{S}^4} |\nabla v_j|^2 \geq \text{const.} \left( \int_{\mathbb{S}^4} |v_j(P) - \bar{v}_j|^4 \right)^{\frac{1}{2}}.$$

(31), (36), (40), and (41) conclude the Claim and (9').

Next we prove the integral estimate (10). This can be seen by looking at the integral version of the equation

$$\int_{\mathbb{S}^4} [2|\nabla v_j|^2 + 2\Delta v_j - 6] \langle \nabla v_j, \nabla \eta \rangle + \Delta \eta |\nabla v_j|^2 + (K \circ \phi_j e^{4v_j} - 6) \eta = 0.$$

If we plug in  $\eta = v_j$ , we obtain

$$\int_{\mathbb{S}^4} (6 - 2|\nabla v_j|^2 - 3\Delta v_j) |\nabla v_j|^2 = \int_{\mathbb{S}^4} (K \circ \phi_j e^{4v_j} - 6) v_j.$$

Using  $\Delta v_j \leq 2 - |\nabla v_j|^2$ , we have

$$(42) \quad \int_{\mathbb{S}^4} |\nabla v_j|^4 \leq \int_{\mathbb{S}^4} (K \circ \phi_j e^{4v_j} - 6) v_j,$$

which converges to 0 by (31), (36), (37), (38) and the Dominated Convergence Theorem.  $\square$

Next, we prove **Theorem 2**. We will first prove that, under our non-degeneracy conditions on  $K$ , there is a bound  $C > 0$  depending on the quantities as in the statements of **Theorem 2**, but uniform in  $0 < s \leq 1$ , such that any solution  $w$  of (3) with  $K^{[s]}$  satisfies  $\max_{\mathbb{S}^4} w_j \leq C$ . Once we have the bound  $\max_{\mathbb{S}^4} w_j \leq C$ , the  $C^{2,\alpha}$  estimates follow from known theory of fully nonlinear elliptic equations.

*Proof of Theorem 2.* Suppose, on the contrary, that  $\max_{\mathbb{S}^4} w_j \rightarrow \infty$  (for a sequence of  $K$ 's, which we write as a single  $K$  for simplicity, satisfying the bounds in **Theorem 2**). Then, as proved above, (9') holds. Let  $P_j, t_j$  be as defined in the earlier part of the proof. We will then prove the following estimates:

$$(43) \quad |\nabla K(P_j)| = \frac{o(1)}{t_j}, \quad \text{as } j \rightarrow \infty,$$

and

$$(44) \quad \Delta K(P_j) \rightarrow 0, \quad \text{as } j \rightarrow \infty.$$

(43) and (44) would contradict our hypotheses on  $K$ .

The main idea is to examine the Kazdan-Warner identity in the light of the asymptotic profile of  $w_j$  as given in **Theorem 1**.

For each  $w_j$ , we choose stereographic coordinates with  $P_j$  as the north pole. For  $P = (x_1, \dots, x_5) \in \mathbb{S}^4$ , let its stereographic coordinates be  $y = (y_1, \dots, y_4)$ . Also set  $x' = (x_1, \dots, x_4)$ . Then

$$(45) \quad \begin{cases} x_i = \frac{2y_i}{1 + |y|^2}, & i = 1, 2, 3, 4, \\ x_5 = \frac{|y|^2 - 1}{|y|^2 + 1}. \end{cases}$$

For any  $\epsilon > 0$ , there exists  $M > 0$  such that for any  $P$  with  $|y(P)| > M$ , we have

$$(46) \quad K(P) = K(P_j) + \sum_{i=1}^4 a_i x_i + \sum_{k,h=1}^4 b_{hk} x_h x_k + r(P),$$

with

$$(47) \quad |r(P)| \leq \epsilon |x'|^2, \quad |x'| |\nabla r(P)| \leq \epsilon |x'|^2, \quad |x'|^2 |\nabla^2 r(Q)| \leq \epsilon |x'|^2.$$

We can identify  $a_i = \nabla_i K(P_j)$ ,  $b_{hk} = \nabla_{hk} K(P_j)$ , and we may assume that  $b_{hk}$  is diagonalized:  $b_{hk} = \delta_{hk} b_h$ . Then, using

$$\nabla x_1 = (1 - x_1^2, -x_1 x_2, \dots, -x_1 x_5), \dots, \nabla x_5 = (-x_5 x_1, \dots, -x_5 x_4, 1 - x_5^2),$$

and

$$\nabla x_i \cdot \nabla x_h = \delta_{ih} - x_i x_h,$$

we have, for  $1 \leq h \leq 4$ ,

$$\langle \nabla K, \nabla x_h \rangle = a_h - \sum_{i=1}^4 a_i x_i x_h + 2b_h x_h - 2 \sum_{i=1}^4 b_i x_i^2 x_h + \nabla r \cdot \nabla x_h.$$

So we can fix  $M$  large such that, when  $|y| > M$ ,

$$(48) \quad \begin{cases} \langle \nabla K, \nabla x_h \rangle = a_h + 2b_h x_h + r_1(P), \\ |r_1(P)| \leq \epsilon |x'|. \end{cases}$$

From the Kazdan-Warner identity, we have

$$\begin{aligned} 0 &= \int_{\mathbb{S}^4} \langle \nabla K^{[s]}, \nabla x_h \rangle e^{4w_j} dvol_{g_c} \\ &= s \int_{\mathbb{S}^4} \langle \nabla K, \nabla x_h \rangle e^{4w_j} dvol_{g_c} \end{aligned}$$

Thus, the deformation parameter  $s$  is divided out from the Kazdan-Warner identity to give

$$\begin{aligned} 0 &= \int_{\mathbb{S}^4} \langle \nabla K, \nabla x_h \rangle e^{4w_j} dvol_{g_c} \\ &= \int_{|y| \leq M} \langle \nabla K, \nabla x_h \rangle e^{4w_j} dvol_{g_c} + \int_{|y| > M} \langle \nabla K, \nabla x_h \rangle e^{4w_j} dvol_{g_c}. \end{aligned}$$

**Remark 5.** *It is this property that  $K$ , not  $K^{[s]}$ , can be used in the Kazdan-Warner identity that allows us to obtain bounds on  $w$  uniform in  $0 < s \leq 1$ . This also applies to the settings in [CGY93] and [L95] to make the estimates there uniform in  $0 < s \leq 1$  in the respective deformations.*

We estimate

$$\begin{aligned} \int_{|y| \leq M} \langle \nabla K, \nabla x_h \rangle e^{4w_j} dvol_{g_c} &\leq C \int_{|y| \leq M} e^{4w_j} \left( \frac{2}{1 + |y|^2} \right)^4 dy \\ &= C \int_{|z| \leq \frac{M}{t_j}} e^{4w_j} \left( \frac{2}{1 + |z|^2} \right)^4 dz \\ &\leq C \left( \frac{M}{t_j} \right)^4, \quad \text{using (37) and (38).} \end{aligned}$$

$$\int_{|y| > M} \langle \nabla K, \nabla x_h \rangle e^{4w_j} dvol_{g_c} = a_h \int_{|y| > M} e^{4w_j} dvol_{g_c} + 2b_h \int_{|y| > M} e^{4w_j} x_h dvol_{g_c} + \int_{|y| > M} e^{4w_j} r_1(P) dvol_{g_c}.$$

The following estimates will complete the proof of (43).

$$(49) \quad \lim_{j \rightarrow \infty} \int_{|y| > M} e^{4w_j} dvol_{g_c} = \frac{6}{K(P_*)} |\mathbb{S}^4|.$$

$$(50) \quad \int_{|y| > M} e^{4w_j} x_h dvol_{g_c} = \frac{o(1)}{t_j}, \quad \text{as } j \rightarrow \infty \ (1 \leq h \leq 4).$$

$$(51) \quad \int_{|y| > M} e^{4w_j} r_1(P) dvol_{g_c} = \frac{o(1)}{t_j}, \quad \text{as } j \rightarrow \infty.$$

Here are the verifications of the above estimates.

$$\int_{|y| > M} e^{4w_j} dvol_{g_c} = \int_{|z| > \frac{M}{t_j}} e^{4v_j} \left( \frac{2}{1 + |z|^2} \right)^4 dz \rightarrow \frac{6}{K(P_*)} |\mathbb{S}^4|,$$

by (31), (37) and (38).

$$\begin{aligned} \int_{|y| > M} e^{4w_j} x_h dvol_{g_c} &= \int_{|z| > \frac{M}{t_j}} \frac{2t_j z_h}{1 + t_j^2 |z|^2} e^{4v_j} \left( \frac{2}{1 + |z|^2} \right)^4 dz \\ &= \int_{|z| > \frac{M}{t_j}} \frac{2t_j z_h}{1 + t_j^2 |z|^2} \left( e^{4v_j} - \frac{6}{K(P_*)} \right) \left( \frac{2}{1 + |z|^2} \right)^4 dz \\ &= \int_{|z| > \delta} + \int_{\delta > |z| > \frac{M}{t_j}}, \end{aligned}$$

with

$$\left| \int_{\delta > |z| > \frac{M}{t_j}} \right| \leq C \int_{\delta > |z| > \frac{M}{t_j}} \frac{1}{t_j |z|} dz \leq \frac{C\delta^3}{t_j}.$$



For any given  $\epsilon > 0$ , we can first fix  $\delta > 0$  such that  $C\delta^3 < \epsilon$ . Then using the convergence of  $v_j$  to  $\frac{1}{4} \ln \frac{6}{K(P_*)}$  on  $|z| > \delta$ , we can fix  $J$  such that when  $j \geq J$ , we have  $\left| e^{4v_j} - \frac{6}{K(P_*)} \right| < \epsilon$ . Then

$$\left| \int_{|z|>\delta} \right| \leq \epsilon \int_{|z|>\delta} \frac{1}{t_j|z|} \left( \frac{2}{1+|z|^2} \right)^4 dz \leq \frac{C\epsilon}{t_j}.$$

These together prove the second estimate above. (51) follows similarly.

Finally

$$\langle \nabla K, \nabla x_5 \rangle = - \sum_{i=1}^4 a_i x_i x_5 - 2 \sum_{i=1}^4 b_i x_i^2 x_5 + \nabla r \cdot \nabla x_5.$$

We may fix  $M$  large so that  $|\nabla r \cdot \nabla x_5| \leq \epsilon |x'|^3$  when  $|y| > M$ . In

$$\begin{aligned} 0 &= \int_{\mathbb{S}^4} \langle \nabla K, \nabla x_5 \rangle e^{4w_j} dvol_{g_c} \\ (52) \quad &= \int_{|y| \leq M} \langle \nabla K, \nabla x_5 \rangle e^{4w_j} dvol_{g_c} + \int_{|y| > M} \langle \nabla K, \nabla x_5 \rangle e^{4w_j} dvol_{g_c}, \end{aligned}$$

$$(53) \quad \left| \int_{|y| \leq M} \langle \nabla K, \nabla x_5 \rangle e^{4w_j} dvol_{g_c} \right| \leq C \left( \frac{M}{t_j} \right)^4,$$

as before.

$$\left| \int_{|y| > M} x_i x_5 e^{4w_j} dvol_{g_c} \right| = \frac{o(1)}{t_j}$$

as in the proof of (50). Thus

$$(54) \quad \left| \int_{|y| > M} a_i x_i x_5 e^{4w_j} dvol_{g_c} \right| = \frac{o(1)}{t_j^2}.$$

$$\begin{aligned}
& \int_{|y|>M} x_i^2 x_5 e^{4w_j} d\text{vol}_{g_c} \\
&= \int_{|z|>\frac{M}{t_j}} \left( \frac{2t_j z_i}{1+t_j|z|^2} \right)^2 \frac{t_j|z|^2-1}{t_j|z|^2+1} e^{4v_j} \left( \frac{2}{1+|z|^2} \right)^4 dz \\
&= \int_{|z|>\frac{M}{t_j}} \left( \frac{2t_j z_i}{1+t_j|z|^2} \right)^2 \frac{t_j|z|^2-1}{t_j|z|^2+1} \left( e^{4v_j} - \frac{6}{K(P_*)} \right) \left( \frac{2}{1+|z|^2} \right)^4 dz \\
&\quad + \frac{6}{K(P_*)} \int_{|z|>\frac{M}{t_j}} \left( \frac{2t_j z_i}{1+t_j|z|^2} \right)^2 \frac{t_j|z|^2-1}{t_j|z|^2+1} \left( \frac{2}{1+|z|^2} \right)^4 dz
\end{aligned}$$

Note that

$$\begin{aligned}
& \int_{|z|>\frac{M}{t_j}} \left( \frac{2t_j z_i}{1+t_j|z|^2} \right)^2 \frac{t_j|z|^2-1}{t_j|z|^2+1} \left( \frac{2}{1+|z|^2} \right)^4 dz \\
&\asymp \frac{4}{t_j^2} \int_{|z|>\frac{M}{t_j}} \frac{z_i^2}{|z|^4} \left( \frac{2}{1+|z|^2} \right)^4 dz \\
(55) \quad &\asymp \frac{1}{t_j^2} \int_{|z|>\frac{M}{t_j}} \frac{1}{|z|^2} \left( \frac{2}{1+|z|^2} \right)^4 dz \\
&\asymp \frac{1}{t_j^2} \left( \int_0^\infty \left( \frac{2}{1+r^2} \right)^4 r dr \right) |\mathbb{S}^3|
\end{aligned}$$

Similarly, we can prove

$$(56) \quad \left| \int_{|z|>\frac{M}{t_j}} \left( \frac{2t_j z_i}{1+t_j|z|^2} \right)^2 \frac{t_j|z|^2-1}{t_j|z|^2+1} \left( e^{4v_j} - \frac{6}{K(P_*)} \right) \left( \frac{2}{1+|z|^2} \right)^4 dz \right| = \frac{o(1)}{t_j^2},$$

and

$$\begin{aligned}
 & \int_{|y|>M} |x'|^3 e^{4w_j} \\
 (57) \quad &= \int_{|z|>\frac{M}{t_j}} \left( \frac{2t_j|z|}{1+t_j^2|z|^2} \right)^3 e^{4w_j} \left( \frac{2}{1+|z|^2} \right)^4 dz \\
 &= \frac{O(1)}{t_j^3}.
 \end{aligned}$$

To put things together, we multiply (52) by  $t_j^2$  and use (53), (54), (55), and (57) to see that

$$0 = o(1) - 2\Delta K(P_j) \left( |\mathbb{S}^3| \int_0^\infty \left( \frac{2}{1+r^2} \right)^4 r dr + o(1) \right) + \frac{o(1)}{t_j},$$

which shows (44). □

*Proof of Proposition 1.* First, by **Theorem 2**, there is a  $C > 0$  depending on  $K$  and  $0 < \alpha < 1$  such that any solution  $w$  to (3) with  $K$  substituted by  $K^{[s]}$  and  $0 < s \leq 1$  satisfies

$$(58) \quad \|w\|_{C^{2,\alpha}(\mathbb{S}^4)} < C.$$

Since  $v = w \circ \varphi_{P,t} + \ln |d\varphi_{P,t}|$  is chosen such that

$$\int_{\mathbb{S}^4} e^{4v(y)} y \, d\text{vol}_{g_c} = 0,$$

we obtain, in terms of  $w$  and  $(P, t)$ ,

$$(59) \quad 0 = \int_{\mathbb{S}^4} e^{4w(x)} \varphi_{P,t}^{-1}(x) \, d\text{vol}_{g_c} = \int_{\mathbb{S}^4} e^{4w(x)} \varphi_{P,t^{-1}}(x) \, d\text{vol}_{g_c},$$

Due to (58), there is a  $\delta > 0$  such that

$$\int_{\mathbb{S}^4} e^{4w(x)} \geq \delta.$$

If there existed a sequence of solutions  $w_j$  for which  $t_j \rightarrow \infty$ , we would have, computing in stereographic coordinates in which  $P_j$  is placed at the north pole,  $\varphi_{P_j, t_j^{-1}}(x) \rightarrow$

$(0, \dots, 0, -1)$  except at  $x = P_j$ , therefore, in view of (58),

$$\int_{\mathbb{S}^4} e^{4w(x)} \varphi_{P, t^{-1}}(x) dvol_{g_c} \rightarrow (0, \dots, 0, -\int_{\mathbb{S}^4} e^{4w(x)} \neq 0,$$

contradicting (59) above. This implies the existence of some  $t_0$  such that  $t \leq t_0$ . Using this and (58) in the relation between  $w$  and  $v$ , we find an upper bound for  $\|v\|_{C^{2,\alpha}(\mathbb{S}^4)}$ . Finally using the equation for  $v$ :

$$\sigma_2(A_v) = K^{[s]} \circ \varphi_{P,t} e^{4v},$$

in which the right hand side has an upper bound in  $C^{2,\alpha}(\mathbb{S}^4)$  due to  $C^{2,\alpha}(\mathbb{S}^4)$  estimates of  $v$  and the bound  $t \leq t_0$ , we find higher derivative bounds for  $v$ . Then as  $s \rightarrow 0$ , a subsequence of  $v$  would converge to a limit  $v_\infty$  in  $C^{2,\alpha}(\mathbb{S}^4)$ , which satisfies

$$\sigma_2(A_{v_\infty}) = 6e^{4v_\infty} \quad \text{and} \quad \int_{\mathbb{S}^4} e^{4v_\infty(x)} x dvol_{g_c} = 0.$$

This implies that  $v_\infty \equiv 0$ . Since this limit  $v_\infty$  is unique, we obtain that  $v \rightarrow 0$  in  $C^{2,\alpha}(\mathbb{S}^4)$  as  $s \rightarrow 0$ , which is the remaining part of (14).  $\square$

#### 4. PROOF OF THE $W^{2,6}$ ESTIMATES OF **Theorem 1** and of **Corollary 1**

For the  $W^{2,6}$  bound for  $v_j$ , we write  $v$  for  $v_j$  and  $\sigma_2$  for  $\sigma_2(e^{-2v_j} g_c^{-1} \circ A_{v_j}) = K \circ \varphi_j$ ,

and adapt the argument for the  $W^{2,p}$  estimates in [CGY02a] of Chang-Gursky-Yang and push the argument to  $p = 6$ . We will first prove a  $W^{2,3}$  estimate for  $v_j$ , with the bound depending on an upper bound of  $\sigma_2 = K \circ \varphi_j$ , a positive lower bound for  $\sigma_2$ , and an upper

bound for  $\int_{\mathbb{S}^4} |\nabla_0(K \circ \varphi_j)|^2 dvol_{g_c}$ . Then we will extend the  $W^{2,3}$  estimate to  $W^{2,6}$  estimate for the  $v_j$  in terms of an upper bound of  $\sigma_2 = K \circ \varphi_j$ , a positive lower bound for  $\sigma_2$ , and an

upper bound for  $\int_{\mathbb{S}^4} |\nabla_0(K \circ \varphi_j)|^4 dvol_{g_c}$ . Since  $\int_{\mathbb{S}^4} |\nabla_0(K \circ \varphi_j)|^4 dvol_{g_c} = \int_{\mathbb{S}^4} |\nabla_0 K|^4 dvol_{g_c}$ , we see that a bound for the  $W^{2,3}$  norm of  $v_j$  is given in terms of an upper bound of  $K$ , a positive lower bound for  $K$ , and an upper bound for  $\int_{\mathbb{S}^4} |\nabla_0 K|^4$ . This will suffice for proving (9).

*Proof of the  $W^{2,6}$  estimates of **Theorem 1**.* First we list a few key ingredients for these  $W^{2,p}$  estimates, mostly adapted from [CGY02a]. As in [CGY02a] we explore two differential identities, which in the case of  $\mathbb{S}^4$ , are

$$(60) \quad \begin{aligned} S_{ij} \nabla_{ij}^2 R &= 6 \operatorname{tr} E^3 + R|E|^2 + 3\Delta\sigma_2 + 3(|\nabla E|^2 - \frac{|\nabla R|^2}{12}) \\ &\geq 6 \operatorname{tr} E^3 + \frac{R^3}{12} - 2\sigma_2 R + 3\Delta\sigma_2 - \frac{3|\nabla\sigma_2|^2}{2\sigma_2}, \end{aligned}$$

following (5.10) of [CGY02a], with

$$S_{ij} = \frac{\partial\sigma_2(A)}{\partial A_{ij}} = -R_{ij} + \frac{1}{2}Rg,$$

and

$$(61) \quad \begin{aligned} &S_{ij} \nabla_{ij}^2 |\nabla v|^2 \\ &= \frac{R^3}{144} - \frac{\operatorname{tr} E^3}{2} - \frac{\sigma_2 R}{12} - \frac{R|\nabla v|^4}{2} - 2S_{ij} \nabla_i |\nabla v|^2 \nabla_j v + S_{ij} \nabla_l A_{ij}^\circ \nabla_l v \\ &\quad - 2e^{-2v} S_{ij} \nabla_i v \nabla_j v + 2Re^{-2v} |\nabla v|^2 + \frac{Re^{-4v}}{2} - \langle \nabla v, \nabla \sigma_2 \rangle - 2\sigma_2 e^{-2v}, \end{aligned}$$

following (5.44) of [CGY02a] and the fact that  $A_{ij}^0 = g_{ij}^0$  in the case of  $\mathbb{S}^4$ . Here the differentiations are in the metric  $g$ .

In (60) and (61) we used  $|E|^2 = \frac{R^2}{12} - 2\sigma_2$  and

$$(62) \quad |\nabla E|^2 - \frac{|\nabla R|^2}{12} \geq -\frac{|\nabla \sigma_2|^2}{2\sigma_2}.$$

(62) can be proven as in (7.26) of [CGY02a], but can also be seen to be based on the general fact that  $\{\sigma_k\}^{1/k}$  is concave in its argument as follows: set  $F(A_{ij}) = \{\sigma_k(A_{ij})\}^{1/k}$ , then

$$(63) \quad S_{ij} = \frac{\partial \sigma_k}{\partial A_{ij}} = kF^{k-1} \frac{\partial F}{\partial A_{ij}}, \quad \text{and} \quad \nabla \sigma_k = S_{ij} \nabla A_{ij} = kF^{k-1} \frac{\partial F}{\partial A_{ij}} \nabla A_{ij}.$$

So

$$\nabla_l S_{ij} = kF^{k-1} \frac{\partial^2 F}{\partial A_{ij} \partial A_{IJ}} \nabla_l A_{IJ} + k(k-1)F^{k-2} \frac{\partial F}{\partial A_{ij}} \frac{\partial F}{\partial A_{IJ}} \nabla_l A_{IJ}.$$

Thus

$$(64) \quad \begin{aligned} & \sum_l \nabla_l S_{ij} \nabla_l A_{ij} \\ &= kF^{k-1} \frac{\partial^2 F}{\partial A_{ij} \partial A_{IJ}} \nabla_l A_{IJ} \nabla_l A_{ij} + k(k-1)F^{k-2} \frac{\partial F}{\partial A_{ij}} \frac{\partial F}{\partial A_{IJ}} \nabla_l A_{IJ} \nabla_l A_{ij} \\ &\leq \frac{(k-1)|\nabla \sigma_k|^2}{k\sigma_k} \quad \text{using concavity of } F \text{ and (63)}. \end{aligned}$$

In the case of  $2k = n = 4$ ,  $A_{ij} = E_{ij} + \frac{R}{12}g_{ij}$ , and  $S_{ij} = \frac{R}{4}g_{ij} - E_{ij}$ . So

$$\sum_l \nabla_l S_{ij} \nabla_l A_{ij} = \sum_l \left\{ \frac{\nabla_l R}{4} g_{ij} - \nabla_l E_{ij} \right\} \left\{ \nabla_l E_{ij} + \frac{\nabla_l R}{12} g_{ij} \right\} = \frac{|\nabla R|^2}{12} - |\nabla E|^2,$$

and by (64)

$$\frac{|\nabla R|^2}{12} - |\nabla E|^2 \leq \frac{|\nabla \sigma_2|^2}{2\sigma_2}.$$

Because of  $S_{ij,j} = 0$ , which is a consequence of Bianchi identity, we can use (60) and (61) to obtain

$$\begin{aligned} 0 &= \int_{\mathbb{S}^4} S_{ij} \nabla_{ij}^2 (R + 12|\nabla v|^2) \\ &\geq \int_{\mathbb{S}^4} \frac{R^3}{6} - 6R|\nabla v|^4 - 24S_{ij} \nabla_i |\nabla v|^2 \nabla_j v \\ &\quad + 12S_{ij} \nabla_l A_{ij}^\circ \nabla_l v + 24Re^{-2v} |\nabla v|^2 - 24e^{-2v} S_{ij} \nabla_i v \nabla_j v \\ &\quad - 12\langle \nabla v, \nabla \sigma_2 \rangle + (6e^{-4v} - 2\sigma_2)R - 24e^{-2v} \sigma_2 - \frac{3|\nabla \sigma_2|^2}{2\sigma_2}, \end{aligned}$$

from which we can estimate  $\int_{\mathbb{S}^4} R^3$  in terms of the other terms:

$$\begin{aligned} \int_{\mathbb{S}^4} \frac{R^3}{6} &\leq \int_{\mathbb{S}^4} 6R|\nabla v|^4 + 24S_{ij} \nabla_i |\nabla v|^2 \nabla_j v \\ (65) \quad &\quad - 12S_{ij} \nabla_l A_{ij}^\circ \nabla_l v - 24Re^{-2v} |\nabla v|^2 + 24e^{-2v} S_{ij} \nabla_i v \nabla_j v \\ &\quad + 12\langle \nabla v, \nabla \sigma_2 \rangle - (6e^{-4v} - 2\sigma_2)R + 24\sigma_2 e^{-2v} + \frac{3|\nabla \sigma_2|^2}{2\sigma_2}. \end{aligned}$$

The integrations are done in the  $g$  metric, but due to the  $L^\infty$  estimates on  $v$ , the integrals in  $g$  metric are comparable to those in  $g_c$ . The terms that require careful treatments are

$$\int_{\mathbb{S}^4} S_{ij} \nabla_i |\nabla v|^2 \nabla_j v$$

and

$$(66) \quad \int_{\mathbb{S}^4} R|\nabla v|^4 \leq \left[ \int_{\mathbb{S}^4} R^3 \right]^{1/3} \left[ \int_{\mathbb{S}^4} |\nabla v|^6 \right]^{2/3} \leq \frac{\epsilon}{3} \int_{\mathbb{S}^4} R^3 + \frac{2\epsilon^{-1/2}}{3} \int_{\mathbb{S}^4} |\nabla v|^6.$$

The term  $\int_{\mathbb{S}^4} S_{ij} \nabla_i |\nabla v|^2 \nabla_j v$  can be estimated as (5.53) in [CGY02a]

$$(67) \quad \begin{aligned} & \int_{\mathbb{S}^4} S_{ij} \nabla_i |\nabla v|^2 \nabla_j v \\ &= - \int_{\mathbb{S}^4} |\nabla v|^2 S_{ij} \nabla_{ij}^2 v \\ &= - \int_{\mathbb{S}^4} |\nabla v|^2 S_{ij} \left\{ -\frac{A_{ij}}{2} + \frac{A_{ij}^0}{2} - \nabla_i v \nabla_j v + \frac{|\nabla v|^2}{2} g_{ij} \right\} \\ &= \int_{\mathbb{S}^4} |\nabla v|^2 \left\{ \sigma_2 + S_{ij} \nabla_i v \nabla_j v - \frac{R|\nabla v|^2}{2} - \frac{S_{ij} A_{ij}^0}{2} \right\} \\ &= \int_{\mathbb{S}^4} |\nabla v|^2 \left\{ \sigma_2 - R_{ij} \nabla_i v \nabla_j v - \frac{S_{ij} A_{ij}^0}{2} \right\} \\ &\leq \int_{\mathbb{S}^4} |\nabla v|^2 \sigma_2. \end{aligned}$$

where in the last line we used  $(R_{ij}) \geq 0$  when  $g \in \Gamma_2^+$  in dimension 4 and  $S_{ij} A_{ij}^0 \geq 0$  on

$\mathbb{S}^4$ . The terms in the second line of (65) can be estimated in terms of  $\int_{\mathbb{S}^4} R e^{-2v} |\nabla v|^2$ , which in turn can be estimated as

$$(68) \quad \int_{\mathbb{S}^4} R e^{-2v} |\nabla v|^2 \lesssim \left\{ \int_{\mathbb{S}^4} R^3 \right\}^{1/3} \left\{ \int_{\mathbb{S}^4} |\nabla v|^3 \right\}^{2/3} \leq \frac{\epsilon}{3} \int_{\mathbb{S}^4} R^3 + \frac{2\epsilon^{-1/2}}{3} \int_{\mathbb{S}^4} |\nabla v|^3.$$

The terms in the last line of (65) can be estimated in terms of upper bound of  $\sigma_2$ , a lower bound of  $\sigma_2$ , and  $\int_{\mathbb{S}^4} |\nabla \sigma_2|^2$ , in a trivial way. The term  $\int_{\mathbb{S}^4} |\nabla v|^6$  in (66) can be



estimated as

$$(69) \quad \int_{\mathbb{S}^4} |\nabla v|^6 \leq \left[ \int_{\mathbb{S}^4} |\nabla v|^4 \right]^{3/4} \left[ \int_{\mathbb{S}^4} |\nabla v|^{12} \right]^{1/4},$$

and, as in (5.73) in [CGY02a],

$$(70) \quad \begin{aligned} & \left[ \int_{\mathbb{S}^4} |\nabla v|^{12} \right]^{1/4} \\ & \lesssim \int_{\mathbb{S}^4} |\nabla^2 v|^3 + |\nabla v|^6 + e^{-3v} |\nabla v|^3 \\ & \lesssim \int_{\mathbb{S}^4} R^3 + |\nabla v|^6 + 1, \end{aligned}$$

here in the last line we used

$$(71) \quad S_{ij} = S_{ij}^0 + 2\nabla_{ij}^2 v - 2(\Delta v)g_{ij} + 2\nabla_i v \nabla_j v + |\nabla v|^2 g_{ij},$$

$$(72) \quad R = R_0 e^{-2v} - 6\Delta v + 6|\nabla v|^2,$$

$$(73)$$

and

$$0 \leq (S_{ij}) \leq (Rg_{ij}).$$

Using (70) in (69) and noting that  $\int_{\mathbb{S}^4} |\nabla v|^4$  is small, we obtain

$$(74) \quad \int_{\mathbb{S}^4} |\nabla v|^6 \lesssim \left[ \int_{\mathbb{S}^4} |\nabla v|^4 \right]^{3/4} \int_{\mathbb{S}^4} R^3 + 1,$$

Using (74), together with (67), (68) and (66) in (65), and noting the smallness of  $\int_{\mathbb{S}^4} |\nabla v|^4$ , we obtain an upper bound for  $\int R^3$  in terms of  $\int |\nabla \sigma_2|^2$ , upper bound for  $\sigma_2$  and

positive lower bound for  $\sigma_2$ . Note that  $\int_{\mathbb{S}^4} |\nabla \sigma_2|^4 = \int_{\mathbb{S}^4} |\nabla_0(K \circ \phi_j)|^4 = \int_{\mathbb{S}^4} |\nabla_0 K|^4 dvol_{g_c}$

and using a transformation law like (74), we can estimate

$$\int |\Delta_0 v|^3 dvol_{g_c} \lesssim \int (R^3 + |\nabla_0 v|^6) dvol_{g_c} \lesssim \int R^3 + 1,$$

bounded above in terms of  $\int |\nabla_0 K|^4 dvol_{g_c}$ , upper bound for  $K$  and positive lower bound

for  $K$ . Then we can use the  $W^{2,p}$  theory for the Laplace operator to obtain the full  $W^{2,3}$  estimates for  $v$ .

**Remark 6.** *In fact, for any solution  $w$  to (3), one can obtain an upper bound for the  $W^{2,3}$  norm of  $w$  in terms of a positive upper and lower bound for  $K$ , an upper bound for  $\int |\nabla_0 K|^2 dvol_{g_c}$ , and an upper bound for  $|w|$  and  $\int |\nabla_0 w|^4 dvol_{g_c}$ . A proof would*

*proceed as above, instead of using the smallness of  $\int |\nabla_0 w|^4 dvol_{g_c}$  in proving (74) and*

*the subsequent bound on  $\int R^3$  via (65), one uses Proposition 5.20, Proposition 5.22, and Lemma 5.24 in [CGY02a] to complete the argument.*

To obtain the  $W^{2,6}$  estimates of  $v$  by iteration, we multiply (60) and (61) by  $R^p$  and estimate  $\int R^{p+3}$  in terms of the other terms:

$$\int R^p S_{ij} \nabla_{ij}^2 R \geq \int \frac{R^{p+3}}{12} + 6R^p \operatorname{tr} E^3 - 2\sigma_2 R^{p+1} + 3R^p \Delta \sigma_2 - \frac{3R^p |\nabla \sigma_2|^2}{2\sigma_2},$$

and

$$\begin{aligned} & \int R^p S_{ij} \nabla_{ij}^2 |\nabla v|^2 \\ &= \int \frac{R^{p+3}}{144} - \frac{R^p \operatorname{tr} E^3}{2} - \frac{\sigma_2 R^{p+1}}{12} - \frac{R^{p+1} |\nabla v|^4}{2} - 2R^p S_{ij} \nabla_i |\nabla v|^2 \nabla_j v \\ & \quad + R^p S_{ij} \nabla_l A_{ij}^\circ \nabla_l v - 2R^p S_{ij} \nabla_i v \nabla_j v + 2R^{p+1} |\nabla v|^2 - R^p \langle \nabla v, \nabla \sigma_2 \rangle - 2\sigma_2 R^p + \frac{R^{p+1}}{2}. \end{aligned}$$

From these we obtain

$$\begin{aligned}
& \int \frac{R^{p+3}}{6} \\
(75) \quad & \leq \int R^p S_{ij} \nabla_{ij}^2 \{R + 12|\nabla v|^2\} - 3R^p \Delta \sigma_2 + 6R^{p+1} |\nabla v|^4 + 24R^p S_{ij} \nabla_i |\nabla v|^2 \nabla_j v \\
& + 3\sigma_2 R^{p+1} + \frac{3R^p |\nabla \sigma_2|^2}{2\sigma_2} - 12R^p S_{ij} \nabla_l A_{ij}^\circ \nabla_l v + 24R^p S_{ij} \nabla_i v \nabla_j v - 24R^{p+1} |\nabla v|^2 \\
& + 12R^p \langle \nabla v, \nabla \sigma_2 \rangle + 24\sigma_2 R^p - 6R^{p+1}.
\end{aligned}$$

The most crucial terms are

$$\begin{aligned}
(76) \quad & \int R^p S_{ij} \nabla_{ij}^2 R = -p \int R^{p-1} S_{ij} \nabla_i R \nabla_j R, \\
& \int R^p S_{ij} \nabla_{ij}^2 |\nabla v|^2 \\
& = -p \int R^{p-1} S_{ij} \nabla_i R \nabla_j |\nabla v|^2 \\
& \leq p \int R^{p-1} [S_{ij} \nabla_i R \nabla_j R]^{1/2} [S_{ij} \nabla_i |\nabla v|^2 \nabla_j |\nabla v|^2]^{1/2} \\
(77) \quad & \leq p \left[ \int R^{p-1} S_{ij} \nabla_i R \nabla_j R \right]^{1/2} \left[ \int R^{p-1} S_{ij} \nabla_i |\nabla v|^2 \nabla_j |\nabla v|^2 \right]^{1/2} \\
& \leq \frac{p}{2} \int R^{p-1} S_{ij} \nabla_i R \nabla_j R + 2p \int R^p |\nabla^2 v|^2 |\nabla v|^2 \\
& \leq \frac{p}{2} \int R^{p-1} S_{ij} \nabla_i R \nabla_j R + C p \int R^p (R^2 + |S_{ij}^0|^2 + |\nabla v|^2) |\nabla v|^2.
\end{aligned}$$

and

$$\begin{aligned}
& \int -R^p \Delta \sigma_2 \\
&= p \int R^{p-1} \nabla \sigma_2 \nabla R \\
(78) \quad & \leq p \left[ \int |\nabla \sigma_2|^4 \right]^{1/4} \left[ \int |\nabla R|^2 R^{p-2} \right]^{1/2} \left[ \int R^{2p} \right]^{1/4} \\
& \leq p \left[ \int |\nabla \sigma_2|^4 \right]^{1/4} \left[ \int \frac{R^{p-1} S_{ij} \nabla_i R \nabla_j R}{3\sigma_2} \right]^{1/2} \left[ \int R^{2p} \right]^{1/4} \\
& \leq \frac{\epsilon p}{2} \int \frac{R^{p-1} S_{ij} \nabla_i R \nabla_j R}{3\sigma_2} + \frac{p}{2\epsilon} \left[ \int |\nabla \sigma_2|^4 \right]^{1/2} \left[ \int R^{2p} \right]^{1/2}
\end{aligned}$$

Next we claim that the Sobolev inequality in dimension 4 implies

$$(79) \quad \left[ \int R^{2p} \right]^{1/2} \lesssim p^2 \int |\nabla R|^2 R^{p-2} + \int R^p |\nabla v|^2 + \int R^p e^{-2v}.$$

Using (79) in (78), we obtain

$$\begin{aligned}
& \int -R^p \Delta \sigma_2 \\
(80) \quad & \leq \epsilon p \int \frac{R^{p-1} S_{ij} \nabla_i R \nabla_j R}{3\sigma_2} + \frac{\epsilon}{2p} \int R^p |\nabla v|^2 + \frac{p^3}{8\epsilon^3} \int |\nabla \sigma_2|^4.
\end{aligned}$$

Using (76), (77), and (80) in (75) and choosing  $\epsilon > 0$  small, we obtain

$$\begin{aligned}
(81) \quad & \int \frac{R^{p+3}}{6} + \frac{p}{4} R^{p-1} S_{ij} \nabla_i R \nabla_j R \\
& \lesssim \int R^{p+1} |\nabla v|^4 + R^{p+2} |\nabla v|^2 + R^p |\nabla \sigma_2|^2 + R^p |\nabla v| |\nabla \sigma_2| + R^{p+1} + 1 \\
& \lesssim \left\{ \int R^{p+3} \right\}^{\frac{p+1}{p+3}} \left\{ \int |\nabla v|^{2(p+3)} \right\}^{\frac{2}{p+3}} + \left\{ \int R^{p+3} \right\}^{\frac{p+2}{p+3}} \left\{ \int |\nabla v|^{2(p+3)} \right\}^{\frac{1}{p+3}} \\
& \quad + \left\{ \int R^{2p} \right\}^{1/2} \left\{ \int |\nabla \sigma_2|^4 \right\}^{1/2} + \left\{ \int R^{2p} \right\}^{1/2} \left\{ \int |\nabla v|^4 \right\}^{1/4} \left\{ \int |\nabla \sigma_2|^4 \right\}^{1/4} + \int R^{p+1} + 1
\end{aligned}$$

Now for  $p \leq 3$ , we have  $2p \leq p+3$  and  $2(p+3) \leq 12$ . Using the earlier bounds on  $\int R^3$

and  $\int |\nabla v|^{12}$  from (70), we obtain an upper bound for  $\int R^6$  in terms of  $\int |\nabla_0 K|^4 dvol_{g_c}$ , an upper bound and a positive lower bound of  $K$ , which again gives a bound for  $v$  in  $W^{2,6}$ .  $\square$

*Proof of Corollary 1.* Let  $\delta > 0$  be small such that the argument for (74) and the subsequent  $W^{2,3}$  estimate for  $v$  via (65) would go through when  $\int_{\mathbb{S}^4} |\nabla v|^4 \leq \delta$ . For any

admissible solution  $w$  to (3), Theorem 1 implies that there is a constant  $B > 0$  depending on the  $C^2$  norm of  $K$ , a positive lower bound of  $K$ , and  $\delta > 0$ , such that if  $\max w = w(Q) > B$ , then the normalized  $v$  defined as in Theorem 1:  $v = w \circ \varphi + \ln |d\varphi|$  with  $v(Q) = \frac{1}{4} \ln \frac{6}{K(Q)}$ , would satisfy

$$(82) \quad \left| v - \frac{1}{4} \ln \frac{6}{K(Q)} \right| \leq \delta, \quad \text{and} \quad \int_{\mathbb{S}^4} |\nabla v|^4 \leq \delta.$$

$v$  also satisfies (8) and then estimate (65), with  $\sigma_2$  standing for  $K \circ \varphi$ , is valid for  $v$ . Then the  $W^{2,3}$  estimate in Theorem 1 would be valid for  $v$ , and one obtains a bound for the  $W^{2,3}$  norm of  $v$  in terms of an upper bound for  $K$ , a positive lower bound for  $K$ , and an upper bound for  $\int_{\mathbb{S}^4} |\nabla_0 K \circ \varphi|^4 dvol_{g_c}$ , and since  $\int_{\mathbb{S}^4} |\nabla_0 K \circ \varphi|^4 dvol_{g_c} = \int_{\mathbb{S}^4} |\nabla_0 K|^4 dvol_{g_c}$ , one

can use this estimate to obtain the bound for  $F[v]$ . Since  $II[w] = II[v]$  and  $Y[w] = Y[v]$ , the bound for  $F[w]$  now follows. When the solution  $w$  satisfies  $w \leq B$ , then one can use the Harnack type estimate in [H04] to obtain a lower bound for  $w$ , and use inequality

(42) to obtain an upper bound for  $\int_{S^4} |\nabla w|^4 dvol_{g_c}$ . Then one can use Remark 6 to obtain

the  $W^{2,3}$  estimate for  $w$  in terms of an upper bound for  $K$ , a positive lower bound for  $K$ , and an upper bound for  $\int_{S^4} |\nabla_0 K|^2 dvol_{g_c}$ . Finally these  $W^{2,3}$  estimates for  $w$  give

directly the bound for  $F[w]$ . □

## REFERENCES

- [A79] T. Aubin, *Meilleures constantes dans le thorme d'inclusion de Sobolev et un thorme de Fredholm non linnaire pour la transformation conforme de la courbure scalaire*, (French. English summary) J. Funct. Anal. **32** (1979), no. 2, 148–174.
- [BC91] A. Bahri and J.-M. Coron, *The scalar-curvature problem on the standard three-dimensional sphere*, J. Funct. Anal. **95** (1991), no. 1, 106–172.
- [CL93] Kung Ching Chang and Jia Quan Liu, *On Nirenberg's problem*, Internat. J. Math. **4** (1993), no. 1, 35–58.
- [CC07] S.-Y. A. Chang and S. Sophie Chen, *On a fully non-linear PDE in conformal geometry*, Mat. Ense. Univ. (N. S.) **15** (2007), suppl. 1, 17–36.
- [CY87] S.-Y. A. Chang and P. Yang, *Prescribing Gaussian curvature on  $S^2$* , Acta Math. **59** (1987), no. 3-4, 215–259.
- [CY91] S.-Y. A. Chang and P. Yang, *A perturbation result in prescribing scalar curvature on  $S^n$* , Duke Math. J. **64** (1991), no. 1, 27–69.
- [CY93] S.-Y. A. Chang and P. Yang, *Addendum to: "A perturbation result in prescribing scalar curvature on  $S^n$ " [Duke Math. J. 64 (1991), no. 1, 27–69; MR1131392 (92m:53063)]*, Duke Math. J. **71** (1993), no. 1, 333–335.
- [CY02] S.-Y. A. Chang and Paul Yang, *Non-linear Partial Differential equations in Conformal Geometry*, Proceedings for ICM 2002, Beijing, volume I, pp 189-209.
- [CGY02a] S.-Y. A. Chang, M. Gursky and P. Yang, *An equation of Monge-Ampère type in conformal geometry, and four-manifolds of positive Ricci curvature*, Annals of Math., **155**(2002), 709-787.
- [CGY02b] S.-Y. A. Chang, M. Gursky and P. Yang, *An a priori estimate for a fully nonlinear equation on four-manifolds*, Dedicated to the memory of Thomas H. Wolff. J. Anal. Math. **87** (2002), 151–186.
- [CGY03] S.-Y. A. Chang, M. Gursky and P. Yang, *Entire solutions of a fully nonlinear equation*, "Lectures in Partial Differential Equations in honor of Louis Nirenberg's 75th birthday", chapter 3. International Press, 2003.
- [CGY93] S.-Y. A. Chang, M. Gursky and P. Yang, *The scalar curvature equation on 2- and 3-spheres*, Calc. Var. Partial Differential Equations **1** (1993), no. 2, 205–229.
- [CHY04] S.-Y. A. Chang, Z. Han, and P. Yang, *Apriori estimates for solutions of the prescribed  $\sigma_2$  curvature equation on  $S^4$* , manuscript, 2004.
- [CHY05] S.-Y. A. Chang, Z. Han, and P. Yang, *Classification of singular radial solutions of the  $\sigma_k$ -Yamabe equation on annular domains in  $\mathbb{R}^n$* , J. Differential Equations **216** (2005), no. 2, 482–501.
- [CL99] C.C. Chen and C.S. Lin, *Blowing up with infinite energy of conformal metrics on  $S^n$* , Comm. PDE **24** (1999), 785–799.
- [CD87] W.-X. Chen and W.-Y. Ding, *Scalar curvatures on  $S^2$* , Trans. Amer. Math. Soc. **303** (1987), no. 1, 365–382.
- [GW03] P. Guan and G. Wang, *Local estimates for a class of fully nonlinear equations arising from conformal geometry*, International Mathematics Research Notices, V. 2003, Issue 26(2003), 1413-1432.
- [H90] Z.-C. Han, *Prescribing Gaussian curvature on  $S^2$* , Duke Math. J. **61**(1990), 679-703.

- [H04] Z.-C. Han, *Local pointwise estimates for solutions of the  $\sigma_2$  curvature equation on 4-manifolds*, Int. Math. Res. Not. 2004, no. 79, 4269–4292.
- [H06] Z.-C. Han, *A Kazdan-Warner type identity for the  $\sigma_k$  curvature*, C. R. Math. Acad. Sci. Paris **342** (2006), no. 7, 475–478.
- [J04] Min Ji, *On positive scalar curvature on  $S^2$* , Calc. Var. Partial Differential Equations **19** (2004), no. 2, 165–182.
- [K85] J. Kazdan, *Prescribing the curvature of a Riemannian manifold*, CBMS Regional Conference Series in Mathematics, 57. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1985. vii+55 pp.
- [KMS09] M. A. Khuri, F. C. Marques, and R. M. Schoen, *A compactness theorem for the Yamabe problem*, J. Diff. Geometry **81** (2009), 143–196.
- [LP87] J. Lee and T. Parker, *The Yamabe problem*, Bull. Amer. Math. Soc. **17** (1987) 37–91.
- [LL03] Aobing Li and YanYan Li, *On some conformally invariant fully nonlinear equations*, Comm. Pure Appl. Math. **56** (2003), 1414–1464.
- [LL05] Aobing Li and YanYan Li, *On some conformally invariant fully nonlinear equations, II: Liouville, Harnack and Yamabe*, Acta Math. **195** (2005), 117–154.
- [L89] YanYan Li, *Degree theory for second order nonlinear elliptic operators and its applications*, Comm. Partial Differential Equations **14** (1989), no. 11, 1541–1578.
- [L95] YanYan Li, *Prescribing scalar curvature on  $S^n$  and related problems. I*, J. Differential Equations **120** (1995), no. 2, 319–410.
- [L96] YanYan Li, *Prescribing scalar curvature on  $S^n$  and related problems, Part II: Existence and Compactness*, Comm. Pure Appl. Math **49** (1996), 541–597.
- [L98] YanYan Li, *Fine analysis of blow up and applications*, First International Congress of Chinese Mathematicians (Beijing, 1998), 411–421, AMS/IP Stud. Adv. Math., 20, Amer. Math. Soc., Providence, RI, 2001.
- [O82] E. Onofri, *On the positivity of the effective action in a theory of random surfaces*, Comm. Math. Phys. **86** (1982), no. 3, 321–326.
- [S91] R. M. Schoen, *A report on some recent progress on nonlinear problems in geometry*, Surveys in Differential Geometry **1** (1991) 201–241.
- [V00a] J. Viaclovsky, *Conformal geometry, contact geometry, and the calculus of variations*, Duke Math. J. **101**, No. 2 (2000), 283–316.
- [V00b] J. Viaclovsky, *Some fully nonlinear equations in conformal geometry*, Differential equations and mathematical physics (Birmingham, AL, 1999), 425–433, AMS/IP Stud. Adv. Math., 16, Amer. Math. Soc., Providence, RI, 2000.
- [V02] J. Viaclovsky, *Estimates and existence results for some fully nonlinear elliptic equations on Riemannian manifolds*, Communications in Analysis and Geometry **10** (2002), no.4, 815–846.
- [V06] J. Viaclovsky, *CONFORMAL GEOMETRY AND FULLY NONLINEAR EQUATIONS*, Inspired by S. S. Chern, 435–460, Nankai Tracts Math., 11, World Sci. Publ., Hackensack, NJ, 2006.
- [Z90] D. Zhang, *New results on geometric variational problems*, Thesis, Stanford University, 1990.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08540

*E-mail address:* chang@math.princeton.edu

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854

*E-mail address:* zchan@math.rutgers.edu

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08540

*E-mail address:* yang@math.princeton.edu